ISSN 0300-9246

JOURNAL OF THE CHEMICAL SOCIETY

Dalton Transactions

A journal of inorganic chemistry

1985

ROYAL SOCIETY OF CHEMISTRY

© The Royal Society of Chemistry, 1985. All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form, or by any means, electronic, mechanical, photographic, recording, or otherwise, without the prior permission of the publishers.

Printed in Great Britain by
Richard Clay (The Chaucer Press) Ltd.,
Bungay, Suffolk

AUTHOR INDEX

Abdullah, Mohammed, 2085
Abel, Edward W., 345, 1561, 1569, 2195
Abrams, Gillian, 2203
Agarwala, Umesh C., 2449
Aggarwal, Aneel, 795
Ahmad, Naseer, 2547
Ahmeti, Xhavit, 1095
Aime, S., 225
Ainscough, Eric W., 151
Airoldi, Claudio, 369, 1103
Akid, Robert, 395
Akitt, J. W., 591
Al-Baker, Salam, 1387, 2655
Al-Hamoud, Saud A. Aziz, 749
Al-Salih, Talib I., 1255
Al-Wassil, Abdul-Aziz I., 1929
Albano, Vincenzo G., 35, 1137, 1309
Alberts, Albert H., 2311
Alcock, Nathaniel W., 219, 517, 1001, 1361, 1997, 2683
Ali, Hapipah Mohd., 1131
Allen, David W., 2505
Allen, Stephen R., 435
Alonso, José Antonio, 1633, 2225
Alonzo, Giuseppe, 523
Alt, Helmut G., 1365
Anderson, Susan, 2247
Andrews, Steven J., 1645
Andriollo, Antida, 1859
Anhaus, Jens, 2453
Antal, Károly, 1191
Aoyagi, Kimitake, 1733
Araki, Koji, 373
Arce, Alejo J., 2479
Arean, Carlos Otero, 2155
Aroney, Manuel J., 1761
Asperger, Smiljko, 1095
Asakawa, Miaki, 101, 107
Ascenzi, Paolo, 1107, 1113
Ashmawy, Fathy M., 1391
Ashok, Rajan F. N., 2449
Aspinall, Helen C., 743
Astheimer, Harald, 315
Atassi, Ghanem, 523
Attali, Serge, 2521
Audisio, Guido, 1547
Awang, M. Rasol, 2009
Bányai, István, 1191
Baggott, James E., 2247
Bailey, Neil A., 1471
Bailey, Stuart I., 1747
Baiocchi, Claudio, 2615
Baird, Gordon J., 1479
Baird, Michael C., 2691
Bajaj, Hari C., 2603
Bakhmutov, Vladimir I., 17
Baldwin, David A., 1613
Bales, John R., 795
Balt, Sijbe, 659, 1495, 2223
Baños, Joaquin, 1975
Bandy, Judith A., 2025, 2037
Bandyopadhyay, Debkumar, 159
Banford, Jon, 1355
Baral, Subhash, 2213
Barbier, Jean-Pierre, 597
Barbieri, Renato, 523
Bard, Allen J., 1303
Barr, Robert D., 2009
Barratt, David S., 135, 2661
Barrett, Jack, 2085
Basato, Marino, 91
Bashkin, Jim, 423
Bataille, Michel, 535, 1201
Batchelor, Raymond J., 1727
Batley, Graeme E., 1967
Bats, Jan W., 1, 2453
Bau, Robert, 2051
Baumann, Nikolaus, 223
Baxter, Paul L., 807, 941
Beck, Mihály T., 1191, 1669
Beckett, Michael A., 1119
Beevor, Robert G., 435
Begley, Michael J., 2433
Belikov, Vasili M., 17
Belokon', Yuri N., 17
Bender, Robert, 711
Benson, Christopher G., 2661
Berardelli, Maria L., 1737
Beringhelli, Tiziana, 1507, 1899
Berry, Frank J., 451
Bethell, Donald, 1405
Betterton, Eric A., 1613, 1619
Bhargava, Suresh K., 345, 1561
Bhattacharjee, Manabendra N., 409
Bigoli, Francesco, 1349
Birchall, Thomas, 1275, 1727, 2671
Blagg, Adrian, 1213
Blair, Peter D., 755
Blake, Antony B., 2509
Blandamer, Michael J., 867
Blaschke, Günther, 827
Blonk, Henk L., 1699
Blower, Philip J., 1533, 2305, 2639, 2647
Boddington, Terrence, 1213
Boden, Peter J., 2551
Boeyens, Jan C. A., 2277
Borish, Edward T., 1789
Botsivali, Maria, 1147
Bottomley, Frank, 1435, 2427
Bould, Jonathan, 1843
Bousseau, Michèle, 783
Bouwman, Elisabeth, 737
Bowman, Karen, 857
Boyar, Esther B., 617, 621, 629, 2113
Braga, Dario, 35, 1137, 1309, 1795
Brant, Patrick, 269
Braterman, P. S., 1081
Braunstein, Pierre, 711
Briant, Clive E., 851, 1693, 2343
Briggs, T. Nigel, 1249
Brint, Paul, 1915, 2591
Brisdon, Brian J., 2191
Brničević, Nevenka, 455, 1531
Broadley, Karen, 777
Brodie, Andrew M., 151
Brookhart, Maurice, 423
Brown, David, 1001, 1853
Brown, David B., 2319
Brown, Michael P., 2421
Brown, Paul L., 723, 1967
Bruce, Michael I., 1223, 1229
Brunori, Maurizio, 1107, 1113
Bruque, Sebastián, 213
Bünzli, Jean-Claude G., 885
Burgess, John, 867
Burgess, Kevin, 85
Burgos, Francisco Sánchez, 31
Byers, Peter K., 981, 1183
Cabeza, Javier A., 573
Calderazzo, Fausto, 1989
Calvaruso, Giuseppe, 1683
Camellini, Marisa Tiripicchio, 1487
Cameron, Alan D., 2691
Cameron, T. Stanley, 345
Campbell, Nicholas J., 1673
Canovese, Luciano, 27, 731
Canty, Allan J., 981, 1183
Carbone, Antonio I., 1683
Cardin, Christine J., 1527
Carl, Lothar, 523
Carlsen, Lars, 879
Carmona-Guzmán, M. Carmen, 1975
Carmona, Daniel, 973
Carr, Stuart W., 1213, 2131
Carriedo, Gabino A., 905
Carvalho, N. N., 2079
Cassoux, Patrick, 783
Castellani, Manola, 1989
Castro, Alicia, 2225
Cattalini, Lucio, 27, 731, 2091
Cavasino, F. Paolo, 1683
Cayuelas, José A., 1163
Cenini, Sergio, 163
Chagas, Aécio P., 1103
Chakravorty, Animesh, 159, 361
Chatt, Joseph, 1131
Chaudhuri, Mihir K., 409
Chaudret, Bruno, 43
Chauhan, Virendra B. S., 2449
Cheeseman, Paul G., 2551
Chemaly, Susan M., 1613, 1619
Chen, Grace S. H., 9
Chessa, Gavino, 2091
Chicote, Maria T., 1163
Chiesa, Anna, 1971
Chini, Paolo, 35
Chung, Chung-Sun, 2217
Ciampolini, Mario, 479, 1179, 1425
Ciani, Gianfranco, 1507
Ciriano, Miguel A., 1891
Clark, Robin J. H., 579, 815, 1775
Clayden, Nigel J., 1811
Clegg, William, 565, 1977, 2161
Cloke, F. Geoffrey N., 423
Clucas, Jennifer A., 1835
Coates, John H., 413
Coffman, Robert E., 891
Cohen, Haim, 641
Cole-Hamilton, David J., 387
Colquhoun, Howard M., 761, 1249
Commenges, Gérard, 1087
Connelly, Neil G., 699, 777, 1019, 1027, 2283
Constable, Edwin C., 333, 1719, 2247, 2687
Cook, Michael J., 1781
Coombe, Vyvyan T., 947
Cooper, Gary R., 1213
Cooper, Mervyn K., 1761
Coronas, Juan Maria, 2333
Coville, Neil J., 2277
Cowley, Alan H., 383, 1303
Cox, David N., 2343
Cox, Kevin, 423
Cradock, Stephen, 755
Crisponi, Guido, 1349
Crocker, Mark, 2145
Crook, Janet E., 2407
Crotti, Corrado, 163
Croud, Vincent B., 815
Cummins, Diane, 493
Curtis, Hilary, 1723
Curtis, Neil F., 151
Curtis, Neville J., 1923
Curzon, Eirian H., 1361, 2195
D'Alfonso, Giuseppe, 1507
Dadkhah, Hengameh, 1523
Dagnall, Stephen P., 2381
Dahan, Françoise, 2521
Daniele, Pier G., 2353
Daolio, Sergio, 1547
Darkwa, James, 1435
Darwent, James R., 395
Datta, Dipankar, 159
Dauter, Zbigniew, 1235
Davidson, Jack L., 2231, 2239
Davies, Alwyn G., 471
Davies, Cathryn E., 669, 2037
Davies, Julian A., 209
Davies, Stephen G., 1479, 2691
Davis, Vivienne A., 1673
Dawes, Helen M., 921, 1775
Dénès, Georges, 1275
Deeming, Antony J., 85, 743, 857, 1037, 1609, 2479
de Graaff, Rudolf A. G., 737
Delgado, Esther, 1323
Deplano, Paola, 1349
de Queiroz, José C., 1103
Derakhshan, B. Mokhtar, 997
Deraniyagala, Samitha P., 1577
De Robertis, Alessandro, 2353
Derome, Andrew E., 423
De Stefano, Concetta, 2353
Dessy, Giulia, 1285
Di Bianca, Francesca, 523
Dillon, Keith B., 1399
Dilworth, Jonathan R., 1523, 1533, 2305, 2639, 2647
Ditzel, Evert J., 555
Di Vaira, Massimo, 291, 2327
Djuran, Miloš I., 861
Dobson, Alan, 611, 621
Dobson, Christopher M., 1811
Dobson, Ian D., 1213
Doddridge, Bruce G., 413
Dokuzović, Zdravko, 1095
Dondi, Stefano, 487
Donovan-Mtunzi, Susan, 1037, 1609, 2473, 2479
Dougherty, Geoffrey, 651
Dove, Michael F. A., 707, 2433, 2551
Downs, Anthony J., 807, 941
Doyle, Michael J., 229
Drew, Michael G. B., 69, 417, 771, 1073, 1771, 1821, 1829
Driessen, Willem L., 737, 1699, 2177
Dubchak, Inna L., 17
Dyason, Jeffrey C., 831, 839

Earl, Graham J., 663
Edwards, John O., 2269
Edwards, Peter G., 2167
Egsgaard, Helge, 879
Ehighaokhuo, Josephine O., 1665
Elbel, Susanne, 879
Ellis, Andrew, 879
Ellis, Arthur F., 1655
Ellis, John, 723, 1967
Ellul, Henriette, 503
Elrington, Margaret, 2407
Emo, Alan T., 1323
Engelhardt, Lutz M., 117, 125, 337, 831, 839, 981, 1747, 2541
Ercolani, Claudio, 1107, 1113
Eshima, Kiyoshi, 275
Espidel, Jouseff, 1859
Espinet, Pablo, 1163
Esteban, Montserrat, 973
Ettorre, Renato, 1271
Evans, Dennis F., 1141, 1147, 1151, 1451, 2525
Fairman, Kevin, 1523
Fanchiang, Yueh-Tai, 1375, 1381
Fanizzi, Francesco P., 1057
Fares, Vincenzo, 1285
Farrar, David H., 2595
Farrugia, Louis J., 177, 645, 2437
Fenton, David E., 1471
Fernández, Vicente, 1281
Fernanda, M., 2079
Fernandez-Baeza, Juan, 1163
Filipczuk, Stephen W., 1761
Filippeschi, Stefania, 523
Finch, Arthur, 997
Fitzgerald, William, 141
Fitzpatrick, Noel J., 1637
Flanders, David J., 517, 1001
Flint, Colin D., 1775
Font-Altaba, Manuel, 1087
Formicka-Kozlowska, Grazyna, 535
Forster, Alan M., 941
Foss, Olav, 231
Frampton, Christopher S., 169
Francis, Keith C., 493
Freeman, Mark J., 1019, 1027, 2283
Freni, Maria, 1507
Frost, Victoria L., 2059
Fuchita, Yoshio, 873
Fuger, Jean, 1853
Fukuzumi, Shunichi, 899
Fumagalli, Alesso, 1137
Furukawa, Haruhiko, 1713
Gabe, Eric J., 151
Gaines, Donald F., 1645
Gainsford, Graeme J., 151
Galas, Anita M. R., 611, 1339
Galli, Beatrice, 1155
Galli, Paola, 1737
Gamelkoorn, Hendrikus J., 659, 1495, 2223
Gans, Peter, 663, 1195
Garbauskas, Mary, 783
Garcia, J. R., 1865
Garcia, Maria P., 2343
Gardiner, Ian M., 669
Garner, C. David, 1977, 2161
Gasparrini, Francesco, 1155
Gates, Peter N., 997
Gavioli, Giovanna Battistuzzi, 2363
Gehring, Stefan, 2609
Germain, Gabriel, 2263
Ghilardi, Carlo A., 605, 2209
Ghosh, Phalguni, 361
Gibb, Terence C., 1455
Gibson, Vernon C., 2025
Giesbrecht, Ernesto, 2469
Gill, J. Bernard, 99, 663, 1213
Gillard, Robert D., 253
Gilmour, David I., 851, 1693
Gilmour, David W., 795
Giorgetti, Aldo, 885
Girolami, Gregory S., 685, 921, 1339
Glaves, Lyn R., 771
Gleria, Mario, 1547
Goddard, Jeffrey P., 471
Goggin, Peter L., 1597
Golding, Bernard T., 1997
Golub, Daniel, 641
Golubev, Sergei N., 17
González, Edgar, 1859
González, Juan M. Rubio, 2155
Goodall, David C., 99, 1213
Goode, Michael J., 941
Goodgame, Margaret, 75
Goodman, Bernard A., 1717
Gott, George A., 2661
Goudsmit, Robert J., 1795
Gracey, Benjamin P., 1935
Graham, Paul G., 777
Graimann, Christof E., 2025
Grandi, Giulia, 2363
Greatrex, Robert, 541
Grebenik, Peter D., 423, 669, 2025
Green, Jennifer C., 423, 669, 2037, 2677
Green, Malcolm L. H., 423, 669, 1585, 2025, 2037
Green, Michael, 177, 435, 2009, 2145, 2343, 2483
Greenhough, Trevor J., 761
Greenwood, Norman N., 541, 953, 1119, 1843, 2397, 2407
Griffith, William P., 717, 1673
Grilli, Ernesta, 2561
Gross, Maurice, 711
Grundy, Kevin R., 1577
Guarido, Carmen G., 1865
Guimon, Claude, 43
Gukathasan, Ravi R., 2595
Gulliver, David J., 1265
Gundersen, Grete, 191
Gupta, Minu, 2449
Gupta, Vishnu D., 2623
Gupta, Yugul Kishore, 2571
Gutiérrez-Puebla, Enrique, 1633
Guzmán, M. Carmen Carmona, 31
Hägele, G., 871
Haase, Wolfgang, 315, 913, 1243, 1909, 2609
Hague, David N., 2381
Hahn, Ekkehardt, 2427
Hails, Michael J., 693, 953
Halfpenny, Joan, 1399
Hall, Kevin P., 1811
Hallam, Malcolm F., 845
Hallett, Christopher, 451
Hanaki, Akira, 1513
Hancock, Robert D., 1877
Hankey, David R., 177
Harbron, Stephen K., 205
Harding, Majorie M., 1835
Hare, Philip M., 423, 2025
Harriman, Anthony, 503
Harris, Gillian W., 2277
Harris, R. K., 871
Harrison, Aidan T., 1173, 1953
Harrison, David, 885
Hašek, Jindřich, 2393
Hasegawa, Etsuo, 275
Hassan, Fatma S. M., 279, 1501
Hasselkus, Cynthia S., 209
Hatfield, William E., 2509
Hathaway, Brian, 141
Haupt, Steven G., 269
Hawkes, Geoffrey E., 225, 2473
Hayashida, Hideki, 873
Haymore, Barry L., 621
Hazel, Nicholas J., 669, 2037
Healy, Peter C., 111, 117, 125, 831, 839, 2531, 2541
Heath, Graham A., 947, 1081
Hegarty, Anthony F., 1915
Hempleman, Andrew J., 1775
Henderson, Richard A., 2059, 2067
Henrick, Kim, 1795
Hensen, Karl, 1, 5
Herak, Rajna, 861
Hibbert, Richard C., 707, 865, 2433, 2551
Higgins, Kevin M., 2195
Higuchi, Masahiro, 101
Hill, Ross H., 2421
Hill, William E., 1387, 2655
Hills, Adrian, 1069
Hiraki, Katsuma, 873
Hitchcock, Peter B., 1295, 1929
Hlatky, Gregory G., 1277
Hodgson, Diane, 1331
Holdcroft, Geoffrey E., 1731
Holden, H. Diane, 85
Hollands, Ronald E., 1527
Hollis, (the late) Stephen, 1829
Holloway, John H., 1853
Homsy, Nayla K., 2205
Honda, Atsunori, 59
Hope, Eric G., 529, 1265, 1443, 2185
Hosseiny, Afshin, 135
Howard, Christopher G., 921
Howard, Judith A. K., 777, 905, 1331, 2009, 2017
Howarth, Oliver W., 1173, 1953, 2195
Howells, Nigel D., 845
Huber, Friedo, 523
Hudson, Michael J., 771, 1655
Huffman, John C., 621
Hugel, René P., 597
Hughes, David L., 459, 1523
Hursthouse, Michael B., 85, 387, 471, 611, 685, 921, 931, 1043, 1339, 1775, 2167, 2321
Husbands, June M., 151
Hussain, M. Sakhawat, 749
Hussain, Wasif, 1131
Hutchinson, John, 1069, 2639, 2647
Hutchinson, John P., 1533
Huttner, Gottfried, 1061
Hutton, Alan T., 1677, 2121
Hynes, Michael J., 1543, 2565
Ibers, James A., 2369
Iftikhar, Khalid, 2547
Iggo, Jonathan A., 1009
Iijima, Kinya, 2555
Imperatori, Patrizia, 1285
Imran, Ahmed, 549
Inani, Krishnadass M., 2571
Innocenti, Paolo, 605, 2209
Interrante, Leonard V., 783
Ishikawa, Kunio, 899
Itoh, Takayuki, 2555
Iwamoto, Kyoko, 987
Iwamoto, Takeo, 81
Izquierdo, Alejo, 2037
Jackson, Peter F., 1795
Jacobsen, Grant B., 1645
James, Brian R., 617
Jeffery, John C., 645, 1315, 1323, 2001
Jeffreys, Brian, 99
Jerez, Antonio, 1633, 2225
Jewiss, Hilary C., 199
Jia-Long, Zhou, 991
Johnson, Brian F. G., 85, 555, 1277, 1723, 1795
Johnson, Owen, 2017
Jones, Christopher J., 401, 405, 1249
Jones, Peter G., 1163, 2417
Jones, Peter J., 529, 1443
Jones, Richard H., 1479
Judge, Austin I., 1853
Jump, Graham A., 541
Jund, Rodolphe, 711
Jurišić, Blaženka, 1095
Jutzi, Peter, 1303
Kabir, Shariff E., 1037, 2479
Kaden, Thomas A., 1169
Kaitner, Branko, 2457
Kakihana, Hidetake, 1733, 2427
Kalatzis, George, 2461
Kamabuchi, Keiko, 635
Kamenar, Boris, 2457
Kan, Chi Tat, 1523
Kanzaki, Tadao, 1713
Katakis, Dimitris, 2461
Katsura, Takashi, 1713
Katti, Kattesh V., 285, 2459
Kaufman, Clifford M., 307
Kazika, Anna I., 17
Kemmitt, Raymond D. W., 259, 549
Kemp, Terence J., 517
Kennard, Colin H. L., 243
Kennedy, John D., 953, 1119, 1843, 2397, 2407
Keszler, Douglas A., 2369
Khan, Fatima K., 333
Kida, Sigeo, 59, 1945, 2375
Kimura, Masaru, 355
Kinomura, Nobukazu, 2349
Kirschenbaum, Louis J., 1789, 2615
Kisenyi, Jonathan M., 69, 1073
Kishii, Noriyuki, 373
Klebe, Gerhard, 1, 5
Kleine-Boymann, Michael, 2493
Kleywegt, Gerard J., 2177
Knox, Selby A. R., 1935
Koga, Genji, 1959
Kohata, Susumu, 2575
Kojić-Prodić, Biserka, 455, 1531
Kondrashov, Yuri D., 17
Konstantatos, John, 2461
Kowalik, Teresa, 1201
Kozlowski, Henryk, 1201
Krasutskii, Pavel A., 17
Kratochvil, Bohumil, 2393
Krishnamurthy, Setharampattu S., 285, 1431, 1881, 2459
Kroto, Harold W., 1767
Kruger, Gert J., 1963
Kückelhaus, W., 871
Kumada, Nobuhiro, 2349
Kurmoo, Mohamedally, 579, 815
Kuroda, Reiko, 795
Kvammen, Frode, 231
Kwiatkowski, Edmund, 803, 2319
Kwiatkowski, Marek, 803
La Ginestra, Aldo, 1737
Laguna, Antonio, 2417
Laguna, Mariano, 2417
Lahiri, Sujit Chandra, 867

Lahoz, Fernando J., 1487, 1891
Laitinen, Risto, 1869
Lambrecht, Günter, 1743
Langrick, C. Richard, 511, 1015, 2121
Lappert, Michael F., 51, 229
Lappin, A. Graham, 2213
Laurie, Jill C. V., 2017
Lavery, Aidan, 1053, 1771
Lawrence, Geoffrey A., 1923
Legros, Jean-Pierre, 783
Lehn, Jean-Marie, 1517, 2311
Leigh, G. Jeffery, 1069, 1131, 2647
Leland, Jonathan K., 1303
Lelievre, Joel, 591
Lemoine, Paul, 711
León, Vladimir, 1859
Leoni, Piero, 2561
Leporati, Enrico, 1605
Leung, Wing-Por, 337, 1747
Levason, William, 199, 205, 529, 1265, 1443, 1735, 2185
Lewis, Diane B., 905, 2001
Lewis, Gregg E., 905, 2001
Lewis, Jack, 85, 333, 555, 1277, 1795
Lian, Lu Yun, 225
Lincoln, Stephen F., 413
Lindsay, Alan J., 2321
Linoh, Haryanto, 1743
Liptrot, Michael C., 333
Liu, Si-Han, 2217
Logan, Norman, 707, 865, 2433, 2551
Lucy, Andrew R., 699
Luke, Michael A., 851
Lyons, David, 587
Männig, Detlef, 1689
Ma, Emei, 2595
MacCurtain, Jacqueline, 2591
MacDonald, Colin J., 2473
MacLean, Gregory K., 1405
Macpherson, Kirsty A., 1935
Maddock, Alfred G., 991
Maeda, Yonezo, 2375, 2575
Maehara, Hiroaki, 1945
Mahmoud, Khalil A., 1365
Maitlis, Peter M., 573, 1555
Makino, Tadashi, 2139
Maleyev, Victor I., 17
Malik, Anis U., 2547
Malik, K. M. Abdul, 2167
Malini-Balakrishnan, Raman, 2291
Mangani, Stefano, 1179, 1425
Manhas, Baldev Singh, 1985
Mani, Fabrizio, 2327
Manivannan, Veeragathy, 2671
Mann, Brian E., 693, 1155, 1555
Manners, Ian, 2283
Manning, Peter J., 85, 1037
Manohar, Hattikudur, 1881, 2051
Manojlović-Muir, Ljubica, 861
Manotti-Lanfredi, Anna M., 1057, 1155
Manzano, Blanca R., 2417
Marøy, Kjartan, 231
Marangoni, Giampaolo, 731, 2091
Marder, Todd B., 2009
Maresca, Luciana, 1057, 1155
Markham, David P., 279
Markwick, Andrew J. W., 549
Marsden, Christine, 645, 1315
Marshall, Andrew, 1399
Marshall, Gregory L., 1265, 2185
Martin-Polo, Jesus J., 2037
Martinengo, Secondo, 35, 1137, 1309
Mason, Joan, 2473
Massucci, Maria A., 1737
Mathieu, René, 1087, 1981, 2521
Matisons, Janis G., 1223
Matsubara, Chiyo, 81
Matsubayashi, Gen-etsu, 465
Matsumoto, Keiji, 2095
Matsumoto, Naohide, 101, 107, 2575
Matsushita, Yoh-ichi, 275
Mawby, Roger J., 1235
Mays, Martin J., 1355
Mazid, Muhammed A., 795
McAdam, Michael E., 2381
McAuliffe, Charles A., 135, 1387, 1391, 2655, 2661
McCleverty, Jon A., 401, 405, 1249
McDonald, Walter S., 953, 1843
McEwan, David M., 1501, 2121
McGinn, Michael A., 1637
McKean, Donald C., 1207
McKee, Vickie, 1771
McKenna, Peter, 259, 549
McNaughton, Donald, 1767
McPartlin, Mary, 1795
McPhail, Donald B., 1717
McQuillan, Geoffrey P., 1207
McWhinnie, William R., 821
Mealli, Carlo, 479
Meanwell, Neil J., 1555
Meina, Dominic G., 1645, 1903
Menabue, Ledi, 2363
Mentasti, Edoardo, 1789, 2615
Meyerstein, Dan, 641
Micheloni, Mauro, 1169
Midollini, Stefano, 605, 2209
Migita, Catharina T., 1077
Miles, Anthony D., 2437
Mills, Nancy K., 111
Mills, Rona M., 699
Minchin, Nigel J., 1183
Mingos, D. Michael P., 845, 851, 1693, 1811, 2343
Miravitlles, Carles, 1981, 2333
Mishra, Anjali, 2449
Missen, Paul H., 1147, 1451
Mittal, Pramod K., 1561, 1569
Miyamura, Kazuo, 2139
Mockford, M. Jane, 717
Molinari, Henriette, 1899
Molloy, Kieran C., 1417
Monacelli, Fabrizio, 1107, 1113
Monge, Angeles, 1633
Moon, Steven D., 1479
Moore, David S., 611, 617
Moore, Peter, 219, 1361
Moreton, Anthony D., 2381
Mori, Nobuo, 1065, 1661
Mori, Wasuke, 2095
Morley, Christopher P., 1303
Morpurgo, Giorgio O., 1285
Morris, John H., 1645, 1903
Morrisson, Andrew R., 1207
Morton, Carolyn E., 2145
Morton, Stephen, 1295
Moser, Ulrich, 1743
Mosset, Alain, 1597
Motevalli, Majid, 685, 921, 931, 2167, 2321
Mtetwa, Victor S. B., 669, 2037
Mueller, Melvin H., 2369
Mukaida, Masao, 1733, 2427
Muller, Guillermo, 2333
Mura, Pasquale, 2101
Murati, Ivo, 1095
Murr, Nabil E. 1249
Murray, Martin, 177
Murray, Stephen G., 1265, 2185
Muto, Fumio, 2349
Mutschler, Ernst, 1743
Mwesigye-Kibende, Samson, 1997
Nagle, Katharine R., 2145
Nakahara, Akitsugu, 2095
Nakamura, Akira, 635
Nakamura, Masaaki, 59
Nakao, Yasuo, 2095
Nakatsuka, Kunio, 2095
Nandana, W. A. Shantha, 1623
Nardelli, Mario, 321, 487
Nardi, Nicoletta, 479, 1179, 1425
Narula, Chaitanya K., 2623
Natile, Giovanni, 1057, 1155
Neaves, Bryan D., 1249, 2647
Neidle, Stephen, 795
Nelson, S. Martin, 1053, 1771
Nelson, William J. H., 1795
Nepveu, Françoise, 315
Nguyen, Minh Tho, 1915
Nicholls, Barry S., 1835
Nicholson, Terry, 2639
Niecke, Edgar, 879
Nishida, Suzuko, 355
Nishida, Yuzo, 1945, 2375
Nishide, Hiroyuki, 65, 275
Nishikawa, Yuji, 81
Nixon, John F., 1295
Nöth, Heinrich, 1689, 2623
Nogami, Hidenori, 101
Nohr, Ronald S., 269
Noltemeyer, Mathias, 565, 2205
Noordik, Jan N., 2177
Norman, Nicholas C., 383, 435, 1303
Normanton, Fred B., 2647
Novikova, Marina I., 17
Nowell, I. W., 2505
Nuber, Bernhard, 2493
Nunn, Michael, 2433
Öhman, Lars-Olof, 2665
Ökawa, Hisashi, 59
O'Brien, Paul, 2085
O'Hare, Dermot, 1585
O'Reilly, Eric J., 243
Oakes, John, 493
Ochi, Naoyuki, 873
Odell, Barbara, 2161
Odiaka, Timothy I., 1049
Ogden, J. Steven, 529, 1443
Ogura, Kotaro, 1077, 2499
Ohkubo, Katsutoshi, 1959
Ohta, Shoichi, 2575
Ohya, Toshie, 987
Ohyoshi, Akira, 101, 107, 2575
Ojo, J. Folorunso, 1665
Okawa, Hisashi, 2575
Okey, J. Nigel, 75
Olby, Bruce G., 2101
Ollerenshaw, Timothy J., 2161
Olubuyide, Olusegun, 1665
Omar, Hadi A. A., 219, 1361
Ondo, Benjamin Mve, 597
Ooi, Shun'ichiro, 2095
Orama, Olli, 2079
Orioli, Pier Luigi, 1179, 1425
Orlandini, Annabella, 605, 2209
Oro, Luis A., 973, 1487, 1891
Orpen, A. Guy, 177, 435, 1019, 1027, 1935, 2145, 2283, 2483
Orrell, Keith G., 345, 1561, 1569, 2195
Osborne, Anthony G., 1527
Ozawa, Toshihiko, 1513
Paap, Frans, 737
Pakawatchai, Chaveng, 117, 125, 831, 839, 2531
Pakulski, Marek, 383
Pal, Amalendu, 867
Pal, Samundranil, 159
Pampaloni, Guido, 1989
Pannetier, Jean, 1275
Paoletti, Piero, 1169
Parish, R. V. (Dick), 1391
Parker, David, 1517, 2311
Parmar, Swarn S., 2221
Pascal, Jean-Louis, 297
Pasquali, Marco, 2561
Passmore, Jack, 9, 1405, 1623
Pastore, Annalisa, 1899
Patrick, Jennifer M., 981, 1183, 1223, 1229
Patrick, Vincent A., 831, 839, 2541
Paulus, Helmut, 913
Pavlović, Dušanka, 1095
Paz-Sandoval, Maria de los Angeles, 2677
Pérez, Pedro L., 973
Peart, Barry J., 795
Pelizzi, Corrado, 215, 321, 487, 2387
Pelizzi, Giancarlo, 215, 321, 487, 2387
Pellacani, Gian Carlo, 2363
Pellinghelli, Maria Angela, 1349
Pennesi, Giovanna, 1107, 1113
Pennington, Mark, 2683
Pepliński, Zbigniew, 2319
Peringer, Paul, 223, 1061
Persson, Ingmar, 1597
Peruzzini, Maurizio, 291
Peters, Roy, 85
Pettit, Leslie D., 535, 1201
Pezeshk, Abbas, 891
Pfister-Guillouzo, Geneviève, 43
Pickardt, Joachim, 2427
Pickett, Christopher J., 1131, 1255, 2079
Pico, Carlos, 2225
Pierpoint, Colin, 219, 1361
Pignatello, Joseph J., 1381
Pilbrow, John R., 651
Pilkington, Roger S., 1405
Pilling, Michael J., 2247
Pimblett, Gillian, 1977
Pinillos, M. Teresa, 1487
Pinkert, Waltraud, 565
Pitteri, Bruno, 2091
Pizzotti, Maddalena, 163
Platt, Andrew W. G., 345
Plazzogna, Gualtiero, 1271
Pocev, Stefan, 2457
Podlaha, Jaroslav, 2393
Podlahová, Jana, 2393
Poilblanc, René, 43
Poli, Rinaldo, 931
Pombeiro, Arm J. L., 2079
Poojary, M. Damodara, 1881
Porta, Francesca, 163
Potier, Jacqueline, 297
Powell, Paul, 2677
Power, Philip P., 51
Pratt, John M., 1613, 1619
Pratten, Stephen J., 1761
Predieri, Giovanni, 487, 2387
Preece, Michael, 617
Pringle, Paul G., 279, 1015, 1501, 1677, 2121
Proud, Graeme P., 857
Prout, Keith, 669, 2025, 2037
Puddephatt, Richard J., 2421
Puga, José, 1795

Purkayastha, Ranendra N. Dutta, 409
Pye, Peter L., 229
Qing-jin, Meng, 2269
Quill, Kieran, 1417
Rábai, Gyula, 1669
Radanović, Dušan J., 861
Raithby, Paul R., 333, 555, 1277, 1355
Ramsden, Christopher A., 795
Randall, Edward W., 225
Rankin, David W. H., 191, 755, 807, 827, 941
Rankin, Douglas A., 1131
Rao, M. N. Sudheedra, 1405
Rao, S. P. Sudhakara, 2051
Raston, Colin L., 337, 831, 1747
Raubenheimer, Helgard G., 1963
Raven, Stephen J., 2283
Read, Gordon, 1591
Reddy, K. Veera, 2603
Reddy, M. Harilatha, 239
Reddy, M. Satyanarayan, 2603
Reddy, P. Rabindra, 239
Reed, David, 1645
Reedijk, Jan, 737, 1699, 2177
Rest, Antony J., 1365
Reynolds, Colin D., 1235
Rice, David A., 417, 1821
Richards, John P. G., 253
Richards, Raymond L., 1523, 2473
Richoux, Marie-Claude, 503
Rigano, Carmelo, 2353
Rimmer, John, 1517
Risse, Wilhelm, 777
Ritchie, Ian M., 1747
Rivarola, Eleonora, 523
Robbins, David J., 1781
Roberts, Roger M. G., 169
Robertson, Heather E., 191, 807, 827, 941
Robinson, Stephen D., 611, 617, 621, 629, 2101, 2113
Robinson, Ward T., 1289
Rocamora, Mercè, 2333
Rodríguez-Castellón, Enrique, 213
Rodríguez-García, Aurora, 213
Rodríguez, Julio, 1865
Roesky, Herbert W., 565, 2205, 2453
Roge, Gerhard, 523
Rojas, Ramón L. Espinoza, 2469
Ros, Josep, 1087, 1981
Rossi, Gentilina, 1113
Rothin, Anne S., 401, 405
Ruebenbauer, Krzysztof, 1275
Ruisi, Giuseppe, 523
Ruiz-Valero, Caridad, 1633
Russell, David R., 259, 549
Ružić-Toroš, Živa, 455, 1531
Ryabov, Alexr D., 2629
Ryzhov, Mikhail G., 17
Šik, Vladimir, 345, 1561, 1569, 2195
Sánchez-Burgos, Francisco, 1975
Sánchez-Delgado, Roberto A., 1859
Sabat, Michal, 479
Sabatini, Antonio, 1195
Saburi, Masahiko, 2139
Sadler, Peter J., 795
Sakaki, Shigeyoshi, 1959
Sakodinskaya, Inna K., 2629
Salagre, Pilar, 2263
Sales, Keith D., 225
Salt, Julian E., 685
Sammartano, Silvio, 2353
Sandström, Magnus, 1597
Sarisaban, Serap, 1929
Sasada, Yoko, 873
Sato, Fumio, 1959
Sato, Mitsunobu, 895
Sato, Mitsuo, 987
Sato, Yoichi, 895
Saunders, David R., 1235
Savage, (née Roquet-Covarrubias), Sonia, 991
Schaverien, Colin J., 2483
Schlüter, Ewald, 1303
Schmidbaur, Hubert, 827
Schriver, Melbourne J., 1405
Schubert, Ulrich, 2079
Schumann, Herbert, 2427
Schwabe, Ludwig, 1909
Scimar, Christiane N., 209
Seddon, Kenneth R., 2247
Seeber, Renato, 601
Seega, J., 871
Seeman, Jeffrey I., 2691
Seip, Ragnhild, 827
Semelhago, Gavin, 2595
Sen, Buddhadev, 307
Sengupta, Dipak, 867
Seregni, Claudio, 1309
Sertić, Mira, 1095
Seseke, Ulrich, 565
Sethulekshmi, C. N., 2509
Shand, Mark A., 517
Sharma, Prem Dutt, 2571
Shaw, Bernard L., 279, 511, 1009, 1015, 1213, 1501, 1677, 2121, 2131
Sheehan, Alan R., 1019
Sheldrick, George M., 565, 1163, 2205, 2417
Sheldrick, William S., 1743
Sheridan, John B., 699, 1019, 1027
Sherry, Lesley J. S., 259, 549
Shibata, Shuzo, 2555
Shimizu, Kunio, 1733
Shimo, Hiroyuki, 1945
Shiraishi, Shinsaku, 373
Shklover, Valeri E., 17
Siddiqui, Md. Rafiq H., 2603
Siddiqui, Zafar A., 2453
Siegfried-Hertli, Liselotte, 1169
Sigel, Helmut, 2291
Silver, Jack, 169
Silvestri, Arturo, 523
Simmons, Charles J., 141
Simmons, Nicholas D., 1323
Simpson, Stephen J., 1479
Singh, Harkesh B., 821
Sironi, Angelo, 1507, 1795
Sjöberg, Staffan, 2665
Skarda, Vladimir, 1781
Skelton, Brian W., 1183
Skorobogaty, Andrew, 651
Smith, Anthony K., 1835
Smith, Deborah J., 1811
Smith, Garry, 387
Smith, Graham, 243
Smith, J. David, 1929
Smith, Thomas D., 651
Sola, Marco, 2363
Solans, Xavier, 1087, 1981, 2263, 2333
Somerville, Richard G., 2343
Sood, Anup, 209
Sosa, Martha E., 475
Sowerby, D. Bryan, 2433
Spalding, Trevor R., 2591
Spencer, Catriona M., 693, 1555
Spreafico, Federico, 523
Sreelatha, Channareddy, 2623
Stacey, Catherine, 135
Stavropoulos, Pericles, 2167
Steel, Ian, 535, 1201
Stephens, Michael, 997
Stephenson, G. Richard, 1723
Stephenson, T. Anthony, 947
Steudel, Ralf, 1869
Stobart, Stephen R., 1935
Stone, F. Gordon A., 177, 645, 905, 1315, 1323, 1331, 2001, 2009, 2017, 2343, 2437
Stoppioni, Piero, 291
Strauss, Eva-Maria, 1869
Struchkov, Yuri T., 17
Strumolo, Donatella, 35, 1309
Suárez, Marta, 1865
Sueiras, Jesús-Eduardo, 2263
Sutcliffe, Leslie H., 1405
Suzuki, Shinnichiro, 2095
Swamy, K. C. Kumara, 1431, 1881
Sweigart, Dwight A., 1019, 2269
Swiridoff, Wolfgang, 2493
Sylva, Ronald N., 723, 1967
Symonds, Andrew N. D., 1027
Symons, Martyn C. R., 379
Szpakowska, Maria, 1849
Tacke, Reinhold, 1743
Tajik, Mahmoud, 199, 529, 1443, 1735
Takahashi, Kazuhiro, 2375
Takamori, Michiko, 1065, 1661
Takamura, Kiyoko, 81
Takemura, Tetsuo, 1065
Tames, Joseph, 1391
Tanaka, Toshio, 465, 899
Tanner, S. Peter, 2661
Taqui Khan, Badar, 2603
Taqui Khan, Mirza M., 2603
Tarasconi, Pieralberto, 215, 321
Tatarowski, Tomasz, 535
Tate, John R., 1471
Tattershall, Bruce W., 1707
Taylor, Brian F., 1555, 2505
Taylor, Michael J., 555
Taylor, Norman, 1213
Taylor, Peter, 9
Tellinghuisen, Patricia C., 1289
Terreros, Pilar, 229
Thöm, Vivienne J., 1877
Thomas, Gillian J. N., 1303
Thomas, Peter M., 1471
Thompson, Andrew J., 1781
Thomson, Colin, 1645
Thorburn, Ian, 617
Thornton-Pett, Mark, 387, 685, 921, 1119, 1339, 2131, 2397, 2407
Tiripicchio-Camellini, Marisa, 973
Tiripicchio, Antonio, 973, 1057, 1155, 1487, 1891
Tobe, Martin L., 27, 475, 731, 2091
Toledo, Jose Hidalgo, 31
Toma, Henrique E., 2469
Tomassini, Marco, 2561
Tomlinson, Anthony A. G., 1655
Tondreau, Gerard A., 2269
Tornero, Jesus D., 1281
Torto, Ivy, 1207
Tossing, G., 871
Townend, Timothy J., 1673
Traldi, Pietro, 1547
Tribolet, Roger, 2291
Trikha, Arun Kumar, 1985
Trogu, Emanuele Filiberto, 1349
Tso, Tze Chung, 1853
Tsuchida, Eishun, 65, 275
Tudela, David, 1281
Turff, Jeremy W., 529, 1443
Turp, Janet E., 2247
Ueyama, Kosuke, 465
Ueyama, Norikazu, 635
Ugo, Renato, 1971
Uma, Venkataraman, 209
Underhill, Allan E., 1731
Upton, Mark W., 1141, 1147, 1151, 2525
Urgelles, Miguel, 1591
Uruska, Irmina, 1849
Usón, Rafael, 2417
Vacca, Alberto, 1195
Valade, Lydie, 783
Valencia, Norma, 1859
Valle, Giovanni, 1271
Van, Driel, Gertjan J., 2177
Vargas, Maria D., 1795
Varshney, Krishna G., 1737
Vasapollo, Guiseppe, 2231, 2239
Vauchskii, Yuri P., 17
Vecchi, Enrico, 1547
Veiga, Maria Luisa, 2225
Velasco, Julian Rodriguez, 31
Vicente, José, 1163
Viljoen, Hendrik W., 1963
Villarroya, B. Eva, 1891
Vitali, Francesca, 2387
Vitt, Sergei V., 17
Vrachnou-Astra, Ersi, 2461
Walker, Alan, 2595
Walker, Nigel P. C., 85, 199, 471, 1043
Wallbridge, Malcolm G. H., 761
Walsh, James, 1543, 2565
Walz, Leonhard, 315, 913, 1243
Wandiga, Shem O., 69
Wardle, Robert W. M., 845, 1693
Warmsley, Joanne F., 169
Warrens, Christopher P., 1043
Watts, Andrew M., 2191
Weber, David C., 269
Weber, Karl-Ludwig, 565
Webster, Michael, 199, 1735
Weiss, Johannes, 2493
Welch, Alan J., 1645, 2343
Went, Michael J., 1331
West, Douglas X., 379
Whidden, Tom K., 9
White, Allan H., 111, 117, 125, 243, 337, 831, 839, 981, 1183, 1223, 1229, 1747, 2531, 2541
White, Peter S., 9, 1405, 1435, 1623
Whiteley, Mark W., 777, 1027
Whiteley, Richard H., 1527
Whitelock, John D., 947
Wickens, Denys A., 2203
Wieghardt, Karl, 2493
Wiesmeijer, Willem G. R., 2177
Wilkins, Cuthbert J., 1289
Wilkinson, Geoffrey, 587, 685, 921, 931, 1339, 2167, 2321
Willey, Gerald R., 69, 1073, 2683
Williams, David M., 417, 1821
Williams, Ian D., 435, 2483
Williams, Michael L., 1229
Willis, Anthony C., 1223
Wilson, Christina L., 1929
Wimmer, Franz L., 253
Wingfield, Jonathan N., 459
Winkler, Peter-Paul, 1061
Woodward, Peter, 699
Woollins, J. Derek, 1043, 2397, 2407
Wynne, Kenneth J., 269
Yamanaka, Shunichiro, 873

Yamasaki, Shinji, 2499
Yamase, Toshihiro, 2585
Yano, Shigenobu, 81, 895
Yates, Paul C., 1829
Yatsimirski, Anatoli, 1971
Yatsimirsky, Anatoly K., 2629
Yavari, Ahmad, 2421, 2509
Yavari, Bahrum, 169
Yellowlees, Lesley J., 947, 1081
Yoshikawa, Sadao, 81, 895, 2139
Yoshimura, Chikako, 2575
Yoshino, Takashi, 1077
Yuasa, Makoto, 65, 275
Yurchenko, Aleks G., 17
Zanazzi, Pier Francesco, 1989
Zanello, Piero, 601
Zanobini, Fabrizio, 479, 1179, 1425
Zhang, Cheng Shan, 297
Zielkiewicz, Jan, 1849
Zsolnai, Laszlo, 1061
Zubieta, Jon A., 1069, 1533, 2639, 2647

SUBJECT INDEX, 1985

ACETALDEHYDE
Reactions of hydride reagents with alkylmolybdenum carbonyl complexes. Reaction of $LiBHEt_3$ with $[MoMe(CO)_3(\eta\text{-}C_5H_5)]$, formation of an anionic acetaldehyde complex, and a stoicheiometric cycle for the synthesis of acetaldehyde, 1815–20

ACETALDOXIME
Stabilisation of copper(I) by an azoimine ligand: redox properties and reactions of bis(phenylazoacetaldoximato)bis(phenylacetaldoxime)dicopper(I), 159–62

ACETATE
The crystal structure and electronic properties of the complex acetatobis(1,10-phenanthroline)copper(II) perchlorate dihydrate, acetatobis(1,10-phenanthroline)copper(II) nitrate dihydrate, and acetatobis(1,10-phenanthroline)zinc(II) tetrafluoroborate dihydrate, 141–50

Actinide structural studies. Part 7. The crystal and molecular structures of (2,2′-bipyridyl)dinitratodioxo-uranium(VI) and -neptunium(VI), and diacetato(2,2′-bipyridyl)dioxo-uranium(VI) and -neptunium(VI), 1001–8

Dirhodium(II,II) tetra-acetate complexes with axially co-ordinated triphenylstibine, triphenylarsine, and dibenzyl sulphide ligands. The syntheses, properties, and *X*-ray crystal structures of $[Rh_2(O_2CMe)_4(SbPh_3)_2]$, $[Rh_2(O_2CMe)_4(AsPh_3)_2]$, and $[Rh_2(O_2CMe)_4\{S(CH_2Ph)_2\}_2]$, 1775–80

Equilibria between mono- and bi-nuclear complexes in $Cu(O_2CMe)_2$–pyridine derivative–diluent systems. The influence of the amine ligand basicity, 1849–52

The easy catalytic reduction of $HClO_4$, HNO_3, and H_5IO_6 by CO in the presence of $[Pd_3(O_2CMe)_6]$, 1971–4

Magnetic and spectroscopic properties of some heterotrinuclear basic acetates of chromium(III), iron(III), and divalent metal ions, 2509–20

Molecular structure of gaseous copper(I) acetate as determined by electron diffraction, 2555–60

Kinetics and mechanism of ortho-palladation of ring-substituted *NN*-dimethylbenzylamines, 2629–38

ACETONITRILE
Isolation and crystal structure of $[Rh(PPh_3)_3(MeCN)][BF_4]$, acetonitriletris(triphenylphosphine)rhodium(I) tetrafluoroborate, 1977–80

ACETYLACETONATE
Electrostatic solvent effect on the formation of the mixed-chelate complex (acetylacetonato)(diethyldithocarbamato)-copper(II), 987–90

ACETYLENEDICARBOXYLATE
Bimetallic systems. Part 10. Synthesis of complexes of type $[(RC{\equiv}C)Pt(\mu\text{-}dppm)_2Pt(C{\equiv}CR)]$ (dppm = $Ph_2PCH_2PPh_2$, R = Ph or *p*-tolyl) and their corresponding 'A frames' $[(RC{\equiv}C)Pt(\mu\text{-}dppm)_2(\mu\text{-}H)Pt(C{\equiv}CR)]Cl$ or $[(RC{\equiv}C)Pt(\mu\text{-}dppm)_2(\mu\text{-}X)Pt(C{\equiv}R)]$ with X = CS_2 or MeOOCC≡CCOOMe, 1015–8

ACETYLIDE
Bimetallic systems. Part 10. Synthesis of complexes of type $[(RC{\equiv}C)Pt(\mu\text{-}dppm)_2Pt(C{\equiv}CR)]$ (dppm = $Ph_2PCH_2PPh_2$, R = Ph or *p*-tolyl) and their corresponding 'A frames' $[(RC{\equiv}C)Pt(\mu\text{-}dppm)_2(\mu\text{-}H)Pt(C{\equiv}CR)]Cl$ or $[(RC{\equiv}C)Pt(\mu\text{-}dppm)_2(\mu\text{-}X)Pt(C{\equiv}R)]$ with X = CS_2 or MeOOCC≡CCOOMe, 1015–8

Bimetallic systems. Part 12. Mixed rhodium(I)–platinum(II) acetylide complexes containing bridging $Ph_2PCH_2PPh_2$. Crystal structures of $[(MeC{\equiv}C)Pt(\mu\text{-}dppm)_2(\sigma,\eta\text{-}C{\equiv}CMe)Rh(CO)]PF_6$ and of $[ClPt(\mu\text{-}dppm)_2(\sigma,\eta\text{-}C{\equiv}CMe)Rh(CO)]PF_6$, 2121–30

ACTINIDE
Actinide structural studies. Part 7. The crystal and molecular structures of (2,2′-bipyridyl)dinitratodioxo-uranium(VI) and -neptunium(VI), and diacetato(2,2′-bipyridyl)dioxo-uranium(VI) and -neptunium(VI), 1001–8

ACTIVITY COEFFICIENT
On the possibility of determining the thermodynamic parameters for the formation of weak complexes using a simple model for the dependence on ionic strength of activity coefficients: Na^+, K^+, and Ca^{2+} complexes of low molecular weight ligands in aqueous solution, 2353–62

ADENOSINE TRIPHOSPHATE
Influence of decreasing solvent polarity (dioxane–water mixtures) on the stability and structure of binary and ternary complexes of adenosine 5′-triphosphate and uridine 5′-triphosphate, 2291–304

ADENOSYL
A new electron spin resonance spectrum for an exchange- and dipole–dipole coupled superoxocobalamin · · · free-radical pair occurring in adenosylcobalamin-containing systems, 891–4

ALCOHOL
Aluminium-27 and hydrogen-1 nuclear magnetic resonance studies of solutions of aluminium salts in alcohol–chloroform mixtures, 591–6

ALKALI METAL
Synthesis and structural assessment of ammonium and caesium difluorodioxoperoxouranates(VI), $A_2[UO_2(O_2)F_2]$ (A = NH_4 or Cs), and alkali-metal difluorodioxoperoxouranate(VI) monohydrates, $A_2[UO_2(O_2)F_2]\cdot H_2O$ (A = K or Rb), 409–12

Co-ordination of alkali metals by open-chain polyethers in transition metal complexes. Part 4. Variation in alkali-metal ion selectivity in cobalt and zinc complexes of 1-(*o*-carboxymethoxyphenoxy)-2-(*o*-hydroxyphenoxy)ethane (HL) and the *X*-ray and molecular structure of $[Zn(NH_4L_2)_2]$, 459–64

Synthesis and properties of alkali-metal tetraoxo-osmate(VII) compounds, 1735–6

On the possibility of determining the thermodynamic parameters for the formation of weak complexes using a simple model for the dependence on ionic strength of activity coefficients: Na^+, K^+, and Ca^{2+} complexes of low molecular weight ligands in aqueous solution, 2353–62

ALKANONE
Reactions of tris[(2*S*)-2-(aminomethyl)pyrrolidine]nickel(II) ion with alk-3-en-2-ones or 4-hydroxyalkan-2-ones: formation of an optically active tetra-aza macrocycle, 2139–44

ALKENONE
Reactions of tris[(2*S*)-2-(aminomethyl)pyrrolidine]nickel(II) ion with alk-3-en-2-ones or 4-hydroxyalkan-2-ones: formation of an optically active tetra-aza macrocycle, 2139–44

ALKYL
Electron-transfer reactions from *cis*-dialkylbis(2,2′-bipyridyl)cobalt(III) complexes to organic oxidants, 899–904

Alkyl, hydrido, and tetrahydroaluminato complexes of manganese with 1,2-bis(dimethylphosphino)ethane (dmpe). *X*-Ray crystal structures of $Mn_2(\mu\text{-}C_6H_{11})_2(C_6H_{11})_2(\mu\text{-dmpe})$, $(dmpe)_2Mn(\mu\text{-}H)_2AlH(\mu\text{-}H)_2AlH(\mu\text{-}H)_2Mn(dmpe)_2$, and $Li_4\{MnH(C_2H_4)[CH_2(Me)PCH_2CH_2PMe_2]_2\}_2\cdot 2Et_2O$, 921–30

Synthetic and nuclear magnetic resonance studies on dialkyl- and diaryl-platinum complexes containing chelating, monodentate, or bridging $Ph_2PCH_2PPh_2$ ligands, 1501–6

Reactions of hydride reagents with alkylmolybdenum carbonyl complexes. Reaction of $LiBHEt_3$ with $[MoMe(CO)_3(\eta\text{-}C_5H_5)]$, formation of an anionic acetaldehyde complex, and a stoicheiometric cycle for the synthesis of acetaldehyde, 1815–20

Some alkyldiplatinum(I) complexes and studies of the photochemical and thermal decomposition of the $[Pt_2Et(\mu\text{-}Ph_2PCH_2PPh_2)_2(Ph_2PCH_2PPh_2\text{-}P)]^+$ cation, 2421–6

ALKYLAMINE
Intercalation of alkylamines into tin(IV) bis(hydrogenphosphate) monohydrate, 213–4

ALKYLCOBALAMIN
Kinetic studies of oxidative dealkylation of alkylcobalamins by hexachloroplatinate(IV), 1375–80

ALKYLCORRINOID
The chemistry of vitamin B_{12}. Part 25. Mechanism of the β-elimination of olefins from alkylcorrinoids; evidence for an initial homolytic fission of the Co–C bond, 1613–8

ALKYLIDENE
Organic chemistry of dinuclear metal centres. Part 8. Organo-iron–ruthenium chemistry. *X*-Ray structure of *trans*-$[FeRu(CO)_2(\mu\text{-}$

ALKYLIDENE (cont'd)
$CO)_2(\eta\text{-}C_5H_5)_2$], 1935–44
ALKYLIDYNE
Formation of substituted cyclopentadienyl ligands on tungsten *via* reactions between the alkyne complexes $[W(CO)(R^1C_2R^2)_3]$ ($R^1 = R^2$ = Ph or Et; R^1 = Me, R^2 = Ph) and the alkylidyne compounds $[W(\equiv CR)(CO)_2(\eta\text{-}C_5H_5)]$ (R = C_6H_4Me-4 or Me); *X*-ray crystal structures of $[W_2(\mu\text{-}CO)_2(CO)(\eta\text{-}PhC_2Ph)(\eta\text{-}C_5Ph_4R)(\eta\text{-}C_5H_5)]$ and $[W_2(\mu\text{-}EtC_2Et)(CO)_4(\eta\text{-}C_5Et_4R)(\eta\text{-}C_5H_5)]$ (R = C_6H_4Me-4), 905–12
ALKYLPEROXO
Routes of formation and crystal structure of an alkylperoxycobaloxime: bis[dimethylglyoximato(1 −)](4-ethoxycarbonylbut-3-en-2-ylperoxo)(pyridine)cobalt(III), 1997–2000
ALKYNE
Formation of substituted cyclopentadienyl ligands on tungsten *via* reactions between the alkyne complexes $[W(CO)(R^1C_2R^2)_3]$ ($R^1 = R^2$ = Ph or Et; R^1 = Me, R^2 = Ph) and the alkylidyne compounds $[W(\equiv CR)(CO)_2(\eta\text{-}C_5H_5)]$ (R = C_6H_4Me-4 or Me); *X*-ray crystal structures of $[W_2(\mu\text{-}CO)_2(CO)(\eta\text{-}PhC_2Ph)(\eta\text{-}C_5Ph_4R)(\eta\text{-}C_5H_5)]$ and $[W_2(\mu\text{-}EtC_2Et)(CO)_4(\eta\text{-}C_5Et_4R)(\eta\text{-}C_5H_5)]$ (R = C_6H_4Me-4), 905–12
Carbon–carbon formation at di-iron centres. Part 2. Reactivity of $[Fe_2(CO)_6(\mu\text{-}COEt)\{\mu\text{-}C(R)C(R)H\}]$ complexes toward MeOC(O)C≡CC(O)OMe (R = Ph) and $CF_3C{\equiv}CCF_3$ (R = Ph or H); *X*-ray crystal structures of $[Fe_2(CO)_5\{\mu\text{-}C(OEt)C[C(O)OMe]C[C(O)OMe]\}\{\mu\text{-}C(Ph)C(Ph)H\}]\cdot H_2O$ and $[Fe_2(CO)_6\{\mu\text{-}C(CF_3)C(CF_3)CHCHCH(OMe)\}]$, 1087–94
Organic chemistry of dinuclear metal centres. Part 8. Organo-iron–ruthenium chemistry. *X*-Ray structure of *trans*-$[FeRu(CO)_2(\mu\text{-}CO)_2(\eta\text{-}C_5H_5)_2]$, 1935–44
Synthesis and nuclear magnetic resonance studies of co-ordinatively unsaturated alkyne complexes of tungsten(II), 2239–46
ALUMINIUM
Partially oxidized group 3B fluorometallophthalocyanines, 269–74
Aluminium-27 and hydrogen-1 nuclear magnetic resonance studies of solutions of aluminium salts in alcohol–chloroform mixtures, 591–6
Complexes of organoaluminium compounds. Part 13. Preparation and nuclear magnetic resonance spectra of the arylamido-compounds $AlMe_2(NHR')$ (R' = Ph, C_6H_4Me-*o*, C_6H_4Me-*p*, $C_6H_3Me_2$-2,6) and the imido-compounds $AlMe(NR')$. Crystal and molecular structures of $[\{AlMe_2(NHC_6H_4Me\text{-}o)\}_2]$ and $[\{AlMe(NPh)\}_6]$, 1929–34
The hydrolysis of metal ions. Part 8. Aluminium(III), 1967–70
Equilibrium and structural studies of silicon(IV) and aluminium(III) in aqueous solution. Part 13. A potentiometric and ^{27}Al nuclear magnetic resonance study of speciation and equilibria in the aluminium(III)–oxalic acid–hydroxide system, 2665–70
ALUMINIUM-27
Aluminium-27 and hydrogen-1 nuclear magnetic resonance studies of solutions of aluminium salts in alcohol–chloroform mixtures, 591–6
Equilibrium and structural studies of silicon(IV) and aluminium(III) in aqueous solution. Part 13. A potentiometric and ^{27}Al nuclear magnetic resonance study of speciation and equilibria in the aluminium(III)–oxalic acid–hydroxide system, 2665–70
AMIDE
Subvalent group 4B metal alkyls and amides. Part 7. Transition-metal chemistry of metal(II) bis(trimethylsilyl)amides $M'(NR_2)_2$ (R = $SiMe_3$; M' = Ge, Sn, or Pb), 51–8
Complexes of organoaluminium compounds. Part 13. Preparation and nuclear magnetic resonance spectra of the arylamido-compounds $AlMe_2(NHR')$ (R' = Ph, C_6H_4Me-*o*, C_6H_4Me-*p*, $C_6H_3Me_2$-2,6) and the imido-compounds $AlMe(NR')$. Crystal and molecular structures of $[\{AlMe_2(NHC_6H_4Me\text{-}o)\}_2]$ and $[\{AlMe(NPh)\}_6]$, 1929–34
AMINE
Determination of the molecular structures and conformations of methylbis(methylsilyl)amine and bis(dimethylsilyl)methylamine in the gas phase by electron diffraction, 191–8
Lewis-base adducts of group 1B metal(I) compounds. Part 17. Synthesis and crystal structures of adducts of copper(I) cyanide with nitrogen bases, 839–44
AMINO ACID
Enantioselectivity of nickel(II) and copper(II) complexes of Schiff bases derived from amino acids and (*S*)-*o*-[(*N*-benzylprolyl)amino]acetophenone or (*S*)-*o*-[(*N*-benzylprolyl)amino]benzaldehyde. Crystal and molecular structures of [Ni{(*S*)-bap-(*S*)-Val}] and [Cu{(*S*)-bap-(*S*)-Val}], 17–26
Glycine complexation with uranyl ion: absorptiometric, luminescence, and *X*-ray structural studies of tetrakis-(glycine)dioxouranium(VI) nitrate, 517–22
Studies on the anti-tumour activity of di- and tri-organotin(IV) complexes of amino acids and related compounds, of 2-mercaptoethanesulphonate, and of purine-6-thiol, 523–8
Electron spin resonance parameters for some copper(II)–bis(amino-acid) complexes, 1717–8
Synthesis and characterization of some chromium(III) complexes with glutathione, 2085–90
The effect of a dansyl group on the co-ordinative ability of *N*-protected amino acids. Part 1. Behaviour of the copper(II) ion–*N*-dansylglycinate system in aqueous and methanolic solution, 2363–8
AMINOETHYLAMMONIUM
Platinum(II) complexes containing a cationic amine ligand: crystal structure of [(2-aminoethyl)ammonium]trichloroplatinum(II), 1057–60
AMINOLYSIS
Studies of phosphazenes. Part 21. Associative and dissociative pathways in the aminolysis reactions of halogenocyclotriphosphazenes, 285–90
Studies of phosphazenes. Part 21. Associative and dissociative pathways in the aminolysis reactions of halogenocyclotriphosphazenes (1985, 285), 2459–60
AMMINE
Solvent dependence of the stereochemistry of base-catalysed solvolysis of *trans*-$[Co(NH_3)_4(^{15}NH_3)X]^{3+/2+}$ ions in dipolar aprotic solvents, 659–62
Kinetics and mechanism of the reductions of tris(oxalato)cobaltate(III) ion by ruthenium(II) species in aqueous solution, 1665–8
Linkage isomerism in penta-ammineruthenium(II),(III) complexes of benzotriazole, 2469–72
AMMONIA
Spectrochemistry of solutions. Part 16. A Raman spectroscopic study of the complexation of mercury(II) by cyanide ligands in liquid ammonia at 293 K, 663–8
AMMONIATION
Steric changes at labelled NH_2 sites during the base-catalysed ammoniation of the *trans*-azido(dimethyl sulphoxide)bis(ethylenediamine)cobalt(III) ion, 1495–500
Steric changes at labelled NH_2 sites during the base-catalysed ammoniation of the *trans*-azido(dimethyl sulphoxide)bis(ethylenediamine)cobalt(III) ion, 2223–4
AMMONIUM
Co-ordination of alkali metals by open-chain polyethers in transition metal complexes. Part 4. Variation in alkali-metal ion selectivity in cobalt and zinc complexes of 1-(*o*-carboxymethoxyphenoxy)-2-(*o*-hydroxyphenoxy)ethane (HL) and the *X*-ray and molecular structure of $[Zn(NH_4L_2)_2]$, 459–64
ANILINE
Transition-metal co-ordination compounds of a novel aniline-based pyrazole derivative. *X*-Ray crystal structures of [*NN*-bis(3,5-dimethylpyrazol-1-ylmethyl)aminobenzene]-dichlorocobalt(II) and -dibromocopper(II), 1699–706
ANTIMONY
Preparation and characterisation of adducts of bismuth pentafluoride and antimony pentafluoride by vibrational spectroscopy, *X*-ray powder diffraction, and single-crystal *X*-ray crystallography, 9–16
Toluene-3,4-dithiol (H_2tdt) complexes of Group 5B halides. Observations of lone-pair stereochemical activity and redox behaviour. Crystal and molecular structures of [AsCl(tdt)] and $[PPh_4][Sb(tdt)_3]$, 69–74
N,N'-Disubstituted dithiomalonamide complexes of antimony(III). Crystal and molecular structure of $[SbCl_3\{C_2H_5NHC(S)CH_2C(S)NHC_2H_5\}]$ with lone-pair occupation of an antimony co-ordination site, 1073–6
The preparation and crystal structure of (β)6$SbF_3\cdot 5SbF_5$, 1623–32
Synthesis and crystal structure of the layer compound Sb_3TeO_6Cl, 1633–6
Synthesis and characterisation of new mixed oxides of antimony and tellurium, 2225–30
ANTIMUSCARINIC AGENT
Sila-pharmaca. Part 32. Crystal and molecular structures of the

ANTIMUSCARINIC AGENT (contd)
(*R*)-enantiomer and the racemate of the antimuscarinic agent (cyclohexyl)phenyl[2-(pyrrolidin-1-yl)ethyl]silanol (sila-procyclidine), 1743–6
ANTI-TUMOUR ACTIVITY
Studies on the anti-tumour activity of di- and tri-organotin(IV) complexes of amino acids and related compounds, of 2-mercaptoethanesulphonate, and of purine-6-thiol, 523–8
ARENE
New carbonyl derivatives of niobium(I) and tantalum(I), 1989–96
Synthesis, structure, and bonding of fulvene complexes of titanium, molybdenum, and tungsten, 2037–50
ARSENATE
Preparation of 1,3,2-dithiazolium hexafluoroarsenate(V), preparation and crystal structures of 5-methyl-1,3,2,4-dithiadiazolium and 4-methyl-1,3,2-dithiazolium hexafluoroarsenate(V) and the reduction of these salts to stable free radicals, 1405–16
Crystalline zirconium(IV) hydrogenarsenate hydrogenphosphate monohydrate: synthesis, ion-exchange properties, and thermal behaviour, 1737–42
ARSENIC
Toluene-3,4-dithiol (H_2tdt) complexes of Group 5B halides. Observations of lone-pair stereochemical activity and redox behaviour. Crystal and molecular structures of [AsCl(tdt)] and $[PPh_4][Sb(tdt)_3]$, 69–74
Purification and *X*-ray crystal structure of bis[tris(trimethylsilyl)methyl]diarsene, 383–6
A disagreement on the explanation of short and long As–O bonds of the $(As–O)_4$ ring in $As_4(CF_3)_6O_6(OH)_2$ in terms of As^{III}–O and As^{V}–O bonds, 2221–2
ARSINE
Chemical and structural aspects of silver–triphenylarsine complexes and silver–tin complex salts, 321–32
Crystal structures and interrelationships of the blue and green conformational isomers of tetrakis(trimethylarsine sulphide)cobalt(II) perchlorate, 1289–94
Dirhodium(II,II) tetra-acetate complexes with axially co-ordinated triphenylstibine, triphenylarsine, and dibenzyl sulphide ligands. The syntheses, properties, and *X*-ray crystal structures of $[Rh_2(O_2CMe)_4(SbPh_3)_2]$, $[Rh_2(O_2CMe)_4(AsPh_3)_2]$, and $[Rh_2(O_2CMe)_4\{S(CH_2Ph)_2\}_2]$, 1775–80
Cationic complexes of ruthenium-(II) and -(III) with uni- and poly-dentate ligands, 2603–8
ARYL
From diarylruthenium complexes to *ortho*-metallated ketones: a mechanistic and crystal structure study, 1235–42
Synthetic and nuclear magnetic resonance studies on dialkyl- and diaryl-platinum complexes containing chelating, monodentate, or bridging $Ph_2PCH_2PPh_2$ ligands, 1501–6
Bimetallic systems. Part 11. Heterobimetallic and unsymmetrical diplatinum complexes from *cis*-$[PtR_2(dppm\text{-}P)_2]$ (dppm = $Ph_2PCH_2PPh_2$; R = Me, 1-naphthyl, or C_6H_4Me-*o*): crystal structure of $[(C_6H_4Me\text{-}o)_2Pt(\mu\text{-}dppm)_2PtMe_2]$, 1677–82
Reactivity of $[NiR(R')L_2]$ compounds and the crystal structure of $[Ni(C_2Cl_3)(C_6H_2Me_3\text{-}2,4,6)(PMe_2Ph)_2]$, 2333–42
ARYLSULPHONYL
Crystal structures and rotameric forms of some diarylsulphonyl-mono-, -di-, and -tri-selanes and their sulphur analogues, 231–8
ASSOCIATION
The peroxodisulphate–hexacyanoferrate(II) reaction. Reactivity and ionic association in isodielectric water–co-solvent mixtures, 1975–6
AZIDE
X-Ray crystal structures and magnetic properties of azide-bridged binuclear copper(II) complexes containing the Schiff-base ligand derived from 2-pyridinecarbaldehyde and histamine. Structure–magnetism relationship, 2095–100
AZIRIDINO
Studies of phosphazenes. Part 25. Synthesis, nuclear magnetic resonance spectroscopy, and mode of formation of (aziridino)(triphenylphosphazenyl)cyclotriphosphazenes. *X*-ray crystal structure and enzyme-inhibiting activity of $N_3P_3(NPPh_3)(NC_2H_4)_5$, 1881–90
BASE CATALYSIS
Steric changes at labelled NH_2 sites during the base-catalysed ammoniation of the *trans*-azido(dimethyl sulphoxide)bis(ethylenediamine)cobalt(III) ion, 1495–500
Steric changes at labelled NH_2 sites during the base-catalysed ammoniation of the *trans*-azido(dimethyl sulphoxide)bis(ethylenediamine)cobalt(III) ion, 2223–4
BENZAMIDINATE
Benzamidinatorhodium complexes. *X*-Ray structures of $[Rh\{CPh(NPh)_2\}(cod)]$ and $[Rh_2\{\mu\text{-}CPh(NPh)_2\}_2(tfbb)_2]$, 1487–94
BENZENETHIOLATE
Crystal structure and spectroscopic and redox properties of the iron–sulphur cluster compound $[NEt_4]_2[Fe_4S_4(SC_6H_4NH_2\text{-}4)_4]$, 2161–6
Rhenium nitrido-, arylimido-, nitrile, and carbonyl complexes with sterically hindered thiolate ligands, 2305–10
Complexes of molybdenum-(II) and -(IV) and tungsten(II) with sterically hindered thiolate ligands. Synthesis, reactivity, and *X*-ray crystal structures of $[PPh_4][Mo(SC_6H_2Pr^i_3\text{-}2,4,6)_3(CO)_2]$ and $[Mo(NNPh)(SC_6H_2Pr^i_3\text{-}2,4,6)_3(NCMe)]$, 2639–46
BENZIMIDAZOLYL
Crystal structures and magnetism of binuclear iron(III) complexes with a linear oxo-bridge, $[Fe_2O(bbimae)_2X_2][NO_3]_2$ {bbimae = 2-[bis(benzimidazol-2-ylmethyl)amino]ethanol, X = Cl or NCS}, 2375–80
BENZOATE
Crystal and molecular structure and magnetic properties of linear trimeric copper(II) complexes with predominant ferromagnetic exchange interaction, 2609–14
BENZODIPHOSPHOLE
Reaction between $[Ru_3(CO)_{12}]$ and the tridentate phosphine ligand $HC(PPh_2)_3$; *X*-ray crystal structures of complexes $[Ru_3(CO)_9\{Ph_2PCHP(Ph)C_6H_4PPh\}]$, $[Ru_2H(CO)_4(Ph_2PCHPPh_2)\{PhPC_6H_4C(O)\}]$, and $[Ru_2(CO)_4Cl(PPh_2)(dppm)]$, 1835–42
BENZOIN
Catalytic air oxidation of benzoin in the presence of dioxo-molybdenum(VI) complexes with sulphur chelate ligands, 635–40
BENZOTRIAZOLE
Linkage isomerism in penta-ammineruthenium(II),(III) complexes of benzotriazole, 2469–72
BENZOYL
Reaction between $[Ru_3(CO)_{12}]$ and the tridentate phosphine ligand $HC(PPh_2)_3$; *X*-ray crystal structures of complexes $[Ru_3(CO)_9\{Ph_2PCHP(Ph)C_6H_4PPh\}]$, $[Ru_2H(CO)_4(Ph_2PCHPPh_2)\{PhPC_6H_4C(O)\}]$, and $[Ru_2(CO)_4Cl(PPh_2)(dppm)]$, 1835–42
BENZYLAMINE
Kinetics and mechanism of ortho-palladation of ring-substituted *NN*-dimethylbenzylamines, 2629–38
BERYLLIUM
Beryllocene: A microwave dielectric loss study, 1761–6
BERYLLOCENE
Beryllocene: A microwave dielectric loss study, 1761–6
BIBICYCLO-OCTADIENE
Reduction–oxidation properties of organotransition-metal complexes. Part 20. Oxidative and thermolytic cyclopropane ring-opening reactions; *X*-ray crystal structure of $[Fe_2(CO)_6(\eta^4:\eta^{4}\text{-}C_{16}H_{18})]$, 699–706
BIPIPERIDINE
Optical resolution of DL-2,2′-bipiperidine through its cobalt(III) complex, 895–8
BIPYRIDINE
Light-induced electron-transfer reactions. Part 3. Kinetics of the oxidation reaction of oxalate ion by peroxodisulphate ion, induced by irradiation with visible light of an aqueous solution containing tris(2,2′-bipyridine)ruthenium(II), 355–60
The extraction and transport of metal ions by 6,6′-diamino-2,2′-bipyridine derivatives, 373–8
Cyclopalladated derivatives of 2,4′-bipyridine, 1719–22
Magnetic circular dichroism spectra of tris-chelate complexes of 2,2′-bipyridyl and 1,10-phenanthroline with iron(II), ruthenium(II), and osmium(II) at 4.2 K, 1781–8
New photoreduction catalysis by $[Cu(N\text{–}N)(PPh_3)_2]^+$ (N–N = 2,9-dimethyl-1,10-phenanthroline or 4,4′,6,6′-tetramethyl-2,2′-bipyridine) and its application to cobalt(III) complexes, 1959–62
Preparation and characterisation of 2,2′-bipyridine-4,4′-disulphonic and -5-sulphonic acids and their ruthenium(II) complexes. Excited-state properties and excited-state electron-transfer reactions of ruthenium(II) complexes containing 2,2′-bipyridine-4,4′-disulphonic acid or 2,2′-bipyridine-4,4′-dicarboxylic acid, 2247–62
Influence of decreasing solvent polarity (dioxane–water mixtures)

BIPYRIDINE (contd)
on the stability and structure of binary and ternary complexes of adenosine 5′-triphosphate and uridine 5′-triphosphate, 2291–304
BIPYRIDYL
Lewis-base adducts of Group 1B metal(I) compounds. Part 13. Crystal structure determinations of tetrakis(triphenyl-phosphine)-copper(I) and -silver(I) perchlorates, bis(pyridine)bis(triphenylphosphine)copper(I) perchlorate, (2,2′-bipyridyl)bis(triphenylphosphine)copper(I) perchlorate, and tetrahydroboratobis(triphenylphosphine)copper(I)–pyridine (1/0.5), 125–34
Electron-transfer reactions from *cis*-dialkylbis(2,2′-bipyridyl)cobalt(III) complexes to organic oxidants, 899–904
Actinide structural studies. Part 7. The crystal and molecular structures of (2,2′-bipyridyl)dinitratodioxo-uranium(VI) and -neptunium(VI), and diacetato(2,2′-bipyridyl)dioxo-uranium(VI) and -neptunium(VI), 1001–8
Absorption and emission in tris(2,2′-bipyridyl)ruthenium(II); effects of excited-state asymmetry, 1081–6
BIQUINOLINE
Characterization of the adducts formed by Cu(CN) and Cu(NCS) with biquinoline. The crystal structure of the polymeric cyano-compound containing both linear and tetrahedrally co-ordinated copper(I), $[\{Cu_3(bq)_2(CN)_3\}_n]$, 1285–8
BIS(DIPHENYLPHOSPHINO)METHANIDE
Synthesis of trinuclear gold(I) and gold(III) complexes containing the tridentate bis(diphenylphosphino)methanide ligand. Crystal structure of $[Cl(C_6F_5)_2Au\{Ph_2PCH(AuNC_5H_5)PPh_2\}AuCl]$, 2417–20
BISMUTH
Preparation and characterisation of adducts of bismuth pentafluoride and antimony pentafluoride by vibrational spectroscopy, *X*-ray powder diffraction, and single-crystal *X*-ray crystallography, 9–16
Kinetics and mechanism of electron-transfer reactions of bismuth(V) in aqueous acidic perchlorate–fluoride media. Part 1. Oxidation of hypophosphorous acid, 2571–4
BORANE
Photoelectron spectra of the aminodifluoroboranes NH_2BF_2, $NHMeBF_2$, and NMe_2BF_2, 1767–70
BORON
A kinetic study of the gas-phase thermolysis of hexaborane(10), 541–8
Carbaborane derivatives of the late- and post-transition elements. Part 3. Structural consequences of ligand substitution in palladadicarbadodecaboranes 3-L_2-3,1,2-$PdC_2B_9H_{11}$. The crystal and molecular structures of 3-$[Me_2N(CH_2)_2NMe_2]$-3,1,2-$PdC_2B_9H_{11}$ and 3-$(PMe_3)_2$-3,1,2-$PdC_2B_9H_{11}$, 761–70
Reactions of 6,6′-bis(*nido*-decaboranyl) oxide and 6-hydroxy-*nido*-decaborane with dihalogenobis(phosphine) complexes of nickel, palladium, and platinum, and some related chemistry; nuclear magnetic resonance investigations and the crystal and molecular structures of bis(dimethylphosphine)-di-μ-(2,3,4-η^3-*nido*-hexaboranyl)-diplatinum(*Pt–Pt*), $[Pt_2(\mu\text{-}\eta^3\text{-}B_6H_9)_2(PMe_2Ph)_2]$, and of 2,4-dichloro-1,1-bis(dimethylphenylphosphine)-*closo*-1-nickeladecaborane, $[(PhMe_2P)_2NiB_9H_7Cl_2]$, 953–72
Polyhedral rhenaborane chemistry: crystal and molecular structures of the *nido*-6-rhenadecaborane cluster compounds [6,6,6,6-$(PMe_2Ph)_3$H-*nido*-6-ReB_9H_{13}] and [2-(PMe_2Ph)-6,6,6,6-$(PMe_2Ph)_2$ClH-*nido*-6-ReB_9H_{12}]; nuclear magnetic resonance parameters of these and other related *nido*-rhenadecaborane cluster species, 1119–30
Studies of 2,5;6,10;8,10-tri-μ-hydro-nonahydro-*nido*-nonaborate(1−), $[B_9H_{12}]^-$: preparation, crystal and molecular structure, nuclear magnetic resonance spectra, electrochemistry, and reactions, 1645–54
Contributions to the chemistry of boron. Part 160. A convenient synthesis of catecholatoborane and diborane, 1689–92
Photoelectron spectra of the aminodifluoroboranes NH_2BF_2, $NHMeBF_2$, and NMe_2BF_2, 1767–70
Electronic structures of the *closo*-thiaboranes 1-SB_9H_9 and 1-$SB_{11}H_{11}$. Electrophilic substitution in 1-SB_9H_9, 2591–4
BORON-11
Bis(triphenylphosphine)copper(I) derivatives of substituted *arachno* nine-vertex borane anions, $Cu(PPh_3)_2(B_9H_{13}X)$ (X = H, NCS, NCSe, $NCBPh_3$, $NCBH_3$, or $NCBH_2NCBH_3$), 1903–8
BROMIDE
Lewis-base adducts of Group 1B metal(I) compounds. Part 11. Synthesis and crystal structure of adducts of silver(I) bromide with monomethyl-substituted pyridine bases, 111–6
Bromine nuclear quadrupole resonance studies of some hexabromostannates: *X*-Ray crystal structure of pyridinium hexabromostannate(IV), $[Hpy]_2[SnBr_6]$, 1399–404
Reactivity of $[NBu_4]_2[Mo_2Br_8]$ with several uni- and poly-dentate phosphines. *X*-Ray structure of $[NBu_4][MoBr_4(Ph_2PCH_2CH_2PPh_2)]$, 2263–8
BUTENYLPEROXO
Routes of formation and crystal structure of an alkylperoxycobaloxime: bis[dimethylglyoximato(1−)](4-ethoxy-carbonylbut-3-en-2-ylperoxo)(pyridine)cobalt(III), 1997–2000
BUTYL
Determination of the molecular structures of tri(t-butyl)phosphine oxide and tri(t-butyl)phosphine imide in the gas phase by electron diffraction, 827–30
CADMIUM
Electron spin resonance spectra of manganese(II) ions in the double-stranded chain polymers $CdLX_2$ (L = pyridine or methylpyridine; X = Cl or Br), 75–80
Thermochemical data for adducts of zinc, cadmium, and mercury halides with hexamethylphosphoramide, 1103–6
Unidentate *versus* symmetrically and unsymmetrically bidentate nitrate co-ordination in pyrazole-containing chelates. The crystal and molecular structures of (nitrato-*O*)[tris(3,5-dimethylpyrazol-1-ylmethyl)amine]copper(II) nitrate, (nitrato-*O*,*O*′)[tris(3,5-dimethylpyrazol-1-ylmethyl)amine]nickel(II)nitrate, and (nitrato-*O*)(nitrato-*O*,*O*′)[tris(3,5-dimethylpyrazol-1-ylmethyl)amine]cadmium(II), 2177–84
CAGE
Phosphorus–phosphorus bond cleavage in the cage molecule P_4S_3: synthesis and crystal structure of the trinuclear platinum complex $[\{Pt(\mu\text{-}P_4S_3)(PPh_3)\}_3]$-$C_6H_6$, 291–6
CALCIUM
Metal–phenoxyalkanoic acid interactions. Part 13. Copper(II)–(2-chlorophenoxy)ethanoic acid complexes. Crystal and molecular structures of *catena*-tetra-μ-[(2-chlorophenoxy)ethanoato-*O*,*O*′]-dicopper(II), *catena*-(2-aminopyrimidine-*N*,*N*′)-tetra-μ-[(2-chlorophenoxy)ethanoato-*O*,*O*′]-dicopper(II), and 1,2,2,2,2-penta-aqua-tetra-μ-[(2-chlorophenoxy)ethanoato-*O*,*O*′]-copper(II)calcium(II), 243–52
On the possibility of determining the thermodynamic parameters for the formation of weak complexes using a simple model for the dependence on ionic strength of activity coefficients: Na^+, K^+, and Ca^{2+} complexes of low molecular weight ligands in aqueous solution, 2353–62
CALORIMETRY
Zinc–sulphur bond enthalpy: its determination in bis(diethyldithiocarbamato)zinc(II), 369–72
CARBENE
Carbene complexes. Part 18. Synthetic routes to electron-rich olefin-derived monocarbenerhodium(I) neutral and cationic complexes and their chemical and physical properties (1984, 2355), 229–30
Sulphur-containing metal complexes. Part 14. Reactions of carbene anions with carbon disulphide or carbon diselenide, 1963–6
CARBIDE
New carbide clusters in the cobalt subgroup. Part 13. Synthesis and chemical characterization of the anions $[Co_6C(CO)_{14}]^-$, $[Co_6C(CO)_{15}]^{2-}$, and $[Co_8C(CO)_{18}]^{2-}$, and crystal structure of μ_6-carbido-ennea-μ-carbonyl-hexacarbonyl-*polyhedro*-hexacobaltate(2−) as its benzyltrimethylammonium salt; a comparison with isostructural species, 35–42
New carbide clusters in the cobalt sub-group. Part 14. Synthesis and structural characterization of the anion $[Co_{13}C_2(CO)_{24}]^{3-}$ as its benzyltrimethylammonium salt, 1137–40
New carbide clusters in the cobalt sub-group. Part 15. Synthesis and crystallographic characterization of di-μ_6-carbido-deca-μ-carbonyl-tridecacarbonyl-*polyhedro*-dodecarhodate(4−) as its tetrapropylammonium salt, $[N(C_3H_7)_4]_4$-$[Rh_{12}C_2(CO)_{23}]$, 1309–14
Reactions of some decaosmium clusters with electrophilic and nucleophilic reagents: *X*-ray structure analyses of $[N(PPh_3)_2][Os_{10}C(CO)_{24}(\mu\text{-}I)]$, $[Os_{10}C(CO)_{24}(\mu\text{-}I)_2]$, $[N(PPh_3)_2]_2[Os_{10}C(CO)_{22}(NO)I]$, $[Os_{10}C(CO)_{23}\{P(OMe)_3\}(\mu\text{-}I)_2]$ and of two isomers of $[Os_{10}C(CO)_{21}\{P(OMe)_3\}_4]$, 1795–810
Synthesis of tetra- and penta-nuclear platinum–osmium carbido-cluster complexes from non-carbido-precursors: *X*-ray structural studies on $[Os_3Pt(\mu\text{-}H)_2(\mu_4\text{-}C)(CO)_{10}\{P(cyclo\text{-}C_6H_{11})_3\}]$, $[Os_3Pt_2(\mu\text{-}H)_2(\mu_5\text{-}C)(\mu\text{-}CO)(CO)_9\{P(cyclo\text{-}C_6H_{11})_3\}_2]$, and

CARBIDE (contd)
$[Os_3Pt_2(\mu\text{-}H)(\mu_3\text{-}C)(\mu\text{-}OMe)(\mu\text{-}CO)(CO)_9\{P(cyclo\text{-}C_6H_{11})_3\}_2]$, 2437–48

CARBON-13
Two-dimensional nuclear magnetic resonance for the analysis of the carbon-13 spectra of carbonyl groups in metallocarbonyls, 225–8
Mercurated and tellurated Schiff bases and phenylhydrazones, 821–6
Geometry-dependent complexation effects in carbon-13 nuclear magnetic resonance spectra of tricarbonyl(3—8-η-[2.2]paracyclophane)chromium and related complexes, 1065–8
Geometry-dependent carbon-13 chemical shifts in (η⁶-[2.2]cyclophane)(η⁵-cyclopentadienyl)iron(II) hexafluorophosphates, 1661–4
Carbon-13 nuclear magnetic resonance evidence of a relaxation process dominated by scalar coupling with a quadrupolar nucleus in $[Re_3(\mu\text{-}H)_4(CO)_{10}]^-$, 1899–902
Carbon-13 nuclear magnetic resonance study of the complexes formed between zinc(II) and triethylenetetramine, 2381–6

CARBONATE
Chemistry of metallacyclobutanones. Part 3. Reactions of heptane-2,4,6-trione and 1,5-diphenylpentane-1,3,5-trione with some carbonate complexes of platinum(II); *X*-ray crystal structures of $[Pt\{CH(COMe)COCH(COMe)\}(PPh_3)_2]$ and $[Pt\{OC(CHCOPh)CHC(Ph)O\}(PPh_3)_2]$, 549–54

CARBON DIOXIDE
Insertion of carbon dioxide, of CO_2-like molecules, and of other unsaturated compounds into the platinum–nitrogen bond of $[Pt(PPh_3)_2(PhNO)]$, 163–8

CARBON DISELENIDE
Sulphur-containing metal complexes. Part 14. Reactions of carbene anions with carbon disulphide or carbon diselenide, 1963–6

CARBON DISULPHIDE
Bimetallic systems. Part 10. Synthesis of complexes of type $[(RC{\equiv}C)Pt(\mu\text{-}dppm)_2Pt(C{\equiv}CR)]$ (dppm = $Ph_2PCH_2PPh_2$, R = Ph or *p*-tolyl) and their corresponding 'A frames' $[(RC{\equiv}C)Pt(\mu\text{-}dppm)_2(\mu\text{-}H)Pt(C{\equiv}CR)]Cl$ or $[(RC{\equiv}C)Pt(\mu\text{-}dppm)_2(\mu\text{-}X)Pt(C{\equiv}R)]$ with X = CS_2 or MeOOCC≡CCOOMe, 1015–8
Sulphur-containing metal complexes. Part 14. Reactions of carbene anions with carbon disulphide or carbon diselenide, 1963–6
Fragmentation of co-ordinated carbon disulphide and dithiocarbene ligands by nucleophilic attack at carbon: the crystal structure of $[(Ph_3P)IPt(\mu\text{-}SMe)(\mu\text{-}CSMe)PtI(PPh_3)]\cdot Me_2CO$, 2595–602

CARBON MONOXIDE
Rate parameters for oxygen and carbon monoxide binding to a liposome-embedded heme under physiological conditions, 65–8
Equilibrium and kinetic study of the reaction between phthalocyaninatoiron(II) and carbon monoxide in dimethyl sulphoxide in the presence of pyridine. Evidence for the formation of a transient, 1113–8
The easy catalytic reduction of $HClO_4$, HNO_3, and H_5IO_6 by CO in the presence of $[Pd_3(O_2CMe)_6]$, 1971–4
Selective conversion of CO into methanol at ordinary temperature. Part 4. Activation by iron(II), iron(III), and chromium(III) complexes, 2499–504

CARBONYL
New carbide clusters in the cobalt subgroup. Part 13. Synthesis and chemical characterization of the anions $[Co_6C(CO)_{14}]^-$, $[Co_6C(CO)_{15}]^{2-}$, and $[Co_8C(CO)_{18}]^{2-}$, and crystal structure of μ_6-carbido-ennea-μ-carbonyl-hexacarbonyl-*polyhedro*-hexacobaltate(2–) as its benzyltrimethylammonium salt; a comparison with isostructural species, 35–42
Metallation of 2-ethenylpyridine at triosmium clusters: *X*-ray crystal structures of the open trinuclear clusters $[Os_3H(CO)_9L(NC_5H_4CH{=}CH)]$ (L = CO or PMe_2Ph), 85–90
Reaction of bis-μ-diethylphosphido-bis(tetracarbonylmetal) (*M–M*) (M = Cr or W) with tri-n-butylphosphine: kinetics and mechanism of a reaction involving seven-co-ordinate complexes, 91–8
Chemistry of the unsaturated cluster compound $[Os_3Pt(\mu\text{-}H)_2(CO)_{10}\{P(cyclo\text{-}C_6H_{11})_3\}]$; *X*-ray crystal structures of $[Os_3Pt(\mu\text{-}H)_2(CO)_{11}\{P(cyclo\text{-}C_6H_{11})_3\}]$, $[Os_3Pt(\mu\text{-}H)_4(CO)_{10}\{P(cyclo\text{-}C_6H_{11})_3\}]$, and $[Os_3Pt(\mu\text{-}H)_2(\mu\text{-}CH_2)(CO)_{10}\{P(cyclo\text{-}C_6H_{11})_3\}]$ (two isomers), 177–90
Two-dimensional nuclear magnetic resonance for the analysis of the carbon-13 spectra of carbonyl groups in metallocarbonyls, 225–8
The preparation and hydridic reduction of dicationic dicarbonyl complexes of osmium(II); the crystal and molecular structure of *trans*-bis[1,2-bis(diphenylphosphino)ethane-*PP'*]carbonylformylosmium(II) hexafluoroantimonate–dichloromethane (1/1), 387–94
Systematic synthesis of tetranuclear osmium clusters by the reaction of trinuclear clusters with $[OsH_2(CO)_4]$; crystal structure of $[Os_4H_3Br(CO)_{13}]$, 555–64
Synthesis and *X*-ray structural studies on the cluster compounds $[RuRh_3(\mu_3\text{-}CO)_2(CO)_3(\eta\text{-}C_5Me_5)_3]$ and $[RuRh_2(\mu\text{-}CO)(\mu_3\text{-}CO)(CO)_2(\eta^4\text{-}C_8H_{10})(\eta\text{-}C_5Me_5)_2]$, 645–50
Synthesis of $[Fe(\eta^4\text{-}C_8H_8)(CO)_2L]$ [L = MeNC, PriNC, ButNC, PhNC, 2,6-$Me_2C_6H_3NC$, or $P(OMe)_3$] and proof of the applicability of the Woodward–Hoffmann rules to the fluxionality of $[Fe(\eta^4\text{-}C_8H_8)(CO)_2(CNPr^i)]$, 693–8
Reduction–oxidation properties of organotransition-metal complexes. Part 20. Oxidative and thermolytic cyclopropane ring-opening reactions; *X*-ray crystal structure of $[Fe_2(CO)_6(\eta^4\text{:}\eta'^4\text{-}C_{16}H_{18})]$, 699–706
Electrochemistry of clusters. Part 4. Redox behaviour of tetrametallic clusters $[M^1{}_2M^2{}_2(\eta^5\text{-}C_5H_5)_2(\mu_3\text{-}CO)_2(\mu\text{-}CO)_4(PR_3)_2]$ (M^1 = Pt or Pd; M^2 = Cr, Mo, or W; R = Me, Et, Bu^n, or Ph), 711–6
The addition of protic acids to $[Mn_2(CO)_8(Ph_2PCH_2PPh_2)_2]$ to give bridging hydrido-compounds, 743–8
Annelation of ring-opened arylcyclopropenium ions to co-ordinated cyclo-octatetraene, and the *X*-ray crystal structure of $[Fe(CO)_3(\sigma,\eta^3\text{-}C_{11}H_9Ph_3)]$, 777–82
Formation of substituted cyclopentadienyl ligands on tungsten *via* reactions between the alkyne complexes $[W(CO)(R^1C_2R^2)_3]$ ($R^1 = R^2$ = Ph or Et; R^1 = Me, R^2 = Ph) and the alkylidyne compounds $[W(\equiv CR)(CO)_2(\eta\text{-}C_5H_5)]$ (R = C_6H_4Me-4 or Me); *X*-ray crystal structures of $[W_2(\mu\text{-}CO)_2(CO)(\eta\text{-}PhC_2Ph)(\eta\text{-}C_5Ph_4R)(\eta\text{-}C_5H_5)]$ and $[W_2(\mu\text{-}EtC_2Et)(CO)_4(\eta\text{-}C_5Et_4R)(\eta\text{-}C_5H_5)]$ (R = C_6H_4Me-4), 905–12
Reaction of transition-metal carbonylate anions and 1,1,1-tris(halogenomethyl)ethane. *X*-Ray crystal structures of tricarbonyl(η⁵-cyclopentadienyl)(1-methylcyclopropylmethyl)tungsten(II), and tetraethylammonium enneacarbonyliododirhenate(O), 931–40
Pyrazolate A-frame rhodium complexes. Crystal structures of $[Rh_2(\mu\text{-}dmpz)(CO)_2(\mu\text{-}dppm)_2][ClO_4]$ and $[Rh_2(\mu\text{-}dmpz)I_2(CO)_2(\mu\text{-}dppm)_2][ClO_4]$, 973–80
Bimetallic systems. Part 9. The synthesis of and nuclear magnetic resonance studies on 10-membered ring complexes of type $[(OC)_4M^1(\mu\text{-}Ph_2PCH_2CH_2PPh_2)_2M^2(CO)_4]$ (M^1, M^2 = Cr, Mo, or W), 1009–14
Reduction–oxidation properties of organotransition-metal complexes. Part 21. Synthesis and *X*-ray structural characterisation of the redox-related pair of cyclohexadienyl complexes $[Mn(CO)(dppe)(\eta^5\text{-}C_6H_6Ph)]$ and $[Mn(CO)(dppe)(\eta^5\text{-}C_6H_6Ph)][PF_6]\cdot 0.5CH_2Cl_2$, 1019–26
Reduction–oxidation properties of organotransition-metal complexes. Part 22. Stereospecific oxidative cyclopropane ring opening and reductive cyclobutane ring formation in polycyclic hydrocarbon complexes of iron; *X*-ray crystal structures of $[Fe_2(CO)_6(\eta^5\text{:}\eta'^5\text{-}C_{16}H_{16})][PF_6]_2\cdot CH_3NO_2$ and $[Fe_2(CO)_6(\eta^4\text{:}\eta'^4\text{-}C_{16}H_{16})]$, 1027–36
New isomers of $[Os_3(CO)_{10}(PMe_2Ph)_2]$ and $[Os_3(CO)_9(PMe_2Ph)_3]$, 1037–42
Mechanism of addition of aryltrimethyl-silanes and -stannanes to tricarbonyl(cyclohexadienyl)ruthenium(II) cation, 1049–52
Geometry-dependent complexation effects in carbon-13 nuclear magnetic resonance spectra of tricarbonyl(3—8-η-[2.2]paracyclophane)chromium and related complexes, 1065–8
Carbon–carbon formation at di-iron centres. Part 2. Reactivity of $[Fe_2(CO)_6(\mu\text{-}COEt)\{\mu\text{-}C(R)C(R)H\}]$ complexes toward MeOC(O)C≡CC(O)OMe (R = Ph) and $CF_3C{\equiv}CCF_3$ (R = Ph or H); *X*-ray crystal structures of $[Fe_2(CO)_5\{\mu\text{-}C(OEt)C[C(O)OMe]C[C(O)OMe]\}\{\mu\text{-}C(Ph)C(Ph)H\}]\cdot H_2O$ and $[Fe_2(CO)_6\{\mu\text{-}C(CF_3)C(CF_3)CHCHCH(OMe)\}]$, 1087–94
New carbide clusters in the cobalt sub-group. Part 14. Synthesis and structural characterization of the anion $[Co_{13}C_2(CO)_{24}]^{3-}$ as its benzyltrimethylammonium salt, 1137–40
Isolated CH stretching frequencies, methyl group geometry, and methyl CH bond lengths and strengths in tricarbonyl(η⁵-cyclopentadienyl)methylchromium(II), -molybdenum(II), and -tungsten(II), 1207–12
A mechanistic study on complexes of type *mer*-$[Cr(CO)_3(\eta^2\text{-}L\text{–}L)(\sigma\text{-}L\text{–}L)]$ (where L–L = $Ph_2PCH_2PPh_2$, $Ph_2PNHPPh_2$, or $Ph_2PNMePPh_2$) using spectroscopic and convolutive

CARBONYL (contd)
electrochemical techniques, 1213–22
Cluster chemistry. Part 31. An $[Ru_3(CO)_{12}]$ derivative with an $[Fe_3(CO)_{12}]$-like structure: preparation and *X*-ray structure of $[Ru_3(\mu\text{-}CO)_2(CO)_8\{PPh(OMe)_2\}_4]$, 1223–8
Cluster chemistry. Part 32. Synthesis and *X*-ray crystal structure of $[Ru_5(\mu_5\text{-}\eta^2\text{-}C_2PPh_2\text{-}P)(\mu\text{-}PPh_2)(CO)_{13}]$, a complex containing an alkynyl ligand in extended interaction with an open Ru_5 cluster, 1229–34
Electron-transfer reactions in nitrogen fixation. Part 1. The electrosynthesis of dinitrogen, hydride, isocyanide, and carbonyl complexes of molybdenum: intermediates, mechanisms, and energetics, 1255–64
Synthesis, structures, and reactivities of some pentamethylcyclopentadienyl–sulphur compounds, 1303–8
New carbide clusters in the cobalt sub-group. Part 15. Synthesis and crystallographic characterization of di-μ_6-carbido-deca-μ-carbonyl-tridecacarbonyl-*polyhedro*-dodecarhodate(4−) as its tetrapropylammonium salt, $[N(C_3H_7)_4]_4\text{-}[Rh_{12}C_2(CO)_{23}]$, 1309–14
Chemistry of di- and tri-metal complexes with bridging carbene or carbyne ligands. Part 33. Reactions of $[W(\equiv CMe)(CO)_2(\eta\text{-}C_5H_5)]$ with the dimetal compounds $[MRh(\mu\text{-}CO)_2(\eta\text{-}C_5Me_5)_2]$ (M = Co or Rh); *X*-ray crystal structure of $[Rh_2W(\mu\text{-}CO)(\mu_3\text{-}CMe)(CO)_2(\eta\text{-}C_5H_4)(\eta\text{-}C_5Me_5)_2]$, 1315–22
Chemistry of di- and tri-metal complexes with bridging carbene or carbyne ligands. Part 34. Reactions of sulphur and selenium with the di-irontungsten complexes $[Fe_2W(\mu_3\text{-}CR)(\mu\text{-}CO)(CO)_8(\eta\text{-}C_5H_5)]$ (R = C_6H_4Me-4 or Me); crystal structures of $[Fe_2W(\mu\text{-}CC_6H_4Me\text{-}4)(\mu_3\text{-}S)(CO)_7(\eta\text{-}C_5H_5)]$ and $[Fe_2W(\mu_3\text{-}SCMe)(CO)_8(\eta\text{-}C_5H_5)]$, 1323–30
Chemistry of di- and tri-metal complexes with bridging carbene or carbyne ligands. Part 35. The synthesis and crystal structures of the compounds $[N(PPh_3)_2][W_2\{\mu\text{-}C(H)C_6H_4Me\text{-}4\}(CO)_7(\eta\text{-}C_5H_5)]\text{-}0.5Et_2O$ and $[W\{=C(H)C_6H_4Me\text{-}4\}\text{-}(SnPh_3)(CO)_2(\eta\text{-}C_5H_5)]$, 1331–8
The reaction of $[Os_3(\mu\text{-}H)_2(CO)_9L]$ (L = CO or PEt_3) with dimethylcyanamide, Me_2NCN: *X*-Ray crystal structure of $[Os_3(\mu\text{-}H)(\mu\text{-}NCHNMe_2)(CO)_{10}]$ and the reactions of this complex with acids, 1355–60
Photochemistry of dicarbonyl(η^5-cyclopentadienyl)-methyl- and -ethyl-iron and -ruthenium complexes in solutions at −30 °C and in frozen gas matrices at 12 K, 1365–74
Reactivity of the unsaturated anion decacarbonyltetra-μ-hydrido-trirhenate(1−) toward phenols. Crystal and molecular structures of the tetraethylammonium salts of the triangular cluster anion $[Re_3(\mu\text{-}H)_3(\mu\text{-}OC_6F_5)(CO)_{10}]^-$ and of the binuclear anion $[Re_2(\mu\text{-}OC_6H_5)_3(CO)_6]^-$, 1507–12
Pyramidal inversions and 1,2-metal shifts in pentacarbonyl-chromium, -molybdenum, and -tungsten derivatives of dialkyl disulphides and dialkyl diselenides. A nuclear magnetic resonance investigation, 1561–8
Dynamic nuclear magnetic resonance study of Group 6 metal pentacarbonyl complexes. Pyramidal inversion and 1,2-metal shifts in the complexes $[M(CO)_5(Me_2\overline{CCH_2EECH_2})]$ (M = Cr, Mo, or W; E = S or Se), 1569–76
Synthesis of bis[μ-bis(diphenylphosphino)methane]-tri-μ-carbonyl-tricarbonyl-*triangulo*-trirhodium(1+) perchlorate, $[Rh_3(CO)_3(\mu\text{-}CO)_3(\mu\text{-}dppm)_2]ClO_4$. An unusual 46-electron cluster possessing the A-frame structure with a bridging $Rh(CO)_3$ fragment, 1577–84
Addition of proton and boron trifluoride at terminal and bridging cyano-ligands in dinuclear manganese compounds, 1609–12
Synthesis and structural characterisation of some *triangulo*-platinum clusters containing isocyanide ligands, 1693–8
Studies on the regioselective preparation of tricarbonyl(cyclohexadienyl)iron(1+) salts, 1723–6
Reactions of some decaosmium clusters with electrophilic and nucleophilic reagents: *X*-ray structure analyses of $[N(PPh_3)_2][Os_{10}C(CO)_{24}(\mu\text{-}I)]$, $[Os_{10}C(CO)_{24}(\mu\text{-}I)_2]$, $[N(PPh_3)_2]_2[Os_{10}C(CO)_{22}(NO)I]$, $[Os_{10}C(CO)_{23}\{P(OMe)_3\}(\mu\text{-}I)_2]$ and of two isomers of $[Os_{10}C(CO)_{21}\{P(OMe)_3\}_4]$, 1795–810
Reactions of hydride reagents with alkylmolybdenum carbonyl complexes. Reaction of $LiBHEt_3$ with $[MoMe(CO)_3(\eta\text{-}C_5H_5)]$, formation of an anionic acetaldehyde complex, and a stoicheiometric cycle for the synthesis of acetaldehyde, 1815–20
Reaction between $[Ru_3(CO)_{12}]$ and the tridentate phosphine ligand $HC(PPh_2)_3$; *X*-ray crystal structures of complexes $[Ru_3(CO)_9\{Ph_2PCHP(Ph)C_6H_4PPh\}]$, $[Ru_2H(CO)_4(Ph_2PCHPPh_2)\{PhPC_6H_4C(O)\}]$, and $[Ru_2(CO)_4Cl(PPh_2)(dppm)]$, 1835–42
1*H*-pyrrolo[2,3-*b*]pyridine (HL) ligands in rhodium(I) and iridium(I) chemistry. Crystal and molecular structures of $[Rh_2(\mu\text{-}L)_2(nbd)_2]$ and $[Rh_4(\mu\text{-}Cl)_2(\mu\text{-}L)_2(\mu\text{-}CO)_2(CO)_2(nbd)_2]$, 1891–8
Carbon-13 nuclear magnetic resonance evidence of a relaxation process dominated by scalar coupling with a quadrupolar nucleus in $[Re_3(\mu\text{-}H)_4(CO)_{10}]^-$, 1899–902
Organic chemistry of dinuclear metal centres. Part 8. Organo-iron–ruthenium chemistry. *X*-Ray structure of *trans*-$[FeRu(CO)_2(\mu\text{-}CO)_2(\eta\text{-}C_5H_5)_2]$, 1935–44
Sulphur-containing metal complexes. Part 14. Reactions of carbene anions with carbon disulphide or carbon diselenide, 1963–6
Reactivity of $[PPh_4][Fe_2(CO)_6\{\mu\text{-}CPhCPhC(CF_3)C(CF_3))H\}]$ toward electrophiles. *X*-Ray structure of $[Fe_2(CO)_6\text{-}\{\mu\text{-}CPhCPhC(CF_3)CHC(OEt)_2\}]$, a product resulting from fluorine abstraction at a CF_3 group and subsequent fluorine substitution by ethoxy-groups, 1981–4
New carbonyl derivatives of niobium(I) and tantalum(I), 1989–96
Chemistry of di- and tri-metal complexes with bridging carbene or carbyne ligands. Part 36. Reactions of the dimetal compounds $[ReM(\equiv CC_6H_4Me\text{-}4)(CO)_9]$ (M = Cr, Mo, or W) with octacarbonyldicobalt; crystal structures of $[Co_2WRe(\mu_3\text{-}CC_6H_4Me\text{-}4)(CO)_{15}]$ and $[Co_2Re(\mu_3\text{-}CC_6H_4Me\text{-}4)(CO)_{10}]$, 2001–8
Chemistry of di- and tri-metal complexes with bridging carbene or carbyne ligands. Part 37. Methylene group transfer to carbon–metal multiple bonds; crystal structures of $[TiW\{\mu\text{-}C(C_6H_4Me\text{-}4)\text{=}CH_2\}(\mu\text{-}CO)(CO)(\eta\text{-}C_5H_5)_3]$ and $[PtW\{\mu\text{-}C(C_6H_4Me\text{-}4)\text{=}CH_2\}(CO)_2(PMe_3)_2(\eta\text{-}C_5H_5)]$, 2009–16
Chemistry of di- and tri-metal complexes with bridging carbene or carbyne ligands. Part 38. Ruthenium–tungsten compounds: crystal structures of $[RuW(\mu\text{-}Cl)(\mu\text{-}CMe)(Cl)(CO)_2(PPh_3)_2(\eta\text{-}C_5H_5)]$ and $[RuW_2(\mu_3\text{-}C_2Me_2)(CO)_7(\eta\text{-}C_5H_5)_2]$, 2017–24
Bimetallic systems. Part 13. Platinum–manganese carbonyl complexes containing bridging $Ph_2PCH_2PPh_2$(dppm) ligands: crystal structure of $[(OC)_3Mn(\mu\text{-}dppm)_2PtH(Br)]BF_4$, 2131–8
Pyramidal inversion in configurational isomers of tetracarbonyl[1,1,2,2-tetrakis(methylthio)ethane]chromium(0): a two-dimensional nuclear magnetic resonance exchange study, 2195–202
Co-ordinatively unsaturated diene complexes of tungsten(II) and their reactions with nucleophiles to give six- and seven-co-ordinate derivatives, 2231–8
Synthesis and nuclear magnetic resonance studies of co-ordinatively unsaturated alkyne complexes of tungsten(II), 2239–46
The palladium-catalysed reaction between $[Re_2(CO)_{10}]$ and phosphines and the crystal and molecular structure of diaxial $[Re_2(CO)_8(PMe_2Ph)_2]$, 2277–82
Synthesis of tetra- and penta-nuclear platinum–osmium carbido-cluster complexes from non-carbido-precursors: *X*-ray structural studies on $[Os_3Pt(\mu\text{-}H)_2(\mu_4\text{-}C)(CO)_{10}\{P(cyclo\text{-}C_6H_{11})_3\}]$, $[Os_3Pt_2(\mu\text{-}H)_2(\mu_5\text{-}C)(\mu\text{-}CO)(CO)_9\{P(cyclo\text{-}C_6H_{11})_3\}_2]$, and $[Os_3Pt_2(\mu\text{-}H)(\mu_5\text{-}C)(\mu\text{-}OMe)(\mu\text{-}CO)(CO)_9\{P(cyclo\text{-}C_6H_{11})_3\}_2]$, 2437–48
Acid-induced displacement of acetaldehyde from a μ-vinyloxy-triosmium cluster, 2479–82
Reactions of co-ordinated ligands. Part 35. Evidence for carbon–carbon double-bond cleavage of cyclopropenes in their reaction with dinuclear cobalt, rhodium, and iridium complexes; crystal structure and protolysis of $[Rh_2(\mu\text{-}CO)(\mu\text{-}COCHCMe_2CH)(\mu\text{-}C_5Me_5)_2]$, 2483–92
The $[Fe_3(\mu_3\text{-}CR)(CO)_{10}]^-$ cluster anions as building blocks for the synthesis of mixed-metal clusters. Part 1. Synthesis of mixed clusters $[MFe_3(\mu_3\text{-}CMe)(CO)_{10}(PPh_3)]$ (M = Cu or Au) and crystal structure of $[CuFe_3(\mu_3\text{-}CMe)(CO)_{10}(PPh_3)]$, 2521–4
Interactions of $[Mo(CO)_6]$ and $[Mo(CO)_3(C_6H_5CH_3)]$ with O_2N_2- and O_2N_3-donor macrocycles and the *X*-ray crystal structure of tetracarbonyl(6,7,16,17-tetrahydro-15*H*-dibenzo[*e*,*n*][1,4,8,12]dioxadiazacyclopentadecine)molybdenum–toluene (2/1), 2561–4
He I and He II photoelectron spectra of open-chain pentadienyl complexes of manganese and rhenium, 2677–82

CARBOXYLATE
Complexes of the platinum metals. Part 23. Synthesis of the nitrosyl carboxylate complexes $[M(O_2CR)_2(NO)(PPh_3)_2]$ (M = Rh or Ir; R = CF_3, C_2F_5, or C_6F_5): crystal and molecular structures of the trifluoroacetate derivatives $[M(O_2CCF_3)_2(NO)(PPh_3)_2]$,

CARBOXYLATE (contd)
611–6
Complexes of the platinum metals. Part 26. Multinuclear nuclear magnetic resonance studies on rhodium(II) carboxylate adducts, 629–34
Complexes of the platinum metals. Part 30. Fragmentation reactions of rhodium and iridium trichloro- and tribromo-acetates, 2113–20
The synthesis, magnetic, electrochemical, and spectroscopic properties of diruthenium(II,II) tetra-μ-carboxylates and their adducts. *X*-Ray structures of $Ru_2(O_2CR)_4L_2$ (R = Me, L = H_2O or tetrahydrofuran; R = Et, L = Me_2CO), 2321–6
CATALYSIS
Heteropolytungstates as catalysts for the photochemical reduction of oxygen and water, 395–400
Kinetic and structural investigations of $[Fe^{III}(edta)]^-$ [edta = ethylenediaminetetra-acetate(4−)] catalysed decomposition of hydrogen peroxide, 493–502
Catalytic air oxidation of benzoin in the presence of dioxomolybdenum(VI) complexes with sulphur chelate ligands, 635–40
Mono-η-cycloheptatrienyltitanium chemistry: synthesis, molecular and electronic structures, and reactivity of the complexes $[Ti(\eta\text{-}C_7H_7)L_2X]$ (L = tertiary phosphine, O- or N-donor ligand; X = Cl or alkyl), 669–84
Studies on cyclic bis(η^5:σ-2-cyclopentadienylidene-ethyl)- and bis(η^5:σ-4-cyclopentadienylidenebutyl)-molybdenum compounds, 1585–90
Oxygenation studies. Part 7. Catalytic dioxygenation of cyclo-octa-1,5-diene at a rhodium centre, 1591–6
The chemistry and catalytic properties of ruthenium and osmium complexes. Part 1. Homogeneous catalysis of organic reactions by bromo(carbonyl)hydridotris(triphenylphosphine)osmium(II), 1859–64
Crystal structures and magnetic properties of binuclear five-co-ordinate copper(II) complexes with a phenolate bridge and their catalytic functions in multielectron redox reactions, 1945–52
New photoreduction catalysis by $[Cu(N\text{–}N)(PPh_3)_2]^+$ (N–N = 2,9-dimethyl-1,10-phenanthroline or 4,4′,6,6′-tetramethyl-2,2′-bipyridine) and its application to cobalt(III) complexes, 1959–62
The easy catalytic reduction of $HClO_4$, HNO_3, and H_5IO_6 by CO in the presence of $[Pd_3(O_2CMe)_6]$, 1971–4
The palladium-catalysed reaction between $[Re_2(CO)_{10}]$ and phosphines and the crystal and molecular structure of diaxial $[Re_2(CO)_8(PMe_2Ph)_2]$, 2277–82
Selective conversion of CO into methanol at ordinary temperature. Part 4. Activation by iron(II), iron(III), and chromium(III) complexes, 2499–504
CATECHOLATE
Contributions to the chemistry of boron. Part 160. A convenient synthesis of catecholatoborane and diborane, 1689–92
CATENADI(AZATHIANE)
Reaction of the anion $[WCl_4(CBu^t)]^-$ with tetrasulphur tetranitride. Formation and crystal structure of $[AsPh_4][WCl_3O(OS_2N_2)]$, 2453–6
C.D.
Correlation between structure and circular dichroism. Structure and absolute configuration of the (−) isomer of lithium (ethylenediamine-*N*,*N*′-diacetato-*N*,*N*′-di-3-propionato)rhodate(III) pentahydrate, 861–4
CHARGE TRANSFER
Synthesis and characterization of a new family of binuclear copper(II) complexes with a flexible bridge and the π–π type charge-transfer interaction with 1,3,5-trinitrobenzene, 107–10
Reaction of bis(morpholinothiocarbonyl) disulphide with iodine. Existence of a 1:1 charge-transfer precursory adduct in an oxidation reaction. Isolation and crystal structure of bis[3,5-di(*N*-morpholinio)-1,2,4-trithiolane] hexadecaiodide, 1349–54
CHEMICAL POTENTIAL
Transfer chemical potentials for Fe^{2+} from water into aqueous methanol: a comment, 867–6
Transfer chemical potentials for Fe^{2+} from water into aqueous methanol: a reply, 867–70
CHIRALITY
Co-ordinative interactions in chelated complexes of silicon. Part 5. Chiralty of five-co-ordinate silicon compounds: crystal and molecular structures of 1-(chlorodimethylsilyl)-and 1-(dichloromethylsilyl)-1,2,3,4-tetrahydro-1,10-phenanthroline, 1–4
Macrocyclic polyphosphane ligands. Cobalt(II) and nickel(II) complexes of the γ and ε diastereoisomers of 4,7,13,16-tetraphenyl-4,7,13,16-tetraphospha-1,10-dithiacyclo-octadecane (L^1) and the crystal structure of $[Co(\varepsilon\text{-}L^1)][BPh_4]_2 \cdot EtOH$, 1425–30
CHLORIDE
The kinetics of the displacement, by chloride, of heterocyclic nitrogen bases (am) from *trans*-$[PtL(am)Cl_2]$ (L = C_2H_4, CO, or PMe_3). The effect of steric hindrance in the leaving group on the *trans* effect of L, 27–30
Sulphur ligand–metal complexes. Part 16. Copper complexes of thioethers and the single-crystal *X*-ray structure of the polymeric mixed-valence complex, penta-μ-chloro-tris-μ-tetrahydrothiophene-tetracopper(I,II), 151–8
X-Ray crystal structures and properties of tris(tetrathiafulvalenium) tetrachlorodimethylstannate(IV) and tetrathiafulvalenium trichlorodimethylstannate (IV), 465–70
Displacement of pyridine-2-methanol from dichloro(pyridine-2-methanolato)gold(III) in acidic solution. Ring opening at oxygen, 731–6
Electrochemically induced ligand substitutions on $[OsCl_3(PMe_2Ph)_3]$: rational pathways to osmium(II) complexes, 947–52
Dinuclear mono-μ-chloro-pyridyldiaza rhodium(I) complexes derived from pyridyldi-imines *via* hydrogen transfer from ethanol, 1053–6
Platinum(II) complexes containing a cationic amine ligand: crystal structure of [(2-aminoethyl)ammonium]trichloroplatinum(II), 1057–60
Preparation and crystal structure of $Na_4[Hg_4(\mu\text{-}Cl)_4\{P(O)(OEt)_2\}_8]\cdot 6H_2O$, 1061–4
Tetrachlorobis(tetrahydrofuran)molybdate(III), a new intermediate for the synthesis of molybdenum(III) complexes: preparation and crystal structure, 1069–72
N,*N*′-Disubstituted dithiomalonamide complexes of antimony(III). Crystal and molecular structure of $[SbCl_3\{C_2H_5NHC(S)CH_2C(S)NHC_2H_5\}]$ with lone-pair occupation of an antimony co-ordination site, 1073–6
The crystal structure of diaquadichlorodimethyltin(IV)–purine (1/4), 1271–4
Homo- and hetero-dinuclear hydride-bridged complexes containing cyclo-octadiene: the crystal and molecular structure of $[(\eta^4\text{-}C_8H_{12})Ir(\mu\text{-}H)(\mu\text{-}Cl)IrH_2(PPh_3)_2]$, 1277–80
Chemistry of di- and tri-metal complexes with bridging carbene or carbyne ligands. Part 38. Ruthenium–tungsten compounds: crystal structures of $[RuW(\mu\text{-}Cl)(\mu\text{-}CMe)(Cl)(CO)_2(PPh_3)_2(\eta\text{-}C_5H_5)]$ and $[RuW_2(\mu_3\text{-}C_2Me_2)(CO)_7(\eta\text{-}C_5H_5)_2]$, 2017–24
A novel pentameric hydrolysis product of $SnMe_2Cl_2$: crystal and molecular structure of $[NHEt_3][(SnMe_2Cl)_5O_3]$, 2683–6
CHLORINE TRIOXIDE
Chlorine trioxide, Cl_2O_6, a most efficient perchlorating reagent in new syntheses of anhydrous metal perchlorates, chloryl and nitryl perchloratometalates of cobalt(II), nickel(II), and copper(II). Reactivity of chlorine trioxide with anhydrous or hydrated chlorides and nitrates, 297–306
CHROMIUM
Reaction of bis-μ-diethylphosphido-bis(tetracarbonylmetal) (*M*–*M*) (M = Cr or W) with tri-n-butylphosphine: kinetics and mechanism of a reaction involving seven-co-ordinate complexes, 91–8
Characterisation of chromium(VI) oxide tetrafluoride, $CrOF_4$, and caesium pentafluoro-oxochromate(VI) $Cs[CrOF_5]$, 529–34
Synthesis and characterisation of 1,2-bis(dimethylphosphino)ethane (dmpe) complexes of chromium-(0) and -(IV): *X*-ray crystal structures of *trans*-$Cr(N_2)_2(dmpe)_2$, *cis*-$Cr(CO)_2(dmpe)_2$, $Cr(C_2Ph_2)_2(dmpe)$, and $CrH_4(dmpe)_2$, 685–92
Difluorophosphate complexes of chromium, manganese, iron, cobalt, and nickel, 707–10
Electrochemistry of clusters. Part 4. Redox behaviour of tetrametallic clusters $[M^1{}_2M^2{}_2(\eta^5\text{-}C_5H_5)_2(\mu_3\text{-}CO)_2(\mu\text{-}CO)_4(PR_3)_2]$ (M^1 = Pt or Pd; M^2 = Cr, Mo, or W; R = Me, Et, Bu^n, or Ph), 711–6
Bimetallic systems. Part 9. The synthesis of and nuclear magnetic resonance studies on 10-membered ring complexes of type $[(OC)_4M^1(\mu\text{-}Ph_2PCH_2CH_2PPh_2)_2M^2(CO)_4]$ (M^1, M^2 = Cr, Mo, or W), 1009–14
Geometry-dependent complexation effects in carbon-13 nuclear magnetic resonance spectra of tricarbonyl(3—8-η-[2.2]paracyclophane)chromium and related complexes, 1065–8
Isolated CH stretching frequencies, methyl group geometry, and

CHROMIUM (contd)
methyl CH bond lengths and strengths in tricarbonyl(η^5-cyclopentadienyl)methylchromium(II), -molybdenum(II), and -tungsten(II), 1207–12
A mechanistic study on complexes of type *mer*-[Cr(CO)$_3$(η^2-L–L)(σ-L–L)] (where L–L = $Ph_2PCH_2PPh_2$, $Ph_2PNHPPh_2$, or $Ph_2PNMePPh_2$) using spectroscopic and convolutive electrochemical techniques, 1213–22
Spectroscopic studies on the higher binary fluorides of chromium: CrF_4, CrF_5, and CrF_6, both in the solid state and isolated in inert gas matrices, 1443–50
Pyramidal inversions and 1,2-metal shifts in pentacarbonyl-chromium, -molybdenum, and -tungsten derivatives of dialkyl disulphides and dialkyl diselenides. A nuclear magnetic resonance investigation, 1561–8
Dynamic nuclear magnetic resonance study of Group 6 metal pentacarbonyl complexes. Pyramidal inversion and 1,2-metal shifts in the complexes [M(CO)$_5$($Me_2\overbrace{CCH_2EECH_2}$)] (M = Cr, Mo, or W; E = S or Se), 1569–76
Sulphur-containing metal complexes. Part 14. Reactions of carbene anions with carbon disulphide or carbon diselenide, 1963–6
Chemistry of di- and tri-metal complexes with bridging carbene or carbyne ligands. Part 36. Reactions of the dimetal compounds [ReM(≡CC$_6$H$_4$Me-4)(CO)$_9$] (M = Cr, Mo, or W) with octacarbonyldicobalt; crystal structures of [$Co_2WRe(\mu_3$-CC$_6$H$_4$Me-4)(CO)$_{15}$] and [$Co_2Re(\mu_3$-CC$_6$H$_4$Me-4)(CO)$_{10}$], 2001–8
Synthesis and characterization of some chromium(III) complexes with glutathione, 2085–90
Pyramidal inversion in configurational isomers of tetracarbonyl[1,1,2,2-tetrakis(methylthio)ethane]chromium(0): a two-dimensional nuclear magnetic resonance exchange study, 2195–202
Selective conversion of CO into methanol at ordinary temperature. Part 4. Activation by iron(II), iron(III), and chromium(III) complexes, 2499–504
Magnetic and spectroscopic properties of some heterotrinuclear basic acetates of chromium(III), iron(III), and divalent metal ions, 2509–20

CIRCULAR DICHROISM
Optically active co-ordination compounds. Part 43. Polysulphide complexes of platinum(IV), 253–8

CLUSTER
New carbide clusters in the cobalt subgroup. Part 13. Synthesis and chemical characterization of the anions [$Co_6C(CO)_{14}$]$^-$, [$Co_6C(CO)_{15}$]$^{2-}$, and [$Co_8C(CO)_{18}$]$^{2-}$, and crystal structure of μ_6-carbido-ennea-μ-carbonyl-hexacarbonyl-*polyhedro*-hexacobaltate(2–) as its benzyltrimethylammonium salt; a comparison with isostructural species, 35–42
Metallation of 2-ethenylpyridine at triosmium clusters: *X*-ray crystal structures of the open trinuclear clusters [$Os_3H(CO)_9L(NC_5H_4CH{=}CH)$] (L = CO or PMe_2Ph), 85–90
Chemistry of the unsaturated cluster compound [$Os_3Pt(\mu$-H)$_2$(CO)$_{10}${P(cyclo-C$_6$H$_{11}$)$_3$}]; *X*-ray crystal structures of [$Os_3Pt(\mu$-H)$_2$(CO)$_{11}${P(cyclo-C$_6$H$_{11}$)$_3$}], [$Os_3Pt(\mu$-H)$_4$(CO)$_{10}${P(cyclo-C$_6$H$_{11}$)$_3$}], and [$Os_3Pt(\mu$-H)$_2(\mu$-CH$_2$)(CO)$_{10}$-{P(cyclo-C$_6$H$_{11}$)$_3$}] (two isomers), 177–90
The synthesis of $NR_4[(Ta_6Cl_{12})(H_2O)_6]X_4$ (R = Me or Et, X = Cl or Br) by the spontaneous reduction of $[Ta_6Cl_{12}]^{4+}$ to $[Ta_6Cl_{12}]^{3+}$ in acidic media: *X*-ray structure analysis of $NMe_4[(Ta_6Cl_{12})(H_2O)_6]Br_4$, 455–8
Systematic synthesis of tetranuclear osmium clusters by the reaction of trinuclear clusters with [$OsH_2(CO)_4$]; crystal structure of [$Os_4H_3Br(CO)_{13}$], 555–64
Synthesis and *X*-ray structural studies on the cluster compounds [$RuRh_2(\mu_3$-CO)$_2$(CO)$_3$(η-C$_5$Me$_5$)$_2$] and [$RuRh_2(\mu$-CO)$(\mu_3$-CO)(CO)$_2$(η^4-C$_8$H$_{10}$)(η-C$_5$Me$_5$)$_2$], 645–50
Electrochemistry of clusters. Part 4. Redox behaviour of tetrametallic clusters [$M^1{}_2M^2{}_2$(η^5-C$_5$H$_4$)$_2(\mu_3$-CO)$_2(\mu$-CO)$_4$(PR$_3$)$_2$] (M^1 = Pt or Pd; M^2 = Cr, Mo, or W; R = Me, Et, Bu^n, or Ph), 711–6
Spectroscopic and magnetic properties of cobalt(II) and nickel(II) clusters obtained from 1-(hydroxymethyl)-3,5-dimethylpyrazole. *X*-Ray structure of tetrakis[chloro(μ_3-3,5-dimethyl-*N*-oxymethylpyrazolato-N^2,μ_3-*O*)(ethanol)nickel(II)] 737–42
Lewis-base adducts of group 1B metal(I) compounds. Part 16. Synthesis, structure, and solid-state phosphorus-31 nuclear magnetic resonance spectra of some novel [$Cu_4X_4L_4$] (X = halogen, L = N,P base) 'cubane' clusters, 831–8
Interconversion of 42- and 44-electron platinum *triangulo*-clusters using chelating tertiary phosphine ligands and the structural characterisation of [1,3-bis(diphenylphosphino)propane]-tris(μ-sulphur dioxide)bis(tricyclohexylphosphine)triplatinum–benzene (1/2), [$Pt_3(\mu$-$SO_2)_3${P(C$_6$H$_{11}$)$_3$}$_2$(dppp)]·2C$_6$H$_6$, 845–50
New isomers of [$Os_3(CO)_{10}(PMe_2Ph)_2$] and [$Os_3(CO)_9(PMe_2Ph)_3$], 1037–42
New carbide clusters in the cobalt sub-group. Part 14. Synthesis and structural characterization of the anion [$Co_{13}C_2(CO)_{24}$]$^{3-}$ as its benzyltrimethylammonium salt, 1137–40
Cluster chemistry. Part 31. An [$Ru_3(CO)_{12}$] derivative with an [$Fe_3(CO)_{12}$]-like structure: preparation and *X*-ray structure of [$Ru_3(\mu$-CO)$_2$(CO)$_6${PPh(OMe)$_2$}$_4$], 1223–8
Cluster chemistry. Part 32. Synthesis and *X*-ray crystal structure of [$Ru_3(\mu_5$-η^2-C_2PPh_2-*P*)(μ-PPh_2)(CO)$_{13}$], a complex containing an alkynyl ligand in extended interaction with an open Ru_5 cluster, 1229–34
Synthesis, structures, and reactivities of some pentamethylcyclopentadienyl–sulphur compounds, 1303–8
New carbide clusters in the cobalt sub-group. Part 15. Synthesis and crystallographic characterization of di-μ_6-carbido-deca-μ-carbonyl-tridecacarbonyl-*polyhedro*-dodecarhodate(4−) as its tetrapropylammonium salt, [$N(C_3H_7)_4]_4$-[$Rh_{12}C_2(CO)_{23}$], 1309–14
Chemistry of di- and tri-metal complexes with bridging carbene or carbyne ligands. Part 33. Reactions of [W(≡CMe)(CO)$_2$(η-C$_5$H$_5$)] with the dimetal compounds [MRh(μ-CO)$_2$(η-C$_5$Me$_5$)$_2$] (M = Co or Rh); *X*-ray crystal structure of [$Rh_2W(\mu$-CO)$(\mu_3$-CMe)(CO)$_2$(η-C$_5$H$_5$)(η-C$_5$Me$_5$)$_2$], 1315–22
Chemistry of di- and tri-metal complexes with bridging carbene or carbyne ligands. Part 34. Reactions of sulphur and selenium with the di-irontungsten complexes [$Fe_2W(\mu_3$-CR)(μ-CO)(CO)$_8$(η-C$_5$H$_5$)] (R = C$_6$H$_4$Me-4 or Me); crystal structures of [$Fe_2W(\mu$-CC$_6$H$_4$Me-4)$(\mu_3$-S)(CO)$_7$(η-C$_5$H$_5$)] and [$Fe_2W(\mu_3$-SCMe)(CO)$_8$(η-C$_5$H$_5$)], 1323–30
The reaction of [$Os_3(\mu$-H)$_2$(CO)$_9$L] (L = CO or PEt_3) with dimethylcyanamide, Me_2NCN: *X*-Ray crystal structure of [$Os_3(\mu$-H)(μ-NCHNMe$_2$)(CO)$_{10}$] and the reactions of this complex with acids, 1355–60
Reactivity of the unsaturated anion decacarbonyltetra-μ-hydrido-trirhenate(1−) toward phenols. Crystal and molecular structures of the tetraethylammonium salts of the triangular cluster anion [$Re_3(\mu$-H)$_3(\mu$-OC$_6$F$_5$)(CO)$_{10}$]$^-$ and of the binuclear anion [$Re_2(\mu$-OC$_6$H$_3$)$_2$(CO)$_6$]$^-$, 1507–12
The synthesis of $NR_4[(Ta_6Cl_{12})(H_2O)_6]X_4$ (R = Me or Et, X = Cl or Br) by the spontaneous reduction of $[Ta_6Cl_{12}]^{4+}$ to $[Ta_6Cl_{12}]^{3+}$ in acidic media: *X*-ray structure analysis of $NMe_4[(Ta_6Cl_{12})(H_2O)_6]Br_4$ (1985, 455), 1531–2
Synthesis of bis[μ-bis(diphenylphosphino)methane]-tri-μ-carbonyl-tricarbonyl-*triangulo*-trirhodium(1+) perchlorate, [$Rh_3(CO)_3(\mu$-CO)$_3(\mu$-dppm)$_2$]ClO$_4$. An unusual 46-electron cluster possessing the A-frame structure with a bridging Rh(CO)$_3$ fragment, 1577–84
Synthesis and structural characterisation of some *triangulo*-platinum clusters containing isocyanide ligands, 1693–8
Reactions of some decaosmium clusters with electrophilic and nucleophilic reagents: *X*-ray structure analyses of [$N(PPh_3)_2$][$Os_{10}C(CO)_{24}(\mu$-I)], [$Os_{10}C(CO)_{24}(\mu$-I)$_2$], [$N(PPh_3)_2]_2$[$Os_{10}C(CO)_{22}$(NO)I], [$Os_{10}C(CO)_{23}${P(OMe)$_3$}$(\mu$-I)$_2$] and of two isomers of [$Os_{10}C(CO)_{21}${P(OMe)$_3$}$_4$], 1795–810
Studies of gold cluster compounds using high-resolution ^{31}P solid-state nuclear magnetic resonance spectroscopy, 1811–4
Carbon-13 nuclear magnetic resonance evidence of a relaxation process dominated by scalar coupling with a quadrupolar nucleus in [$Re_3(\mu$-H)$_4$(CO)$_{10}$]$^-$, 1899–902
Chemistry of di- and tri-metal complexes with bridging carbene or carbyne ligands. Part 36. Reactions of the dimetal compounds [ReM(≡CC$_6$H$_4$Me-4)(CO)$_9$] (M = Cr, Mo, or W) with octacarbonyldicobalt; crystal structures of [$Co_2WRe(\mu_3$-CC$_6$H$_4$Me-4)(CO)$_{15}$] and [$Co_2Re(\mu_3$-CC$_6$H$_4$Me-4)(CO)$_{10}$], 2001–8
Chemistry of di- and tri-metal complexes with bridging carbene or carbyne ligands. Part 38. Ruthenium–tungsten compounds: crystal structures of [RuW(μ-Cl)(μ-CMe)(Cl)(CO)$_2$(PPh$_3$)$_2$(η-C$_5$H$_5$)] and [$RuW_2(\mu_3$-C_2Me_2)(CO)$_7$(η-C$_5$H$_5$)$_2$], 2017–24
Crystal structure and spectroscopic and redox properties of the iron–sulphur cluster compound [$NEt_4]_2$[Fe_4S_4(SC$_6$H$_4$NH$_2$-4)$_4$],

CLUSTER (contd)
2161–6
Synthesis of tetra- and penta-nuclear platinum–osmium carbido-cluster complexes from non-carbido-precursors: *X*-ray structural studies on $[Os_3Pt(\mu\text{-}H)_2(\mu_4\text{-}C)(CO)_{10}\{P(cyclo\text{-}C_6H_{11})_3\}]$, $[Os_3Pt_2(\mu\text{-}H)_2(\mu_5\text{-}C)(\mu\text{-}CO)(CO)_9\{P(cyclo\text{-}C_6H_{11})_3\}_2]$, and $[Os_3Pt_2(\mu\text{-}H)(\mu_5\text{-}C)(\mu\text{-}OMe)(\mu\text{-}CO)(CO)_9\{P(cyclo\text{-}C_6H_{11})_3\}_2]$, 2437–48
Acid-induced displacement of acetaldehyde from a μ-vinyloxy-triosmium cluster, 2479–82
The $[Fe_3(\mu_3\text{-}CR)(CO)_{10}]^-$ cluster anions as building blocks for the synthesis of mixed-metal clusters. Part 1. Synthesis of mixed clusters $[MFe_3(\mu_3\text{-}CMe)(CO)_{10}(PPh_3)]$ (M = Cu or Au) and crystal structure of $[CuFe_3(\mu_3\text{-}CMe)(CO)_{10}(PPh_3)]$, 2521–4

COBALAMIN
A new electron spin resonance spectrum for an exchange- and dipole–dipole coupled superoxocobalamin · · · free-radical pair occurring in adenosylcobalamin-containing systems, 891–4
Self-association of organocobalamins in aqueous solution, 1381–6

COBALOXIME
Routes of formation and crystal structure of an alkylperoxycobaloxime: bis[dimethylglyoximato(1−)](4-ethoxycarbonylbut-3-en-2-ylperoxo)(pyridine)cobalt(III), 1997–2000

COBALT
New carbide clusters in the cobalt subgroup. Part 13. Synthesis and chemical characterization of the anions $[Co_6C(CO)_{14}]^-$, $[Co_6C(CO)_{15}]^{2-}$, and $[Co_8C(CO)_{18}]^{2-}$, and crystal structure of μ_6-carbido-ennea-μ-carbonyl-hexacarbonyl-*polyhedro*-hexacobaltate(2−) as its benzyltrimethylammonium salt; a comparison with isostructural species, 35–42
Chlorine trioxide, Cl_2O_6, a most efficient perchlorating reagent in new syntheses of anhydrous metal perchlorates, chloryl and nitryl perchlorato-metalates of cobalt(II), nickel(II), and copper(II). Reactivity of chlorine trioxide with anhydrous or hydrated chlorides and nitrates, 297–306
Ligand substitution on (*N*,*N*-dimethylformamide)[2,2′,2″-tri(*N*,*N*-dimethylamino)triethylamine]cobalt(II), 413–6
Co-ordination of alkali metals by open-chain polyethers in transition metal complexes. Part 4. Variation in alkali-metal ion selectivity in cobalt and zinc complexes of 1-(*o*-carboxymethoxyphenoxy)-2-(*o*-hydroxyphenoxy)ethane (HL) and the *X*-ray and molecular structure of $[Zn(NH_4L)_2]$, 459–64
Proton exchange and base hydrolysis of *syn,anti-cis*-dichloro(1,4,7,10-tetra-azacyclododecane)cobalt(III) cations, 475–8
Macrocyclic polyphosphane ligands. Iron(II), cobalt(II), and nickel(II) complexes of (4*RS*,7*RS*,13*SR*,16*SR*)-tetraphenyl-1,10-dipropyl-1,10-diaza-4,7,13,16-tetraphosphacyclo-octadecane: crystal structures of their tetraphenylborate derivatives, 479–86
Synthesis and characterization of several mononuclear cobalt(IV) dithiocarbamates, $[Co(S_2CNRR')_3]ClO_4$ [R = R′ = Me, Et, or CH_2Ph; R = Me, R′ = Ph; R,R′ = $-(CH_2)_{4-}$], 597–600
Solvent dependence of the stereochemistry of base-catalysed solvolysis of *trans*-$[Co(NH_3)_4(^{15}NH_3)X]^{3+/2+}$ ions in dipolar aprotic solvents, 659–62
Difluorophosphate complexes of chromium, manganese, iron, cobalt, and nickel, 707–10
Spectroscopic and magnetic properties of cobalt(II) and nickel(II) clusters obtained from 1-(hydroxymethyl)-3,5-dimethylpyrazole. *X*-Ray structure of tetrakis[chloro(μ$_3$-3,5-dimethyl-*N*-oxymethylpyrazolato-N^2,μ$_3$-*O*)(ethanol)nickel(II)] 737–42
Optical resolution of DL-2,2′-bipiperidine through its cobalt(III) complex, 895–8
Electron-transfer reactions from *cis*-dialkylbis(2,2′-bipyridyl)cobalt(III) complexes to organic oxidants, 899–904
Reactions of cobalt(II) protoporphyrin IX dimethyl ester, $[Co^{II}P]$, and $[Co^{III}P(Cl)]$ in co-ordinating aliphatic alcohols, 1095–102
New carbide clusters in the cobalt sub-group. Part 14. Synthesis and structural characterization of the anion $[Co_{13}C_2(CO)_{24}]^{3-}$ as its benzyltrimethylammonium salt, 1137–40
Macrocyclic polyphosphane ligands. Cobalt(II) and nickel(II) complexes of 5*RS*,8*RS*,16*RS*,19*RS*-tetraphenyl-5,8,16,19-tetraphospha-1,12-dithiacyclodocosane (δ-L^1) and the crystal structure of $[Co(\delta\text{-}L^1)][BF_4]_2{\cdot}0.5H_2O$, 1179–82
Crystal structures and interrelationships of the blue and green conformational isomers of tetrakis(trimethylarsine sulphide)cobalt(II) perchlorate, 1289–94
Chemistry of di- and tri-metal complexes with bridging carbene or carbyne ligands. Part 33. Reactions of $[W(\equiv CMe)(CO)_2(\eta\text{-}C_5H_5)]$ with the dimetal compounds $[MRh(\mu\text{-}CO)_2(\eta\text{-}C_5Me_5)_2]$ (M = Co or Rh); *X*-ray crystal structure of $[Rh_2W(\mu\text{-}CO)(\mu_3\text{-}CMe)(CO)_2(\eta\text{-}C_5H_5)(\eta\text{-}C_5Me_5)_2]$, 1315–22
Kinetic studies of oxidative dealkylation of alkylcobalamins by hexachloroplatinate(IV), 1375–80
Self-association of organocobalamins in aqueous solution, 1381–6
Macrocyclic polyphosphane ligands. Cobalt(II) and nickel(II) complexes of the γ and ε diastereoisomers of 4,7,13,16-tetraphenyl-4,7,13,16-tetraphospha-1,10-dithiacyclo-octadecane (L^1) and the crystal structure of $[Co(\varepsilon\text{-}L^1)][BPh_4]_2{\cdot}$ EtOH, 1425–30
Steric changes at labelled NH_2 sites during the base-catalysed ammoniation of the *trans*-azido(dimethyl sulphoxide)bis(ethylenediamine)cobalt(III) ion, 1495–500
Dismutation of superoxide ion in an aprotic solvent by 5,10,15,20-tetra-*p*-tolylporphyrinatocobalt(II), 1513–6
Reactions of metal ions with triketones in solution. Part 1. Formation constants for the systems of heptane-2,4,6-trione, 1-phenylhexane-1,3,5-trione, 1,5-diphenylpentane-1,3,5-trione, and 2,2′-dihydroxybenzophenone with proton, nickel(II), and cobalt(II), 1543–6
Potentiometric study of the complex-formation equilibria of manganese(II), cobalt(II), nickel(II), copper(II), and zinc(II) with ethylenediamine-*N*-acetic acid, 1605–8
The chemistry of vitamin B_{12}. Part 25. Mechanism of the β-elimination of olefins from alkylcorrinoids; evidence for an initial homolytic fission of the Co–C bond, 1613–8
The chemistry of vitamin B_{12}. Part 26. Co-ordination of the malonitrile anion by Co^{III} corrinoids: first experimental determination of equilibrium constants for the co-ordination of a tetrahedral carbanion by a transition–metal ion, 1619–22
Kinetics and mechanism of the reductions of tris(oxalato)cobaltate(III) ion by ruthenium(II) species in aqueous solution, 1665–8
Transition-metal co-ordination compounds of a novel aniline-based pyrazole derivative. *X*-Ray crystal structures of [*NN*-bis(3,5-dimethylpyrazol-1-ylmethyl)aminobenzene]-dichlorocobalt(II) and -dibromocopper(II), 1699–706
Studies in the flexibility of macrocycle ligands. Calculation of macrocycle cavity size by force-field methods. Crystal and molecular structures of $[CoLCl][ClO_4]_2$ and $[CuL][PF_6]_2$ {L = 2,13-dimethyl-3,6,9,12,18-penta-azabicyclo[12.3.1]octadeca-1(18),14,16-triene}, 1829–34
Synthesis, characterization and acid hydrolysis of [*N*,*N*-bis(2-aminoethyl)-1,2-ethanediamine]bis(solvent)cobalt(III) complexes, 1923–8
New photoreduction catalysis by $[Cu(N\text{–}N)(PPh_3)_2]^+$ (N–N = 2,9-dimethyl-1,10-phenanthroline or 4,4′,6,6′-tetramethyl-2,2′-bipyridine) and its application to cobalt(III) complexes, 1959–62
Routes of formation and crystal structure of an alkylperoxycobaloxime: bis[dimethylglyoximato(1−)](4-ethoxy carbonylbut-3-en-2-ylperoxo)(pyridine)cobalt(III), 1997–2000
Chemistry of di- and tri-metal complexes with bridging carbene or carbyne ligands. Part 36. Reactions of the dimetal compounds $[ReM(\equiv CC_6H_4Me\text{-}4)(CO)_9]$ (M = Cr, Mo, or W) with octacarbonyldicobalt; crystal structures of $[Co_2WRe(\mu_3\text{-}CC_6H_4Me\text{-}4)(CO)_{15}]$ and $[Co_2Re(\mu_3\text{-}CC_6H_4Me\text{-}4)(CO)_{10}]$, 2001–8
Steric changes at labelled NH_2 sites during the base-catalysed ammoniation of the *trans*-azido(dimethyl sulphoxide)bis(ethylenediamine)cobalt(III) ion, 2223–4
Investigation into aroylhydrazones as chelating agents. Part 7. Synthesis and spectroscopic characterization of complexes of Mn^{II}, Co^{II}, Ni^{II}, Cu^{II}, and Zn^{II} with 2,6-diacetylpyridine bis(2-aminobenzoylhydrazone) and *X*-ray structure of chloro[2,6-diacetylpyridine bis(2-aminobenzoylhydrazone)](methanol)manganese(II) chloride monohydrate, 2387–92
Crystal structures of the difluorophosphate complexes, $Co(O_2PF_2)_2{\cdot}2MeCN$ and $Cu(O_2PF_2)_2$, 2433–6
Reactions of co-ordinated ligands. Part 35. Evidence for carbon–carbon double-bond cleavage of cyclopropenes in their reaction with dinuclear cobalt, rhodium, and iridium complexes; crystal structure and protolysis of $[Rh_2(\mu\text{-}CO)(\mu\text{-}COCHCMe_2CH)(\mu\text{-}C_5Me_5)_2]$, 2483–92
Reactions of metal ions with triketones. Part 2. Kinetics and mechanisms of the reactions of nickel(II) and cobalt(II) with heptane-2,4,6-trione in methanol–water (70:30 v/v), 2565–70

CONDUCTIVITY
X-Ray crystal structures and properties of tris(tetrathiafulvalenium) tetrachlorodimethylstannate(IV) and tetrathiafulvalenium trichlorodimethylstannate (IV), 465–70
Molecular structure and solid-state properties of the two-dimensional conducting mixed-valence complex $[NBu_4]_{0.29}[Ni(dmit)_2]$ and the neutral $[Ni(dmit)_2]$ (H_2dmit = 4,5-dimercapto-1,3-dithiole-2-thione); members of an electron-transfer series, 783–94
Copper(II) salts of metal dithiolates, 1731–2
Electrical and magnetic properties of $K_3Cu_8S_6$, 2319–20
CONFIGURATION
Sila-pharmaca. Part 32. Crystal and molecular structures of the (*R*)-enantiomer and the racemate of the antimuscarinic agent (cyclohexyl)phenyl[2-(pyrrolidin-1-yl)ethyl]silanol (sila-procyclidine), 1743–6
CONFORMATION
Conformation analysis of compounds of the type $[Fe(\eta^5\text{-}C_5H_5)(CO)(PPh_3)(CH_2R)]$ (R = alkyl or aryl): a comment, 2691
Conformation analysis of compounds of the type $[Fe(\eta^5\text{-}C_5H_5)(CO)(PPh_3)(CH_2R)]$ (R = alkyl or aryl): a reply, 2691
COPPER
Enantioselectivity of nickel(II) and copper(II) complexes of Schiff bases derived from amino acids and (*S*)-*o*-[(*N*-benzylprolyl)amino]acetophenone or (*S*)-*o*-[(*N*-benzylprolyl)amino]benzaldehyde. Crystal and molecular structures of [Ni{(*S*)-bap-(*S*)-Val}] and [Cu{(*S*)-bap-(*S*)-Val}], 17–26
Synthesis and characterization of copper(II) complexes with unsymmetrical quadridentate Schiff bases derived from β-diketone, diamine, and *o*-hydroxybenzaldehyde ligands, 101–6
Synthesis and characterization of a new family of binuclear copper(II) complexes with a flexible bridge and the π–π type charge-transfer interaction with 1,3,5-trinitrobenzene, 107–10
Lewis-base adducts of Group 1B metal(I) compounds. Part 12. Structural studies of some bis(methyl-substituted pyridine)-copper(I) and -silver(I) nitrates and perchlorates, 117–24
Lewis-base adducts of Group 1B metal(I) compounds. Part 13. Crystal structure determinations of tetrakis(triphenyl-phosphine)-copper(I) and -silver(I) perchlorates, bis(pyridine)bis(triphenylphosphine)copper(I) perchlorate, (2,2′-bipyridyl)bis(triphenylphosphine)copper(I) perchlorate, and tetrahydroboratobis(triphenylphosphine)copper(I)–pyridine (1/0.5), 125–34
The crystal structure and electronic properties of the complex acetatobis(1,10-phenanthroline)copper(II) perchlorate dihydrate, acetatobis(1,10-phenanthroline)copper(II) nitrate dihydrate, and acetatobis(1,10-phenanthroline)zinc(II) tetrafluoroborate dihydrate, 141–50
Sulphur ligand–metal complexes. Part 16. Copper complexes of thioethers and the single-crystal *X*-ray structure of the polymeric mixed-valence complex, penta-μ-chloro-tris-μ-tetrahydrothiophene-tetracopper(I,II), 151–8
Stabilisation of copper(I) by an azoimine ligand: redox properties and reactions of bis(phenylazoacetaldoximato)bis(phenylacetaldoxime)dicopper(I), 159–62
Preparation of 11-(2′-dimethylaminoethyl)-1,4,7-trimethyl-1,4,7,11-tetra-azacyclotetradecane, and characterisation of its nickel(II), copper(II), and zinc(II) complexes, 219–22
Metal–phenoxyalkanoic acid interactions. Part 13. Copper(II)–(2-chlorophenoxy)ethanoic acid complexes. Crystal and molecular structures of *catena*-tetra-μ-[(2-chlorophenoxy)ethanoato-*O*,*O*′]-dicopper(II), *catena*-(2-aminopyrimidine-*N*,*N*′)-tetra-μ-[(2-chlorophenoxy)ethanoato-*O*,*O*′]-dicopper(II), and 1,2,2,2,2-penta-aqua-tetra-μ-[(2-chlorophenoxy)ethanoato-*O*,*O*′]-copper(II)calcium(II), 243–52
Chlorine trioxide, Cl_2O_6, a most efficient perchlorating reagent in new syntheses of anhydrous metal perchlorates, chloryl and nitryl perchloratometalates of cobalt(II), nickel(II), and copper(II). Reactivity of chlorine trioxide with anhydrous or hydrated chlorides and nitrates, 297–306
Crystal and molecular structures and magnetic properties of tetrameric copper(II) complexes with 3-hydroxy-5-hydroxymethyl-4-(4′-hydroxy-4′-phenyl-2′-azabut-1′-en-1′-yl)-2-methylpyridine (H_2L^3), $[Cu_4L^3{}_4]$·9CH_3OH and 3-hydroxy-5-hydroxymethyl-4-(4′-hydroxy-3′-methyl-4′-phenyl-2′-azabut-1′-en-1′-yl)-2-methylpyridine (H_2L^1), $[Cu_4L^1{}_4]$·8CH_3CH_2OH: two complexes with ferromagnetic ground states, 315–20
Comparison between the bis(2-thiopyridine *N*-oxide) derivatives of Cu^{II} and Ni^{II}: an electron spin resonance study, 379–82
The L-proline residue as a 'break-point' in metal–peptide systems, 535–40
Synthesis and physico-chemical properties of cationic derivatives of phthalocyaninatocopper(II), 651–8
Reaction of some aliphatic diamines with four-co-ordinated unsymmetrical ketoenamine copper(II)and nickel(II) complexes, 803–6
Lewis-base adducts of group 1B metal(I) compounds. Part 16. Synthesis, structure, and solid-state phosphorus-31 nuclear magnetic resonance spectra of some novel $[Cu_4X_4L_4]$ (X = halogen, L = N,P base) 'cubane' clusters, 831–8
Lewis-base adducts of group 1B metal(I) compounds. Part 17. Synthesis and crystal structures of adducts of copper(I) cyanide with nitrogen bases, 839–44
Crystal and molecular structures and magnetic properties of bromo(3-dimethylaminopropan-1-olato)copper(II), chloro(3-dimethylaminopropan-1-olato)copper(II), and (3-diethylaminopropan-1-olato)isocyanatocopper(II), 913–20
Co-ordination chemistry of pyridyl and *N*-methylimidazolyl ketones. Synthetic and *X*-ray structural studies of copper(II), nickel(II), and dimethylgold(III) complexes, 981–6
Electrostatic solvent effect on the formation of the mixed-chelate complex (acetylacetonato)(diethyldithocarbamato)-copper(II), 987–90
Solution chemistry of macrocycles. Part 4. Thermodynamics of protonation and complexation of several N_2S_2 macrocycles, 1169–72
Specific binding of the tyrosine residue in copper(II) complexes of Tyr-Pro-Gly-Tyr and Tyr-Gly-Pro-Tyr, 1201–6
Crystal and molecular structures and magnetic properties of four new exchange-coupled copper(II) complexes derived from different 3-*N*,*N*-dialkylamino-1-propanols and pseudohalogens, 1243–8
Characterization of the adducts formed by Cu(CN) and Cu(NCS) with biquinoline. The crystal structure of the polymeric cyano-compound containing both linear and tetrahedrally co-ordinated copper(I), $[\{Cu_3(bq)_2(CN)_3\}_n]$, 1285–8
Compartmental ligands. Part 11. Copper(II) complexes of 'half-unit' Schiff bases: crystal and molecular structures of one monomeric and of one dimeric complex, 1471–8
Potentiometric study of the complex-formation equilibria of manganese(II), cobalt(II), nickel(II), copper(II), and zinc(II) with ethylenediamine-*N*-acetic acid, 1605–8
The concentration and selective extraction of copper(II), rhodium(III), and iridium(III) using a copolymer functionalised with dithiocarbamate groups. Spectroscopic evidence for the nature of the binding sites, 1655–60
Transition-metal co-ordination compounds of a novel aniline-based pyrazole derivative. *X*-Ray crystal structures of [*NN*-bis(3,5-dimethylpyrazol-1-ylmethyl)aminobenzene]-dichlorocobalt(II) and -dibromocopper(II), 1699–706
Electron spin resonance parameters for some copper(II)–bis(amino-acid) complexes, 1717–8
Iodine-127 Mössbauer spectroscopy of copper(I) iodide–phosphine adducts, 1727–30
Copper(II) salts of metal dithiolates, 1731–2
The structure of a dinuclear copper(I) complex of a Schiff-base ligand containing a copper–copper bond, 1771–4
Studies in the flexibility of macrocycle ligands. Calculation of macrocycle cavity size by force-field methods. Crystal and molecular structures of $[CoLCl][ClO_4]_2$ and $[CuL][PF_6]_2$ {L = 2,13-dimethyl-3,6,9,12,18-penta-azabicyclo[12.3.1]octadeca-1(18),14,16-triene}, 1829–34
Equilibria between mono- and bi-nuclear complexes in $Cu(O_2CMe)_2$–pyridine derivative–diluent systems. The influence of the amine ligand basicity, 1849–52
Bis(triphenylphosphine)copper(I) derivatives of substituted *arachno* nine-vertex borane anions, $Cu(PPh_3)_2(B_9H_{13}X)$ (X = H, NCS, NCSe, $NCBPh_3$, $NCBH_3$, or $NCBH_2NCBH_3$), 1903–8
Magnetic behaviour of tetrakis[(2-diethylaminoethanolato)isocyanatocopper(II)], a complex with an antiferromagnetic ground state; the crystal and molecular structure of the triclinic modification, 1909–14
Crystal structures and magnetic properties of binuclear five-co-ordinate copper(II) complexes with a phenolate bridge and their catalytic functions in multielectron redox reactions, 1945–52
New photoreduction catalysis by $[Cu(N\text{–}N)(PPh_3)_2]^+$ (N–N = 2,9-

COPPER (contd)
dimethyl-1,10-phenanthroline or 4,4′,6,6′-tetramethyl-2,2′-bipyridine) and its application to cobalt(III) complexes, 1959–62
Metal complexes of vitamin B_6 related compounds. Crystal and molecular structures of aqua(5′-phosphopyridoxylideneglycinato)copper(II) trihydrate and bis(pyridoxylideneglycinato)nickel(II) hexahydrate, 2051–8
X-Ray crystal structures and magnetic properties of azide-bridged binuclear copper(II) complexes containing the Schiff-base ligand derived from 2-pyridinecarbaldehyde and histamine. Structure–magnetism relationship, 2095–100
Unidentate *versus* symmetrically and unsymmetrically bidentate nitrate co-ordination in pyrazole-containing chelates. The crystal and molecular structures of (nitrato-*O*)[tris(3,5-dimethylpyrazol-1-ylmethyl)amine]copper(II) nitrate, (nitrato-*O*,*O*′)[tris(3,5-dimethylpyrazol-1-ylmethyl)amine]nickel(II)nitrate, and (nitrato-*O*)(nitrato-*O*,*O*′)[tris(3,5-dimethylpyrazol-1-ylmethyl)amine]cadmium(II), 2177–84
Copper(II) complexes of *N*,*N*′-bis(2-carbamoylethyl)ethylenediamine in methanol–water, 2217–20
Influence of decreasing solvent polarity (dioxane–water mixtures) on the stability and structure of binary and ternary complexes of adenosine 5′-triphosphate and uridine 5′-triphosphate, 2291–304
Synthesis and cryptate complexes of azathia macropolycyclic ligands based on 12-membered N_2S_2 and 15-membered N_2S_3 macrocyclic subunits, 2311–8
Electrical and magnetic properties of $K_3Cu_8S_6$, 2319–20
Copper complexes with quadridentate bis(pyrazolyl)thioether amine and tris(pyrazolyl)amine ligands. Structural characterization of the complexes $[Cu(NCS)(tpea)][Cu(NCS)_2]$ and $[CuCl(bdma)]Cl{\cdot}2H_2O$, 2327–32
The effect of a dansyl group on the co-ordinative ability of *N*-protected amino acids. Part 1. Behaviour of the copper(II) ion–*N*-dansylglycinate system in aqueous and methanolic solution, 2363–8
Investigation into aroylhydrazones as chelating agents. Part 7. Synthesis and spectroscopic characterization of complexes of Mn^{II}, Co^{II}, Ni^{II}, Cu^{II}, and Zn^{II} with 2,6-diacetylpyridine bis(2-aminobenzoylhydrazone) and *X*-ray structure of chloro[2,6-diacetylpyridine bis(2-aminobenzoylhydrazone)](methanol)manganese(II) chloride monohydrate, 2387–92
Crystal structures of the difluorophosphate complexes, $Co(O_2PF_2)_2{\cdot}2MeCN$ and $Cu(O_2PF_2)_2$, 2433–6
The $[Fe_3(\mu_3\text{-}CR)(CO)_{10}]^-$ cluster anions as building blocks for the synthesis of mixed-metal clusters. Part 1. Synthesis of mixed clusters $[MFe_3(\mu_3\text{-}CMe)(CO)_{10}(PPh_3)]$ (M = Cu or Au) and crystal structure of $[CuFe_3(\mu_3\text{-}CMe)(CO)_{10}(PPh_3)]$, 2521–4
Lewis-base adducts of Group 1B metal(I) compounds. Part 18. Stereochemistries and structures of the 1:1 neutral complexes of Cu^IX with 1,10-phenanthroline (X = I) or 2,9-dimethyl-1,10-phenanthroline (X = I, Br, or Cl), 2531–40
Lewis-base adducts of Group 1B metal(I) compounds. Part 19. Crystal structures of bis(1,10-phenanthroline)copper(I) perchlorate and dibromocuprate(I), 2541–6
Molecular structure of gaseous copper(I) acetate as determined by electron diffraction, 2555–60
Crystal and molecular structure and magnetic properties of linear trimeric copper(II) complexes with predominant ferromagnetic exchange interaction, 2609–14

CORRINOID
The chemistry of vitamin B_{12}. Part 26. Co-ordination of the malonitrile anion by Co^{III} corrinoids: first experimental determination of equilibrium constants for the co-ordination of a tetrahedral carbanion by a transition–metal ion, 1619–22

CORROSION
Electrochemical studies in HNO_3–N_2O_4 mixtures: corrosion of stainless steel in HNO_3–N_2O_4 mixtures and the effect of inhibitors, 2551–4

CROWN ETHER
Complexes of lanthanoid salts with macrocyclic ligands. Part 17. Synthesis and crystal and molecular structure of a hydroxide-bridged praseodymium trifluoroacetate complex with 15-crown-5-ether, $[Pr_2(CF_3CO_2)_3(OH)(C_{10}H_{20}O_5)_2]{\cdot}[Pr_2(CF_3CO_2)_8]$, 885–90

CRYPTATE
Synthesis and cryptate complexes of azathia macropolycyclic ligands based on 12-membered N_2S_2 and 15-membered N_2S_3 macrocyclic subunits, 2311–8

CRYSTAL STRUCTURE
Co-ordinative interactions in chelated complexes of silicon. Part 5. Chiralty of five-co-ordinate silicon compounds: crystal and molecular structures of 1-(chlorodimethylsilyl)-and 1-(dichloromethylsilyl)-1,2,3,4-tetrahydro-1,10-phenanthroline, 1–4
Preparation and characterisation of adducts of bismuth pentafluoride and antimony pentafluoride by vibrational spectroscopy, *X*-ray powder diffraction, and single-crystal *X*-ray crystallography, 9–16
Enantioselectivity of nickel(II) and copper(II) complexes of Schiff bases derived from amino acids and (*S*)-*o*-[(*N*-benzylprolyl)amino]acetophenone or (*S*)-*o*-[(*N*-benzylprolyl)amino]benzaldehyde. Crystal and molecular structures of [Ni{(*S*)-bap-(*S*)-Val}] and [Cu{(*S*)-bap-(*S*)-Val}], 17–26
New carbide clusters in the cobalt subgroup. Part 13. Synthesis and chemical characterization of the anions $[Co_6C(CO)_{14}]^-$, $[Co_6C(CO)_{15}]^{2-}$, and $[Co_8C(CO)_{18}]^{2-}$, and crystal structure of μ_6-carbido-ennea-μ-carbonyl-hexacarbonyl-*polyhedro*-hexacobaltate(2–) as its benzyltrimethylammonium salt; a comparison with isostructural species, 35–42
Toluene-3,4-dithiol (H_2tdt) complexes of Group 5B halides. Observations of lone-pair stereochemical activity and redox behaviour. Crystal and molecular structures of [AsCl(tdt)] and $[PPh_4][Sb(tdt)_3]$, 69–74
Metallation of 2-ethenylpyridine at triosmium clusters: *X*-ray crystal structures of the open trinuclear clusters $[Os_3H(CO)_9L(NC_5H_4CH{=}CH)]$ (L = CO or PMe_2Ph), 85–90
Synthesis and characterization of copper(II) complexes with unsymmetrical quadridentate Schiff bases derived from β-diketone, diamine, and *o*-hydroxybenzaldehyde ligands, 101–6
Lewis-base adducts of Group 1B metal(I) compounds. Part 11. Synthesis and crystal structure of adducts of silver(I) bromide with monomethyl-substituted pyridine bases, 111–6
Lewis-base adducts of Group 1B metal(I) compounds. Part 12. Structural studies of some bis(methyl-substituted pyridine)-copper(I) and -silver(I) nitrates and perchlorates, 117–24
Lewis-base adducts of Group 1B metal(I) compounds. Part 13. Crystal structure determinations of tetrakis(triphenyl-phosphine)-copper(I) and -silver(I) perchlorates, bis(pyridine)bis(triphenylphosphine)copper(I) perchlorate, (2,2′-bipyridyl)bis(triphenylphosphine)copper(I) perchlorate, and tetrahydroboratobis(triphenylphosphine)copper(I)–pyridine (1/0.5), 125–34
The crystal structure and electronic properties of the complex acetatobis(1,10-phenanthroline)copper(II) perchlorate dihydrate, acetatobis(1,10-phenanthroline)copper(II) nitrate dihydrate, and acetatobis(1,10-phenanthroline)zinc(II) tetrafluoroborate dihydrate, 141–50
Sulphur ligand–metal complexes. Part 16. Copper complexes of thioethers and the single-crystal *X*-ray structure of the polymeric mixed-valence complex, penta-μ-chloro-tris-μ-tetrahydrothiophene-tetracopper(I,II), 151–8
Chemistry of the unsaturated cluster compound $[Os_3Pt(\mu\text{-}H)_2(CO)_{10}\{P(cyclo\text{-}C_6H_{11})_3\}]$; *X*-ray crystal structures of $[Os_3Pt(\mu\text{-}H)_2(CO)_{11}\{P(cyclo\text{-}C_6H_{11})_3\}]$, $[Os_3Pt(\mu\text{-}H)_4(CO)_{10}\{P(cyclo\text{-}C_6H_{11})_3\}]$, and $[Os_3Pt(\mu\text{-}H)_3(\mu\text{-}CH_2)(CO)_{10}\{P(cyclo\text{-}C_6H_{11})_3\}]$ (two isomers), 177–90
Co-ordination chemistry of higher oxidation states. Part 13. Synthesis and properties of alkali-metal hydroxo-oxo-osmate(VIII) compounds and the molecular structure of $Cs[O_4Os(\mu\text{-}OH)OsO_4]$, 199–204
The template synthesis and crystal and molecular structure of a seven-co-ordinate manganese(II) complex with 2,6-diacetylpyridine mono(2-aminobenzoylhydrazone), 215–8
Crystal structures and rotameric forms of some diarylsulphonyl-mono-, -di-, and -tri-selanes and their sulphur analogues, 231–8
Metal–phenoxyalkanoic acid interactions. Part 13. Copper(II)–(2-chlorophenoxy)ethanoic acid complexes. Crystal and molecular structures of *catena*-tetra-μ-[(2-chlorophenoxy)ethanoato-*O*,*O*′]-dicopper(II), *catena*-(2-aminopyrimidine-*N*,*N*′)-tetra-μ-[(2-chlorophenoxy)ethanoato-*O*,*O*′]-dicopper(II), and 1,2,2,2,2-penta-aqua-tetra-μ-[(2-chlorophenoxy)ethanoato-*O*,*O*′]-copper(II)calcium(II), 243–52
Chemistry of metallacyclobutanones. Part 2. Synthesis and ring inversion of some highly puckered metallacyclobutan-3-one (slipped oxodimethylenemethane) complexes of palladium; crystal structures of 2,4-bis(methoxycarbonyl)-1,1-bis(triphenylphosphine)palladacyclobutan-3-one, 2,4-

CRYSTAL STRUCTURE (contd)
bis(methoxycarbonyl)-1,1-bis(triphenylarsine)palladacyclobutan-3-one, and 1,1-(2′,2″-bipyridyl)-2,4-bis(methoxycarbonyl)palladacyclobutan-3-one, 259–68
Phosphorus–phosphorus bond cleavage in the cage molecule P_4S_3: synthesis and crystal structure of the trinuclear platinum complex $[\{Pt(\mu\text{-}P_4S_3)(PPh_3)\}_3]\cdot C_6H_6$, 291–6
Crystal and molecular structures and magnetic properties of tetrameric copper(II) complexes with 3-hydroxy-5-hydroxymethyl-4-(4′-hydroxy-4′-phenyl-2′-azabut-1′-en-1′-yl)-2-methylpyridine (H_2L^3), $[Cu_4L^3{}_4]\cdot 9CH_3OH$ and 3-hydroxy-5-hydroxymethyl-4-(4′-hydroxy-3′-methyl-4′-phenyl-2′-azabut-1′-en-1′-yl)-2-methylpyridine (H_2L^1), $[Cu_4L^1{}_4]\cdot 8CH_3CH_2OH$: two complexes with ferromagnetic ground states, 315–20
Chemical and structural aspects of silver–triphenylarsine complexes and silver–tin complex salts, 321–32
The preparation and co-ordination chemistry of 2,2′:6′,2″-terpyridine macrocycles. Part 4. Structural characterisation of an intermediate in a transient template reaction, 333–6
Synthesis of α-lithioarylmethanes of *m*-xylene and its α-trimethylsilyl derivatives; crystal structure of $[\{Li(Me_2NCH_2CH_2NMe_2)\}_2\{C_6H_4(CHSiMe_3)_2\text{-}m\}]$, 337–44
Static and dynamic nuclear magnetic resonance studies of complexes of trimethylplatinum(IV) halides with olefinic thio-and seleno-ethers. *X*-Ray crystal structures of $[PtXMe_3(MeSeCH{=}CHSeMe)]$ (X = Cl or I), 345–54
Purification and *X*-ray crystal structure of bis[tris(trimethylsilyl)methyl]diarsene, 383–6
The preparation and hydridic reduction of dicationic dicarbonyl complexes of osmium(II); the crystal and molecular structure of *trans*-bis[1,2-bis(diphenylphosphino)ethane-*PP′*]carbonylformylosmium(II) hexafluoroantimonate–dichloromethane (1/1), 387–94
Niobium(IV) sulphidohalides: preparation of $Nb_2X_4S_3$ and $Nb_2X_4S\text{-}nL$[X = Br or Cl; *n* = 4, L = NCMe, SMe_2, or tetrahydrothiophene (tht); *n* = 2, L = $PhSCH_2CH_2SPh$]. Crystal and molecular structure of $Nb_2Cl_4S_2\text{-}4tht$, 417–22
Hexakis(trimethylphosphine)molybdenum chemistry: dinitrogen, ethylene, butadiene, η-cyclopentadienyl, and related derivatives, 423–34
Reactions of co-ordinated ligands. Part 33. Mononuclear η^2-vinyl complexes: synthesis, structure, and reactivity, 435–50
The synthesis of $NR_4[(Ta_6Cl_{12})(H_2O)_6]X_4$ (R = Me or Et, X = Cl or Br) by the spontaneous reduction of $[Ta_6Cl_{12}]^{4+}$ to $[Ta_6Cl_{12}]^{3+}$ in acidic media: *X*-ray structure analysis of $NMe_4[(Ta_6Cl_{12})(H_2O)_6]Br_4$, 455–8
Co-ordination of alkali metals by open-chain polyethers in transition metal complexes. Part 4. Variation in alkali-metal ion selectivity in cobalt and zinc complexes of 1-(*o*-carboxymethoxyphenoxy)-2-(*o*-hydroxyphenoxy)ethane (HL) and the *X*-ray and molecular structure of $[Zn(NH_4L_2)_2]$, 459–64
X-Ray crystal structures and properties of tris(tetrathiafulvalenium) tetrachlorodimethylstannate(IV) and tetrathiafulvalenium trichlorodimethylstannate (IV), 465–70
The structure of pentach' rocyclopentadienylphenylmercury(II) in the solid state and in solution; the effect of *X*-ray absorption and crystal decay on molecular geometry parameters, 471–4
Macrocyclic polyphosphane ligands. Iron(II), cobalt(II), and nickel(II) complexes of (4*RS*,7*RS*,13*SR*,16*SR*)-tetraphenyl-1,10-dipropyl-1,10-diaza-4,7,13,16-tetraphosphacyclo-octadecane: crystal structures of their tetraphenylborate derivatives, 479–86
Investigation into diphosphine oxides as ligands in diorganotin(IV) adducts. Part 3. Synthesis and crystal structure of two adducts of dinitratodiphenyltin(IV) with *cis*- and *trans*-1,2-bis(diphenylphosphoryl)ethylene, 487–92
Glycine complexation with uranyl ion: absorptiometric, luminescence, and *X*-ray structural studies of tetrakis-(glycine)dioxouranium(VI) nitrate, 517–22
Chemistry of metallacyclobutanones. Part 3. Reactions of heptane-2,4,6-trione and 1,5-diphenylpentane-1,3,5-trione with some carbonate complexes of platinum(II); *X*-ray crystal structures of $[Pt\{CH(COMe)COCH(COMe)\}(PPh_3)_2]$ and $[Pt\{OC(CHCOPh)CHC(Ph)O\}(PPh_3)_2]$, 549–54
Systematic synthesis of tetranuclear osmium clusters by the reaction of trinuclear clusters with $[OsH_2(CO)_4]$; crystal structure of $[Os_4H_3Br(CO)_{13}]$, 555–64
Structural and nuclear magnetic resonance studies of short selenium–nitrogen bonds, 565–72
Synthesis, characterization, and structure of the complex $[FeH(H_2BH_2)\{CH_3C(CH_2PPh_2)_3\}]$, 605–10
Complexes of the platinum metals. Part 23. Synthesis of the nitrosyl carboxylate complexes $[M(O_2CR)_2(NO)(PPh_3)_2]$ (M = Rh or Ir; R = CF_3, C_2F_5, or C_6F_5): crystal and molecular structures of the trifluoroacetate derivatives $[M(O_2CCF_3)_2(NO)(PPh_3)_2]$, 611–6
Complexes of the platinum metals. Part 25. Reactions of ruthenium and osmium nitrosyls $[MH(NO)(PPh_3)_3]$ and $[M(NO)_2(PPh_3)_2]$ with perfluorocarboxylic acids: *X*-ray crystal structure determination of nitrosyl(trifluoroacetato)-(trifluoroacetohydroximato-*OO′*)bis(triphenylphosphine)osmium(II)–dichloromethane (1/1), 621–8
Synthesis and *X*-ray structural studies on the cluster compounds $[RuRh_3(\mu_3\text{-}CO)_2(CO)_3(\eta\text{-}C_5Me_5)_3]$ and $[RuRh_2(\mu\text{-}CO)(\mu_3\text{-}CO)(CO)_2(\eta^4\text{-}C_8H_{10})(\eta\text{-}C_5Me_5)_2]$, 645–50
Mono-η-cycloheptatrienyltitanium chemistry: synthesis, molecular and electronic structures, and reactivity of the complexes $[Ti(\eta\text{-}C_7H_7)L_2X]$ (L = tertiary phosphine, O- or N-donor ligand; X = Cl or alkyl), 669–84
Synthesis and characterisation of 1,2-bis(dimethylphosphino)ethane (dmpe) complexes of chromium-(0) and -(IV): *X*-ray crystal structures of *trans*-$Cr(N_2)_2(dmpe)_2$, *cis*-$Cr(CO)_2(dmpe)_2$, $Cr(C_2Ph_2)_2(dmpe)$, and $CrH_4(dmpe)_2$, 685–92
Reduction–oxidation properties of organotransition-metal complexes. Part 20. Oxidative and thermolytic cyclopropane ring-opening reactions; *X*-ray crystal structure of $[Fe_2(CO)_6(\eta^4\text{:}\eta^{\prime 4}\text{-}C_{16}H_{18})]$, 699–706
Spectroscopic and magnetic properties of cobalt(II) and nickel(II) clusters obtained from 1-(hydroxymethyl)-3,5-dimethylpyrazole. *X*-Ray structure of tetrakis[chloro(μ_3-3,5-dimethyl-*N*-oxymethylpyrazolato-N^2,μ_3-*O*)(ethanol)nickel(II)], 737–42
Short hydrogen bonds: diadducts of substituted pyridine *N*-oxides: synthesis, spectroscopic studies, and *X*-ray structure, 749–54
Carbaborane derivatives of the late- and post-transition elements. Part 3. Structural consequences of ligand substitution in palladadicarbadodecaboranes 3-L_2-3,1,2-$PdC_2B_9H_{11}$. The crystal and molecular structures of 3-$[Me_2N(CH_2)_2NMe_2]$-3,1,2-$PdC_2B_9H_{11}$ and 3-$(PMe_3)_2$-3,1,2-$PdC_2B_9H_{11}$, 761–70
Solvent extraction of gold and platinum-group metals using 2-nonylpyridine 1-oxide, and the crystal and molecular structure of bis(2-nonylpyridine 1-oxide)hydrogen(1+) tetrachloroaurate(III), 771–6
Annelation of ring-opened arylcyclopropenium ions to co-ordinated cyclo-octatetraene, and the *X*-ray crystal structure of $[Fe(CO)_3(\sigma,\eta^3\text{-}C_{11}H_9Ph_3)]$, 777–82
Molecular structure and solid-state properties of the two-dimensional conducting mixed-valence complex $[NBu_4]_{0.29}[Ni(dmit)_2]$ and the neutral $[Ni(dmit)_2]$ (H_2dmit = 4,5-dimercapto-1,3-dithiole-2-thione); members of an electron-transfer series, 783–94
Platinum(II) complexes of nitroimidazoles: synthesis, characterisation, and *X*-ray crystal structures of *cis*-dichlorobis[1-(2′-hydroxyethyl)-2-hydroxymethyl-5-nitroimidazole)platinum(II) and *trans*-dichlorobis[1-(2′-hydroxy-3′-methoxypropyl)-2-nitroimidazole]platinum(II), 795–802
Palladium(II,IV) mixed-valence complexes of 1,2-diaminoethane, 1,3-diaminopropane, and diethylenetriamine: syntheses, electronic, infrared, Raman, and resonance Raman spectra and *X*-ray studies, 815–20
Lewis-base adducts of group 1B metal(I) compounds. Part 16. Synthesis, structure, and solid-state phosphorus-31 nuclear magnetic resonance spectra of some novel $[Cu_4X_4L_4]$ (X = halogen, L = N,P base) 'cubane' clusters, 831–8
Lewis-base adducts of group 1B metal(I) compounds. Part 17. Synthesis and crystal structures of adducts of copper(I) cyanide with nitrogen bases, 839–44
Interconversion of 42- and 44-electron platinum *triangulo*-clusters using chelating tertiary phosphine ligands and the structural characterisation of [1,3-bis(diphenylphosphino)propane]-tris(μ-sulphur dioxide)bis(tricyclohexylphosphine)triplatinum–benzene (1/2), $[Pt_3(\mu\text{-}SO_2)_3\{P(C_6H_{11})_3\}_2(dppp)]\cdot 2C_6H_6$, 845–50
Chemistry of platinum sulphido-complexes. Part 5. Synthesis and crystal and molecular structure of 3-(η-cyclo-octa-1,5-diene)bis(μ_3-sulphido)-1,1,2,2-tetrakis(triphenylphosphine)diplatinum(II)rhodium(I) hexafluorophosphate–dichloromethane (1/1), $[Pt_2Rh(\mu_3\text{-}S)_2(PPh_3)_4(\eta\text{-}C_8H_{12})]PF_6\cdot CH_2Cl_2$, 851–6

CRYSTAL STRUCTURE (contd)

Correlation between structure and circular dichroism. Structure and absolute configuration of the (−) isomer of lithium (ethylenediamine-*N*,*N*′-diacetato-*N*,*N*′-di-3-propionato)rhodate(III) pentahydrate, 861–4

Complexes of lanthanoid salts with macrocyclic ligands. Part 17. Synthesis and crystal and molecular structure of a hydroxide-bridged praseodymium trifluoroacetate complex with 15-crown-5-ether, $[Pr_2(CF_3CO_2)_3(OH)(C_{10}H_{20}O_5)_2]$-$[Pr_2(CF_3CO_2)_6]$, 885–90

Formation of substituted cyclopentadienyl ligands on tungsten *via* reactions between the alkyne complexes $[W(CO)(R^1C_2R^2)_3]$ ($R^1 = R^2$ = Ph or Et; R^1 = Me, R^2 = Ph) and the alkylidyne compounds $[W(\equiv CR)(CO)_2(\eta\text{-}C_5H_5)]$ (R = C_6H_4Me-4 or Me); *X*-ray crystal structures of $[W_2(\mu\text{-}CO)_2(CO)(\eta\text{-}PhC_2Ph)(\eta\text{-}C_5Ph_4R)(\eta\text{-}C_5H_5)]$ and $[W_2(\mu\text{-}EtC_2Et)(CO)_4(\eta\text{-}C_5Et_4R)(\eta\text{-}C_5H_5)]$ (R = C_6H_4Me-4), 905–12

Crystal and molecular structures and magnetic properties of bromo(3-dimethylaminopropan-1-olato)copper(II), chloro(3-dimethylaminopropan-1-olato)copper(II), and (3-diethylaminopropan-1-olato)isocyanatocopper(II), 913–20

Alkyl, hydrido, and tetrahydroaluminato complexes of manganese with 1,2-bis(dimethylphosphino)ethane (dmpe). *X*-Ray crystal structures of $Mn_2(\mu\text{-}C_6H_{11})_2(C_6H_{11})_2(\mu\text{-dmpe})$, $(dmpe)_2Mn(\mu\text{-}H)_2AlH(\mu\text{-}H)_2Mn(dmpe)_2$, and $Li_4\{MnH(C_2H_4)[CH_2(Me)PCH_2CH_2PMe_2]_2\}_2\cdot 2Et_2O$, 921–30

Reaction of transition-metal carbonylate anions and 1,1,1-tris(halogenomethyl)ethane. *X*-Ray crystal structures of tricarbonyl(η^5-cyclopentadienyl)(1-methylcyclopropylmethyl)tungsten(II), and tetraethylammonium enneacarbonyliododirhenate(0), 931–40

Reactions of 6,6′-bis(*nido*-decaboranyl) oxide and 6-hydroxy-*nido*-decaborane with dihalogenobis(phosphine) complexes of nickel, palladium, and platinum, and some related chemistry; nuclear magnetic resonance investigations and the crystal and molecular structures of bis(dimethylphosphine)-di-μ-(2,3,4-η^3-*nido*-hexaboranyl)-diplatinum(*Pt–Pt*), $[Pt_2(\mu\text{-}\eta^3\text{-}B_6H_9)_2(PMe_2Ph)_2]$, and of 2,4-dichloro-1,1-bis(dimethylphenylphosphine)-*closo*-1-nickeladecaborane, $[(PhMe_2P)_2NiB_9H_7Cl_2]$, 953–72

Pyrazolate A-frame rhodium complexes. Crystal structures of $[Rh_2(\mu\text{-dmpz})(CO)_2(\mu\text{-dppm})_2][ClO_4]$ and $[Rh_2(\mu\text{-dmpz})I_2(CO)_2(\mu\text{-dppm})_2][ClO_4]$, 973–80

Co-ordination chemistry of pyridyl and *N*-methylimidazolyl ketones. Synthetic and *X*-ray structural studies of copper(II), nickel(II), and dimethylgold(III) complexes, 981–6

Actinide structural studies. Part 7. The crystal and molecular structures of (2,2′-bipyridyl)dinitratodioxo-uranium(VI) and -neptunium(VI), and diacetato(2,2′-bipyridyl)dioxo-uranium(VI) and -neptunium(VI), 1001–8

Reduction–oxidation properties of organotransition-metal complexes. Part 21. Synthesis and *X*-ray structural characterisation of the redox-related pair of cyclohexadienyl complexes $[Mn(CO)(dppe)(\eta^5\text{-}C_6H_6Ph)]$ and $[Mn(CO)(dppe)(\eta^5\text{-}C_6H_6Ph)][PF_6]\cdot 0.5CH_2Cl_2$, 1019–26

Reduction–oxidation properties of organotransition-metal complexes. Part 22. Stereospecific oxidative cyclopropane ring opening and reductive cyclobutane ring formation in polycyclic hydrocarbon complexes of iron; *X*-ray crystal structures of $[Fe_2(CO)_6(\eta^5\text{:}\eta'^5\text{-}C_{16}H_{16})][PF_6]_2\cdot CH_3NO_2$ and $[Fe_2(CO)_6(\eta^4\text{:}\eta'^4\text{-}C_{16}H_{16})]$, 1027–36

Reaction of trithiazyl trichloride, $(NSCl)_3$, with triphenylphosphine or triphenylphosphine metal complexes. *X*-Ray crystal structure of aminotriphenylphosphonium chloride–dichloromethane (1/1), $[Ph_3PNH_2]Cl\cdot CH_2Cl_2$, 1043–8

Platinum(II) complexes containing a cationic amine ligand: crystal structure of [(2-aminoethyl)ammonium]trichloroplatinum(II), 1057–60

Preparation and crystal structure of $Na_4[Hg_4(\mu\text{-}Cl)_4\{P(O)(OEt)_2\}_8]\cdot 6H_2O$, 1061–4

Tetrachlorobis(tetrahydrofuran)molybdate(III), a new intermediate for the synthesis of molybdenum(III) complexes: preparation and crystal structure, 1069–72

N,*N*′-Disubstituted dithiomalonamide complexes of antimony(III). Crystal and molecular structure of $[SbCl_3\{C_2H_5NHC(S)CH_2C(S)NHC_2H_5\}]$ with lone-pair occupation of an antimony co-ordination site, 1073–6

Carbon–carbon formation at di-iron centres. Part 2. Reactivity of $[Fe_2(CO)_6(\mu\text{-}COEt)\{\mu\text{-}C(R)C(R)H\}]$ complexes toward $MeOC(O)C{\equiv}CC(O)OMe$ (R = Ph) and $CF_3C{\equiv}CCF_3$ (R = Ph or H); *X*-ray crystal structures of $[Fe_2(CO)_5\{\mu\text{-}C(OEt)C[C(O)OMe]C[C(O)OMe]\}\{\mu\text{-}C(Ph)C(Ph)H\}]\cdot H_2O$ and $[Fe_2(CO)_6\{\mu\text{-}C(CF_3)C(CF_3)CHCHCH(OMe)\}]$, 1087–94

Polyhedral rhenaborane chemistry: crystal and molecular structures of the *nido*-6-rhenadecaborane cluster compounds [6,6,6,6-$(PMe_2Ph)_3$H-*nido*-6-ReB_9H_{13}] and [2-(PMe_2Ph)-6,6,6,6-$(PMe_2Ph)_2$ClH-*nido*-6-ReB_9H_{12}]; nuclear magnetic resonance parameters of these and other related *nido*-rhenadecaborane cluster species, 1119–30

New carbide clusters in the cobalt sub-group. Part 14. Synthesis and structural characterization of the anion $[Co_{13}C_2(CO)_{24}]^{3-}$ as its benzyltrimethylammonium salt, 1137–40

Dynamic behaviour of carbon-metallated palladium hydrazone complexes. Crystal structures of $[\{Pd[CH_2CMe_2C(=N\text{–}NMePh)Me]Cl\}_2]$ and $[\{Pd[CH_2C(=N\text{–}NMePh)Bu^t]Cl\}_2]$, 1155–62

Synthesis of gold-(I) and -(III) complexes with carbonyl-stabilized phosphorus ylides. Crystal structure of $[\{Au(PPh_3)\}_2\{\mu\text{-}C(PPh_3)CO_2Et\}]ClO_4$, 1163–8

Macrocyclic polyphosphane ligands. Cobalt(II) and nickel(II) complexes of 5*RS*,8*RS*,16*RS*,19*RS*-tetraphenyl-5,8,16,19-tetraphospha-1,12-dithiacyclododecosane (δ-L^1) and the crystal structure of $[Co(\delta\text{-}L^1)][BF_4]_2\cdot 0.5H_2O$, 1179–82

Co-ordination chemistry of dimethylgold(III). Synthesis, structural studies, and fluxional behaviour of complexes with polydentate ligands, 1183–90

Cluster chemistry. Part 31. An $[Ru_3(CO)_{12}]$ derivative with an $[Fe_3(CO)_{12}]$-like structure: preparation and *X*-ray structure of $[Ru_3(\mu\text{-}CO)_2(CO)_6\{PPh(OMe)_2\}_4]$, 1223–8

Cluster chemistry. Part 32. Synthesis and *X*-ray crystal structure of $[Ru_5(\mu_5\text{-}\eta^2\text{-}C_2PPh_2\text{-}P)(\mu\text{-}PPh_2)(CO)_{13}]$, a complex containing an alkynyl ligand in extended interaction with an open Ru_5 cluster, 1229–34

From diarylruthenium complexes to *ortho*-metallated ketones: a mechanistic and crystal structure study, 1235–42

Crystal and molecular structures and magnetic properties of four new exchange-coupled copper(II) complexes derived from different 3-*N*,*N*-dialkylamino-1-propanols and pseudohalogens, 1243–8

The crystal structure of diaquadichlorodimethyltin(IV)–purine (1/4), 1271–4

Homo- and hetero-dinuclear hydride-bridged complexes containing cyclo-octadiene: the crystal and molecular structure of $[(\eta^4\text{-}C_8H_{12})Ir(\mu\text{-}H)(\mu\text{-}Cl)IrH_2(PPh_3)_2]$, 1277–80

Characterization of the adducts formed by Cu(CN) and Cu(NCS) with biquinoline. The crystal structure of the polymeric cyano-compound containing both linear and tetrahedrally co-ordinated copper(I), $[\{Cu_3(bq)_2(CN)_3\}_n]$, 1285–8

Crystal structures and interrelationships of the blue and green conformational isomers of tetrakis(trimethylarsine sulphide)cobalt(II) perchlorate, 1289–94

Fluorophosphine complexes of rhodium(I) and iridium(I): towards the design of systems with extended metal–metal interactions. The crystal structure of $[\{IrCl(PF_3)_2\}_2]$, 1295–302

Synthesis, structures, and reactivities of some pentamethylcyclopentadienyl–sulphur compounds, 1303–8

New carbide clusters in the cobalt sub-group. Part 15. Synthesis and crystallographic characterization of di-μ_6-carbido-deca-μ-carbonyl-tridecacarbonyl-*polyhedro*-dodecarhodate(4−) as its tetrapropylammonium salt, $[N(C_3H_7)_4]_4\text{-}[Rh_{12}C_2(CO)_{23}]$, 1309–14

Chemistry of di- and tri-metal complexes with bridging carbene or carbyne ligands. Part 33. Reactions of $[W(\equiv CMe)(CO)_2(\eta\text{-}C_5H_5)]$ with the dimetal compounds $[MRh(\mu\text{-}CO)_2(\eta\text{-}C_5Me_5)_2]$ (M = Co or Rh); *X*-ray crystal structure of $[Rh_2W(\mu\text{-}CO)(\mu_3\text{-}CMe)(CO)_2(\eta\text{-}C_5H_5)(\eta\text{-}C_5Me_5)_2]$, 1315–22

Chemistry of di- and tri-metal complexes with bridging carbene or carbyne ligands. Part 34. Reactions of sulphur and selenium with the di-irontungsten complexes $[Fe_2W(\mu_3\text{-}CR)(\mu\text{-}CO)(CO)_8(\eta\text{-}C_5H_5)]$ (R = C_6H_4Me-4 or Me); crystal structures of $[Fe_2W(\mu\text{-}CC_6H_4Me\text{-}4)(\mu_3\text{-}S)(CO)_7(\eta\text{-}C_5H_5)]$ and $[Fe_2W(\mu_3\text{-}SCMe)(CO)_8(\eta\text{-}C_5H_5)]$, 1323–30

Chemistry of di- and tri-metal complexes with bridging carbene or carbyne ligands. Part 35. The synthesis and crystal structures of the compounds $[N(PPh_3)_2][W_2\{\mu\text{-}C(H)C_6H_4Me\text{-}4\}(CO)_7(\eta\text{-}C_5H_5)]\cdot 0.5Et_2O$ and $[W\{=C(H)C_6H_4Me\text{-}4\}\text{-}(SnPh_3)(CO)_2(\eta\text{-}C_5H_5)]$, 1331–8

Synthesis and properties of the divalent 1,2-

CRYSTAL STRUCTURE (contd)
bis(dimethylphosphino)ethane (dmpe) complexes $MCl_2(dmpe)_2$ and $MMe_2(dmpe)_2$ (M = Ti, V, Cr, Mn, or Fe). *X*-Ray crystal structures of $MCl_2(dmpe)_2$ (M = Ti, V, or Cr), $MnBr_2(dmpe)_2$, $TiMe_{1.3}Cl_{0.7}(dmpe)_2$, and $CrMe_2(dmpe)_2$, 1339–48
Reaction of bis(morpholinothiocarbonyl) disulphide with iodine. Existence of a 1:1 charge-transfer precursory adduct in an oxidation reaction. Isolation and crystal structure of bis[3,5-di(*N*-morpholinio)-1,2,4-trithiolane] hexadecaiodide, 1349–54
The reaction of $[Os_3(\mu\text{-}H)_2(CO)_9L]$ (L = CO or PEt_3) with dimethylcyanamide, Me_2NCN: *X*-Ray crystal structure of $[Os_3(\mu\text{-}H)(\mu\text{-}NCHNMe_2)(CO)_{10}]$ and the reactions of this complex with acids, 1355–60
Bromine nuclear quadrupole resonance studies of some hexabromostannates: *X*-Ray crystal structure of pyridinium hexabromostannate(IV), $[Hpy]_2[SnBr_6]$, 1399–404
Preparation of 1,3,2-dithiazolium hexafluoroarsenate(V), preparation and crystal structures of 5-methyl-1,3,2,4-dithiadiazolium and 4-methyl-1,3,2-dithiazolium hexafluoroarsenate(V) and the reduction of these salts to stable free radicals, 1405–16
Macrocyclic polyphosphane ligands. Cobalt(II) and nickel(II) complexes of the γ and ε diastereoisomers of 4,7,13,16-tetraphenyl-4,7,13,16-tetraphospha-1,10-dithiacyclo-octadecane (L^1) and the crystal structure of $[Co(\varepsilon\text{-}L^1)][BPh_4]_2$· EtOH, 1425–30
Reactions of bis(cyclopentadienyl)vanadium derivatives with nitrogen mono-oxide and the crystal structure of an oxo-bridged nitrosyl complex of vanadium, 1435–42
Compartmental ligands. Part 11. Copper(II) complexes of 'half-unit' Schiff bases: crystal and molecular structures of one monomeric and of one dimeric complex, 1471–8
Bimetallic cyano-bridged cations: preparation and hydride reduction of $[(\eta^5\text{-}C_5H_5)L_2Ru(\mu\text{-}CN)ML'_2(\eta^5\text{-}C_5H_5)]PF_6$ [L_2, L'_2 = $(PPh_3)_2$, $Ph_2PCH_2CH_2PPh_2$; M = Ru or Fe]. Formation of $[Ru(\eta^5\text{-}C_5H_5)(PPh_3)H_2]$ and *X*-ray crystal structure of $[(\eta^5\text{-}C_5H_5)(Ph_2PCH_2CH_2PPh_2)Ru(\mu\text{-}CN)Ru(PPh_3)_2(\eta^5\text{-}C_5H_5)]PF_6$, 1479–86
Benzamidinatorhodium complexes. *X*-Ray structures of $[Rh\{CPh(NPh)_2\}(cod)]$ and $[Rh_2\{\mu\text{-}CPh(NPh)_2\}_2(tfbb)_2]$, 1487–94
Reactivity of the unsaturated anion decacarbonyltetra-μ-hydrido-trirhenate(1−) toward phenols. Crystal and molecular structures of the tetraethylammonium salts of the triangular cluster anion $[Re_3(\mu\text{-}H)_3(\mu\text{-}OC_6F_5)(CO)_{10}]^-$ and of the binuclear anion $[Re_2(\mu\text{-}OC_6H_3)_3(CO)_6]^-$, 1507–12
Preparation and properties of *cis*-$[W(N_2)_2(Ph_2PCH_2PPh_2)_2]$ and *trans*-$[M(N_2)_2(Ph_2PCH{=}CHPPh_2)_2]$ (M = Mo or W), and the crystal structure of *cis*-$[W(N_2)_2(PMe_2Ph)_4]$, 1523–6
Synthesis and structural studies of some 1,1′-dichloroferrocene derivatives of platinum(II). Crystal and molecular structure of 2,2-μ-[(1—2,5—6-η-*cis*,*cis*-cyclo-octa-1,5-diene)platinio]-bis(1,1′-dichloroferrocene), 1527–30
The synthesis of $NR_4[(Ta_6Cl_{12})(H_2O)_6]X_4$ (R = Me or Et, X = Cl or Br) by the spontaneous reduction of $[Ta_6Cl_{12}]^{4+}$ to $[Ta_6Cl_{12}]^{3+}$ in acidic media: *X*-ray structure analysis of $NMe_4[(Ta_6Cl_{12})(H_2O)_6]Br_4$ (1985, 455), 1531–2
Rhenium nitrosyl complexes with simple and with sterically demanding aromatic thiolate ligands: *X*-ray crystal structures of $[PPh_4][Re_2(SC_6H_4Me\text{-}4)_7(NO)_2]\text{-}CH_2Cl_2$ and $[Re(SC_6H_3Pr^i_2\text{-}2,6)_4(NO)]$, 1533–42
Structure and solvation of mercury(II) iodide, bromide, and chloride in pyridine solution; refinement of the crystal structure of di-iodobis(pyridine)mercury(II), $[HgI_2(py)_2]$, 1597–604
The preparation and crystal structure of (β)6SbF_3·5SbF_5, 1623–32
Synthesis and crystal structure of the layer compound Sb_3TeO_6Cl, 1633–6
Studies of 2,5;6,10;8,10-tri-μ-hydro-nonahydro-*nido*-nonaborate(1−), $[B_9H_{12}]^-$: preparation, crystal and molecular structure, nuclear magnetic resonance spectra, electrochemistry, and reactions, 1645–54
Bimetallic systems. Part 11. Heterobimetallic and unsymmetrical diplatinum complexes from *cis*-$[PtR_2(dppm\text{-}P)_2]$ (dppm = $Ph_2PCH_2PPh_2$; R = Me, 1-naphthyl, or $C_6H_4Me\text{-}o$): crystal structure of $[(C_6H_4Me\text{-}o)_2Pt(\mu\text{-}dppm)_2PtMe_2]$, 1677–82
Synthesis and structural characterisation of some *triangulo*-platinum clusters containing isocyanide ligands, 1693–8
Transition-metal co-ordination compounds of a novel aniline-based pyrazole derivative. *X*-Ray crystal structures of [*NN*-bis(3,5-dimethylpyrazol-1-ylmethyl)aminobenzene]-dichlorocobalt(II) and -dibromocopper(II), 1699–706
Sila-pharmaca. Part 32. Crystal and molecular structures of the (*R*)-enantiomer and the racemate of the antimuscarinic agent (cyclohexyl)phenyl[2-(pyrrolidin-1-yl)ethyl]silanol (sila-procyclidine), 1743–6
Axially asymmetric metal alkyls. Part 3. Chemical, electrochemical, and structural studies of group 5A $d^{0,1,2}$ metallepines $[M\{(2\text{-}CH_2C_6H_4)_2\}(\eta\text{-}C_5H_5)_2]^z$ (M = Nb or Ta; $z = 1-$, 0, or 1+); synthesis of $[Nb\{2\text{-}CH_2(C_6H_4)_2Me\text{-}2'\}X(\eta\text{-}C_5H_5)_2]$ (X = Cl or CO) and $[\{(\eta\text{-}C_5H_5)_2ClV\}_2\{(2\text{-}CH_2C_6H_4)_2\}]$, and electrochemistry of $[M(CH_2Ph)_{2-n}Cl_n(\eta\text{-}C_5H_5)_2]$ (n = 0, M = Nb, or Ta; n = 1, M = V), 1747–60
The structure of a dinuclear copper(I) complex of a Schiff-base ligand containing a copper–copper bond, 1771–4
Dirhodium(II,II) tetra-acetate complexes with axially co-ordinated triphenylstibine, triphenylarsine, and dibenzyl sulphide ligands. The syntheses, properties, and *X*-ray crystal structures of $[Rh_2(O_2CMe)_4(SbPh_3)_2]$, $[Rh_2(O_2CMe)_4(AsPh_3)_2]$, and $[Rh_2(O_2CMe)_4\{S(CH_2Ph)_2\}_2]$, 1775–80
Reactions of some decaosmium clusters with electrophilic and nucleophilic reagents: *X*-ray structure analyses of $[N(PPh_3)_2][Os_{10}C(CO)_{24}(\mu\text{-}I)]$, $[Os_{10}C(CO)_{24}(\mu\text{-}I)_2]$, $[N(PPh_3)_2]_2[Os_{10}C(CO)_{22}(NO)I]$, $[Os_{10}C(CO)_{23}\{P(OMe)_3\}(\mu\text{-}I)_2]$ and of two isomers of $[Os_{10}C(CO)_{21}\{P(OMe)_3\}_4]$, 1795–810
High-co-ordination-number compounds of niobium and tantalum: Reactions of niobium and tantalum halides and sulphido-halides with sodium diethyldithiocarbamate. The crystal structures of $[Nb(S_2CNEt_2)_4]Br$, $Nb(S_2CNEt_2)_3S$, and $Ta(S_2CNEt_2)_3(S_2)$, 1821–8
Studies in the flexibility of macrocycle ligands. Calculation of macrocycle cavity size by force-field methods. Crystal and molecular structures of $[CoLCl][ClO_4]_2$ and $[CuL][PF_6]_2$ {L = 2,13-dimethyl-3,6,9,12,18-penta-azabicyclo[12.3.1]octadeca-1(18),14,16-triene}, 1829–34
Reaction between $[Ru_3(CO)_{12}]$ and the tridentate phosphine ligand $HC(PPh_2)_3$; *X*-ray crystal structures of complexes $[Ru_3(CO)_9\{Ph_2PCHP(Ph)C_6H_4PPh\}]$, $[Ru_2H(CO)_4(Ph_2PCHPPh_2)\{PhPC_6H_4C(O)\}]$, and $[Ru_2(CO)_4Cl(PPh_2)(dppm)]$, 1835–42
The crystal and molecular structure of $[1,1,1,1\text{-}(CO)H(PPh_3)_2\text{-}$*arachno*-1-$IrB_3H_7]$ and some bonding considerations in *arachno*-type four-vertex metal 'π-allyl' and 'borallyl' clusters, 1843–8
Studies of phosphazenes. Part 25. Synthesis, nuclear magnetic resonance spectroscopy, and mode of formation of (aziridino)(triphenylphosphazenyl)cyclotriphosphazenes. *X*-ray crystal structure and enzyme-inhibiting activity of $N_3P_3(NPPh_3)(NC_2H_4)_5$, 1881–90
1*H*-pyrrolo[2,3-*b*]pyridine (HL) ligands in rhodium(I) and iridium(I) chemistry. Crystal and molecular structures of $[Rh_2(\mu\text{-}L)_2(nbd)_2]$ and $[Rh_4(\mu\text{-}Cl)_2(\mu\text{-}L)_2(\mu\text{-}CO)_2(CO)_2(nbd)_2]$, 1891–8
Magnetic behaviour of tetrakis[(2-diethylaminoethanolato)isocyanatocopper(II)], a complex with an antiferromagnetic ground state; the crystal and molecular structure of the triclinic modification, 1909–14
Complexes of organoaluminium compounds. Part 13. Preparation and nuclear magnetic resonance spectra of the arylamido-compounds $AlMe_2(NHR')$ (R′ = Ph, $C_6H_4Me\text{-}o$, $C_6H_4Me\text{-}p$, $C_6H_3Me_2\text{-}2,6$) and the imido-compounds AlMe(NR′). Crystal and molecular structures of $[\{AlMe_2(NHC_6H_4Me\text{-}o)\}_2]$ and $[\{AlMe(NPh)\}_6]$, 1929–34
Organic chemistry of dinuclear metal centres. Part 8. Organo-iron–ruthenium chemistry. *X*-Ray structure of *trans*-$[FeRu(CO)_2(\mu\text{-}CO)_2(\eta\text{-}C_5H_5)_2]$, 1935–44
Crystal structures and magnetic properties of binuclear five-co-ordinate copper(II) complexes with a phenolate bridge and their catalytic functions in multielectron redox reactions, 1945–52
Isolation and crystal structure of $[Rh(PPh_3)_3(MeCN)][BF_4]$, acetonitriletris(triphenylphosphine)rhodium(I) tetrafluoroborate, 1977–80
Reactivity of $[PPh_4][Fe_2(CO)_6\{\mu\text{-}CPhCPhC(CF_3)C(CF_3))H\}]$ toward electrophiles. *X*-Ray structure of $[Fe_2(CO)_6\text{-}\{\mu\text{-}CPhCPhC(CF_3)CHC(OEt)_2\}]$, a product resulting from fluorine abstraction at a CF_3 group and subsequent fluorine substitution by ethoxy-groups, 1981–4
New carbonyl derivatives of niobium(I) and tantalum(I), 1989–96
Routes of formation and crystal structure of an alkylperoxycobaloxime: bis[dimethylglyoximato(1−)](4-

CRYSTAL STRUCTURE (contd)
ethoxycarbonylbut-3-en-2-ylperoxo)(pyridine)cobalt(III), 1997–2000
Chemistry of di- and tri-metal complexes with bridging carbene or carbyne ligands. Part 36. Reactions of the dimetal compounds [ReM(≡CC$_6$H$_4$Me-4)(CO)$_9$] (M = Cr, Mo, or W) with octacarbonyldicobalt; crystal structures of [Co$_2$WRe(μ$_3$-CC$_6$H$_4$Me-4)(CO)$_{15}$] and [Co$_2$Re(μ$_3$-CC$_6$H$_4$Me-4)(CO)$_{10}$], 2001–8
Chemistry of di- and tri-metal complexes with bridging carbene or carbyne ligands. Part 37. Methylene group transfer to carbon–metal multiple bonds; crystal structures of [TiW{μ-C(C$_6$H$_4$Me-4)=CH$_2$}(μ-CO)(CO)(η-C$_5$H$_5$)$_2$] and [PtW{μ-C(C$_6$H$_4$Me-4)=CH$_2$}(CO)$_2$(PMe$_2$Ph)$_2$(η-C$_5$H$_5$)], 2009–16
Chemistry of di- and tri-metal complexes with bridging carbene or carbyne ligands. Part 38. Ruthenium–tungsten compounds: crystal structures of [RuW(μ-Cl)(μ-CMe)(Cl)(CO)$_2$(PPh$_3$)$_2$(η-C$_5$H$_5$)] and [RuW$_2$(μ$_3$-C$_2$Me$_2$)(CO)$_7$(η-C$_5$H$_5$)$_2$], 2017–24
Trimethylphosphine as a reactive solvent: synthesis, crystal structures, and reactions of [Ta(PMe$_3$)$_3$(η^2-CH$_2$PMe$_2$)(η^2-CHPMe$_2$)] and [W(PMe$_3$)$_4$(η^2-CH$_2$PMe$_2$)H] and related studies, 2025–36
Synthesis, structure, and bonding of fulvene complexes of titanium, molybdenum, and tungsten, 2037–50
Metal complexes of vitamin B$_6$ related compounds. Crystal and molecular structures of aqua(5′-phosphopyridoxylideneglycinato)copper(II) trihydrate and bis(pyridoxylideneglycinato)nickel(II) hexahydrate, 2051–8
Preparation and properties of *mer*-[ReCl(N$_2$)(CNR){P(OMe)$_3$}$_3$] (R = Me, Et, Bu^t, C$_6$H$_4$Me-4, or C$_6$H$_4$Cl-4) and [ReCl(N$_2$)(CNMe)(PPh$_3$){P(OEt)$_3$}$_2$]. *X*-Ray crystal structure of *mer*-[ReCl(N$_2$)(CNMe){P(OMe)$_3$}$_3$] and reductive cleavage of the isocyanide ligands to primary amines upon protonation, 2079–84
X-Ray crystal structures and magnetic properties of azide-bridged binuclear copper(II) complexes containing the Schiff-base ligand derived from 2-pyridinecarbaldehyde and histamine. Structure–magnetism relationship, 2095–100
Complexes of the platinum metals. Part 29. Pyridine-2-thiolate derivatives of ruthenium and osmium: *X*-ray crystal structures of [Ru(C$_5$H$_4$NS)$_2$(CO)$_2$(PPh$_3$)] and [Ru(C$_5$H$_4$NS)$_2$(CO)(PPh$_3$)], 2101–12
Bimetallic systems. Part 12. Mixed rhodium(I)–platinum(II) acetylide complexes containing bridging Ph$_2$PCH$_2$PPh$_2$. Crystal structures of [(MeC≡C)Pt(μ-dppm)$_2$(σ,η-C≡CMe)Rh(CO)]PF$_6$ and of [ClPt(μ-dppm)$_2$(σ,η-C≡CMe)Rh(CO)]PF$_6$, 2121–30
Bimetallic systems. Part 13. Platinum–manganese carbonyl complexes containing bridging Ph$_2$PCH$_2$PPh$_2$(dppm) ligands: crystal structure of [(OC)$_3$Mn(μ-dppm)$_2$PtH(Br)]BF$_4$, 2131–8
Reactions of co-ordinated ligands. Part 34. Synthesis, structure, and reactivity of cationic dieneruthenium complexes; crystal structures of [Ru(η^4-C$_6$H$_8$)(CO)(η-C$_5$H$_5$)][BF$_4$] and [Ru(η^5-C$_6$H$_9$)(CO)(η-C$_5$H$_5$)], 2145–54
Crystal structure and spectroscopic and redox properties of the iron–sulphur cluster compound [NEt$_4$]$_2$[Fe$_4$S$_4$(SC$_6$H$_4$NH$_2$-4)$_4$], 2161–6
Oxoalkyls of rhenium-(V) and -(VI). *X*-Ray crystal structures of (Me$_4$ReO)$_2$Mg(thf)$_4$, [(Me$_3$SiCH$_2$)$_4$ReO]$_2$Mg(thf)$_2$, Re$_2$O$_3$Me$_6$, and Re$_2$O$_3$(CH$_2$SiMe$_3$)$_6$, 2167–76
Unidentate *versus* symmetrically and unsymmetrically bidentate nitrate co-ordination in pyrazole-containing chelates. The crystal and molecular structures of (nitrato-*O*)[tris(3,5-dimethylpyrazol-1-ylmethyl)amine]copper(II) nitrate, (nitrato-*O*,*O*′)[tris(3,5-dimethylpyrazol-1-ylmethyl)amine]nickel(II)nitrate, and (nitrato-*O*)(nitrato-*O*,*O*′)[tris(3,5-dimethylpyrazol-1-ylmethyl)amine]cadmium(II), 2177–84
Preparation and crystal structure of [AsPh$_4$]$_2$[(WCl$_5$)$_2${μ-NC(CF$_3$)$_2$N}], 2205–8
Synthesis and *X*-ray crystal structure of the asymmetric trinuclear complex [Ni$_3$(μ$_3$-S)$_2$(H$_2$O)(PPh$_3$)$_5$][PF$_6$]$_2$, 2209–12
A disagreement on the explanation of short and long As–O bonds of the (As–O)$_4$ ring in As$_8$(CF$_3$)$_6$O$_6$(OH)$_2$ in terms of As^{III}–O and As^V–O bonds, 2221–2
Reactivity of [NBu$_4$]$_2$[Mo$_2$Br$_6$] with several uni- and poly-dentate phosphines. *X*-Ray structure of [NBu$_4$][MoBr$_4$(Ph$_2$PCH$_2$CH$_2$PPh$_2$)], 2263–8
The palladium-catalysed reaction between [Re$_2$(CO)$_{10}$] and phosphines and the crystal and molecular structure of diaxial [Re$_2$(CO)$_8$(PMe$_2$Ph)$_2$], 2277–82
Stereochemical consequences of the two-electron oxidation of a dirhodium fulvalene complex: the *X*-ray crystal structures of *trans*-[Rh$_2$(CO)$_2$(PPh$_3$)$_2$(η^5:η′^{5}-C$_{10}$H$_8$)] and *cis*-[Rh$_2$(CO)$_2$(PPh$_3$)$_2$(η^5:η′^{5}-C$_{10}$H$_8$)][PF$_6$]$_2$, 2283–90
The synthesis, magnetic, electrochemical, and spectroscopic properties of diruthenium(II,II) tetra-μ-carboxylates and their adducts. *X*-Ray structures of Ru$_2$(O$_2$CR)$_4$L$_2$ (R = Me, L = H$_2$O or tetrahydrofuran; R = Et, L = Me$_2$CO), 2321–6
Copper complexes with quadridentate bis(pyrazolyl)thioether amine and tris(pyrazolyl)amine ligands. Structural characterization of the complexes [Cu(NCS)(tpea)][Cu(NCS)$_2$] and [CuCl(bdma)]Cl·2H$_2$O, 2327–32
Reactivity of [NiR(R′)L$_2$] compounds and the crystal structure of [Ni(C$_2$Cl$_3$)(C$_6$H$_2$Me$_3$-2,4,6)(PMe$_2$Ph)$_2$], 2333–42
Metallaborane chemistry. Part 14.[1] Icosahedral η^6-arene carbametallaboranes of iron and ruthenium; molecular structures of *closo*-[1-(η^6-C$_6$H$_5$Me)-2,4-Me$_2$-1,2,4-FeC$_2$B$_9$H$_9$] and *closo*-[3-(η^6-C$_6$H$_6$)-3,1,2-RuC$_2$B$_9$H$_{11}$], 2343–8
Crystal structures and magnetism of binuclear iron(III) complexes with a linear oxo-bridge, [Fe$_2$O(bbimae)$_2$X$_2$][NO$_3$]$_2$ {bbimae = 2-[bis(benzimidazol-2-ylmethyl)amino]ethanol, X = Cl or NCS}, 2375–80
Investigation into aroylhydrazones as chelating agents. Part 7. Synthesis and spectroscopic characterization of complexes of Mn^{II}, Co^{II}, Ni^{II}, Cu^{II}, and Zn^{II} with 2,6-diacetylpyridine bis(2-aminobenzoylhydrazone) and *X*-ray structure of chloro[2,6-diacetylpyridine bis(2-aminobenzoylhydrazone)](methanol)manganese(II) chloride monohydrate, 2387–92
Trigonal bipyramidal penta-aquazinc(II): crystal structure of penta-aquazinc(II) bis(3,3′,3″-phosphinetriyltripropionato)dizincate(II,II) heptahydrate, 2393–6
Polyhedral ruthenaborane chemistry: characterization of several new ruthenaboranes by nuclear magnetic resonance spectroscopy, and the crystal and molecular structure of [5,6,6-(PPh$_3$)$_3$-6-H-*nido*-6-RuB$_9$H$_{12}$], 2397–406
Two unusual *closo*-type ruthenaboranes: preparation, molecular structure, and nuclear magnetic resonance properties of [1,1,1-(PPh$_3$)HCl-1-RuB$_9$H$_7$-3,5-(PPh$_3$)$_2$] and [1,1-(PPh$_3$)$_2$-1-RuB$_{10}$H$_8$-2,5-(OEt)$_2$], 2407–16
Synthesis of trinuclear gold(I) and gold(III) complexes containing the tridentate bis(diphenylphosphino)methanide ligand. Crystal structure of [Cl(C$_6$F$_5$)$_2$Au{Ph$_2$PCH(AuNC$_5$H$_5$)PPh$_2$}AuCl], 2417–20
Electrophilic nitrosyls: preparation, structure, and reactivity of *cis*-chloronitrosylbis(pyridine-2-carboxylato)ruthenium and related complexes, 2427–32
Crystal structures of the difluorophosphate complexes, Co(O$_2$PF$_2$)$_2$·2MeCN and Cu(O$_2$PF$_2$)$_2$, 2433–6
Synthesis of tetra- and penta-nuclear platinum–osmium carbido-cluster complexes from non-carbido-precursors: *X*-ray structural studies on [Os$_3$Pt(μ-H)$_2$(μ$_4$-C)(CO)$_{10}${P(cyclo-C$_6$H$_{11}$)$_3$}], [Os$_3$Pt$_2$(μ-H)$_2$(μ$_5$-C)(μ-CO)(CO)$_9${P(cyclo-C$_6$H$_{11}$)$_3$}$_2$], and [Os$_3$Pt$_2$(μ-H)(μ$_5$-C)(μ-OMe)(μ-CO)(CO)$_9${P(cyclo-C$_6$H$_{11}$)$_3$}$_2$], 2437–48
Reaction of the anion [WCl$_4$(CBu^t)]$^-$ with tetrasulphur tetranitride. Formation and crystal structure of [AsPh$_4$][WCl$_3$O(OS$_2$N$_2$)], 2453–6
Crystal structure of tris(methylmercurio)sulphonium perchlorate, 2457–8
Reactions of co-ordinated ligands. Part 35. Evidence for carbon–carbon double-bond cleavage of cyclopropenes in their reaction with dinuclear cobalt, rhodium, and iridium complexes; crystal structure and protolysis of [Rh$_2$(μ-CO)(μ-COCHCMe$_2$CH)(μ-C$_5$Me$_5$)$_2$], 2483–92
Kinetics and mechanism of the equilibration reaction between (2,2′,2″-nitrilotriethoxy)nitrosylvanadate(1–) and cyanide. Crystal structures of sodium (2,2′,2″-nitrilotriethoxy)nitrosylvanadate(I)–sodium perchlorate tetrahydrate and of barium cyano(2,2′,2″-nitrilotriethoxy)nitrosylvanadate(I) pentahydrate, 2493–8
The chemistry of heteroarylphosphorus compounds. Part 16. Unusual substituent effects on selenium-77 nuclear magnetic resonance chemical shifts of heteroaryl- and aryl-phosphine selenides. *X*-Ray crystal structure of tri(2-furyl)phosphine selenide, 2505–8
The [Fe$_3$(μ$_3$-CR)(CO)$_{10}$]$^-$ cluster anions as building blocks for the synthesis of mixed-metal clusters. Part 1. Synthesis of mixed clusters [MFe$_3$(μ$_3$-CMe)(CO)$_{10}$(PPh$_3$)] (M = Cu or Au) and crystal structure of [CuFe$_3$(μ$_3$-CMe)(CO)$_{10}$(PPh$_3$)], 2521–4

CRYSTAL STRUCTURE (contd)
Lewis-base adducts of Group 1B metal(I) compounds. Part 18. Stereochemistries and structures of the 1:1 neutral complexes of $Cu^{I}X$ with 1,10-phenanthroline (X = I) or 2,9-dimethyl-1,10-phenanthroline (X = I, Br, or Cl), 2531–40
Lewis-base adducts of Group 1B metal(I) compounds. Part 19. Crystal structures of bis(1,10-phenanthroline)copper(I) perchlorate and dibromocuprate(I), 2541–6
Interactions of $[Mo(CO)_6]$ and $[Mo(CO)_3(C_6H_5CH_3)]$ with O_2N_2- and O_2N_3-donor macrocycles and the *X*-ray crystal structure of tetracarbonyl(6,7,16,17-tetrahydro-15*H*-dibenzo[*e*,*n*][1,4,8,12]dioxadiazacyclopentadecine)molybdenum–toluene (2/1), 2561–4
Studies on spin-equilibrium iron(III) complexes. Part 1. Syntheses and magnetic properties of a new family of spin cross-over iron(III) complexes with a unidentate ligand over a wide range of the spectrochemical series and a quinquedentate ligand derived from salicylaldehyde and di(3-aminopropyl)amine. *X*-Ray crystal structure of [4-azaheptamethylene-1,7-bis(salicylideneiminato)](4-methylpyridine)iron(III) tetraphenylborate, 2575–84
Fragmentation of co-ordinated carbon disulphide and dithiocarbene ligands by nucleophilic attack at carbon: the crystal structure of $[(Ph_3P)IPt(\mu\text{-}SMe)(\mu\text{-}CSMe)PtI(PPh_3)]\cdot Me_2CO$, 2595–602
Crystal and molecular structure and magnetic properties of linear trimeric copper(II) complexes with predominant ferromagnetic exchange interaction, 2609–14
Indium derivatives of monothio-β-diketones and the *X*-ray structure of tris[benzoyl(thiobenzoyl)methanato-*O*,*S*]indium(III), 2623–8
Complexes of molybdenum-(II) and -(IV) and tungsten(II) with sterically hindered thiolate ligands. Synthesis, reactivity, and *X*-ray crystal structures of $[PPh_4][Mo(SC_6H_2Pr^{i}_3\text{-}2,4,6)_3(CO)_2]$ and $[Mo(NNPh)(SC_6H_2Pr^{i}_3\text{-}2,4,6)_3(NCMe)]$, 2639–46
Molybdenum, rhenium, and tungsten complexes with bi- and tridentate phosphinothiolato-ligands; structures of $[Mo\{PhP(CH_2CH_2S)_2\}_2]$ and $[Mo(NNMe_2)\{PhP(CH_2CH_2S)_2\}_2]$, 2647–54
A novel pentameric hydrolysis product of $SnMe_2Cl_2$: crystal and molecular structure of $[NHEt_3][(SnMe_2Cl)_5O_3]$, 2683–6

CYANIDE
Bimetallic systems. Part 7. Platinum and palladium dicyanides containing terminal or bridging $Ph_2PCH_2PPh_2$ and heterobimetallics with silver, gold, mercury, rhodium, iridium, or molybdenum, 279–84
Spectrochemistry of solutions. Part 16. A Raman spectroscopic study of the complexation of mercury(II) by cyanide ligands in liquid ammonia at 293 K, 663–8
Studies on transition-metal cyano-complexes. Part 4. Cyanide hydride complexes of Groups 6A and 8, 717–22
Lewis-base adducts of group 1B metal(I) compounds. Part 17. Synthesis and crystal structures of adducts of copper(I) cyanide with nitrogen bases, 839–44
On the effect of cyanide ion on the reaction of pentacyanonitrosylferrate(2−) with cysteine, 1191–4
Characterization of the adducts formed by Cu(CN) and Cu(NCS) with biquinoline. The crystal structure of the polymeric cyano-compound containing both linear and tetrahedrally co-ordinated copper(I), $[\{Cu_3(bq)_2(CN)_3\}_n]$, 1285–8
Bimetallic cyano-bridged cations: preparation and hydride reduction of $[(\eta^5\text{-}C_5H_5)L_2Ru(\mu\text{-}CN)ML'_2(\eta^5\text{-}C_5H_5)]PF_6$ [L_2, L'_2 = $(PPh_3)_2$, $Ph_2PCH_2CH_2PPh_2$; M = Ru or Fe]. Formation of $[Ru(\eta^5\text{-}C_5H_5)(PPh_3)H_3]$ and *X*-ray crystal structure of $[(\eta^5\text{-}C_5H_5)(Ph_2PCH_2CH_2PPh_2)Ru(\mu\text{-}CN)Ru(PPh_3)_2(\eta^5\text{-}C_5H_5)]PF_6$, 1479–86
Addition of proton and boron trifluoride at terminal and bridging cyano-ligands in dinuclear manganese compounds, 1609–12
The kinetics and stoicheiometry of silver(III) reduction by the octacyano-complexes of molybdenum(IV) and tungsten(IV), 1789–94
Kinetics and mechanism of the equilibration reaction between (2,2′,2″-nitrilotriethoxy)nitrosylvanadate(1−) and cyanide. Crystal structures of sodium (2,2′,2″-nitrilotriethoxy)nitrosylvanadate(I)–sodium perchlorate tetrahydrate and of barium cyano(2,2′,2″-nitrilotriethoxy)nitrosylvanadate(I) pentahydrate, 2493–8

CYANOETHYL
The co-ordination chemistry of manganese. Part 14. Synthesis of manganese(II) complexes of tertiary phosphine ligands containing 2-cyanoethyl groups, 135–40

CYCLOBUTANE
Reduction–oxidation properties of organotransition-metal complexes. Part 22. Stereospecific oxidative cyclopropane ring opening and reductive cyclobutane ring formation in polycyclic hydrocarbon complexes of iron; *X*-ray crystal structures of $[Fe_2(CO)_6(\eta^5:\eta'^5\text{-}C_{16}H_{16})][PF_6]_2\cdot CH_3NO_2$ and $[Fe_2(CO)_6(\eta^4:\eta'^4\text{-}C_{16}H_{16})]$, 1027–36

CYCLOHEPTATRIENYL
Mono-η-cycloheptatrienyltitanium chemistry: synthesis, molecular and electronic structures, and reactivity of the complexes $[Ti(\eta\text{-}C_7H_7)L_2X]$ (L = tertiary phosphine, O- or N-donor ligand; X = Cl or alkyl), 669–84

CYCLOHEXADIENYL
Reduction–oxidation properties of organotransition-metal complexes. Part 21. Synthesis and *X*-ray structural characterisation of the redox-related pair of cyclohexadienyl complexes $[Mn(CO)(dppe)(\eta^5\text{-}C_6H_6Ph)]$ and $[Mn(CO)(dppe)(\eta^5\text{-}C_6H_6Ph)][PF_6]\cdot 0.5CH_2Cl_2$, 1019–26
Mechanism of addition of aryltrimethyl-silanes and -stannanes to tricarbonyl(cyclohexadienyl)ruthenium(II) cation, 1049–52
Studies on the regioselective preparation of tricarbonyl(cyclohexadienyl)iron(1+) salts, 1723–6

CYCLOHEXENYL
A theoretical study of three-centre M–H–C interactions, 1637–44

CYCLOHEXYL
Organotin biocides. Part 2. Variable-temperature ^{119}Sn Mössbauer study of phenyl-and cyclohexyl-tin compounds, 1417–24

CYCLO-OCTADIENE
Chemistry of platinum sulphido-complexes. Part 5. Synthesis and crystal and molecular structure of 3-(η-cyclo-octa-1,5-diene)bis(μ$_3$-sulphido)-1,1,2,2-tetrakis(triphenylphosphine)diplatinum(II)rhodium(I) hexafluorophosphate–dichloromethane (1/1), $[Pt_2Rh(\mu_3\text{-}S)_2(PPh_3)_4(\eta\text{-}C_8H_{12})]PF_6\cdot CH_2Cl_2$, 851–6
Homo- and hetero-dinuclear hydride-bridged complexes containing cyclo-octadiene: the crystal and molecular structure of $[(\eta^4\text{-}C_8H_{12})Ir(\mu\text{-}H)(\mu\text{-}Cl)IrH_2(PPh_3)_2]$, 1277–80
Synthesis and structural studies of some 1,1′-dichloroferrocene derivatives of platinum(II). Crystal and molecular structure of 2,2-μ-[(1—2,5—6-η-*cis*,*cis*-cyclo-octa-1,5-diene)platinio]-bis(1,1′-dichloroferrocene), 1527–30
Oxygenation studies. Part 7. Catalytic dioxygenation of cyclo-octa-1,5-diene at a rhodium centre, 1591–6

CYCLO-OCTATETRAENE
Synthesis of $[Fe(\eta^4\text{-}C_8H_8)(CO)_2L]$ [L = MeNC, $Pr^{i}NC$, $Bu^{t}NC$, PhNC, 2,6-$Me_2C_6H_3NC$, or $P(OMe)_3$] and proof of the applicability of the Woodward–Hoffmann rules to the fluxionality of $[Fe(\eta^4\text{-}C_8H_8)(CO)_2(CNPr^{i})]$, 693–8

CYCLOPENTADIENYL
Reactions of co-ordinated ligands. Part 33. Mononuclear η^2-vinyl complexes: synthesis, structure, and reactivity, 435–50
Formation of substituted cyclopentadienyl ligands on tungsten *via* reactions between the alkyne complexes $[W(CO)(R^1C_2R^2)_3]$ (R^1 = R^2 = Ph or Et; R^1 = Me, R^2 = Ph) and the alkylidyne compounds $[W(\equiv CR)(CO)_2(\eta\text{-}C_5H_5)]$ (R = C_6H_4Me-4 or Me); *X*-ray crystal structures of $[W_2(\mu\text{-}CO)_2(CO)(\eta\text{-}PhC_2Ph)(\eta\text{-}C_5Ph_4R)(\eta\text{-}C_5H_5)]$ and $[W_2(\mu\text{-}EtC_2Et)(CO)_4(\eta\text{-}C_5Et_4R)(\eta\text{-}C_5H_5)]$ (R = C_6H_4Me-4), 905–12
Isolated CH stretching frequencies, methyl group geometry, and methyl CH bond lengths and strengths in tricarbonyl(η^5-cyclopentadienyl)methylchromium(II), -molybdenum(II), and -tungsten(II), 1207–12
Chemistry of di- and tri-metal complexes with bridging carbene or carbyne ligands. Part 35. The synthesis and crystal structures of the compounds $[N(PPh_3)_2][W_2\{\mu\text{-}C(H)C_6H_4Me\text{-}4\}(CO)_7(\eta\text{-}C_5H_5)]\cdot 0.5Et_2O$ and $[W\{\equiv C(H)C_6H_4Me\text{-}4\}\text{-}(SnPh_3)(CO)_2(\eta\text{-}C_5H_5)]$, 1331–8
Photochemistry of dicarbonyl(η^5-cyclopentadienyl)-methyl- and -ethyl-iron and -ruthenium complexes in solutions at −30 °C and in frozen gas matrices at 12 K, 1365–74
Reactions of bis(cyclopentadienyl)vanadium derivatives with nitrogen mono-oxide and the crystal structure of an oxo-bridged nitrosyl complex of vanadium, 1435–42
Geometry-dependent carbon-13 chemical shifts in (η^6-[2.2]cyclophane)(η^5-cyclopentadienyl)iron(II) hexafluorophosphates, 1661–4
Chemistry of di- and tri-metal complexes with bridging carbene or carbyne ligands. Part 37. Methylene group transfer to carbon–metal multiple bonds; crystal structures of $[TiW\{\mu\text{-}C(C_6H_4Me\text{-}4)=CH_2\}(\mu\text{-}CO)(CO)(\eta\text{-}C_5H_5)_2]$ and $[PtW\{\mu\text{-}C(C_6H_4Me\text{-}4)=CH_2\}(CO)_2(PMe_3)_2(\eta\text{-}C_5H_5)]$, 2009–16

CYCLOPENTADIENYLIDENEALKYL
Studies on cyclic bis(η^5:σ-2-cyclopentadienylidene-ethyl)- and bis(η^5:σ-4-cyclopentadienylidenebutyl)-molybdenum compounds, 1585–90
CYCLOPHANE
Geometry-dependent carbon-13 chemical shifts in (η^6-[2.2]cyclophane)(η^5-cyclopentadienyl)iron(II) hexafluorophosphates, 1661–4
CYCLOPROPANE
Reduction–oxidation properties of organotransition-metal complexes. Part 20. Oxidative and thermolytic cyclopropane ring-opening reactions; *X*-ray crystal structure of $[Fe_2(CO)_6(\eta^4:\eta^{-4}\text{-}C_{16}H_{18})]$, 699–706
Reduction–oxidation properties of organotransition-metal complexes. Part 22. Stereospecific oxidative cyclopropane ring opening and reductive cyclobutane ring formation in polycyclic hydrocarbon complexes of iron; *X*-ray crystal structures of $[Fe_2(CO)_6(\eta^5\text{:}\ \text{-}C_{16}H_{16})][PF_6]_2\cdot CH_3NO_2$ and $[Fe_2(CO)_6(\eta^4\text{:}\eta^{-4}\text{-}C_{16}H_{16})]$, 1027–36
CYCLOPROPENE
Reactions of co-ordinated ligands. Part 35. Evidence for carbon–carbon double-bond cleavage of cyclopropenes in their reaction with dinuclear cobalt, rhodium, and iridium complexes; crystal structure and protolysis of $[Rh_2(\mu\text{-}CO)(\mu\text{-}COCHCMe_2CH)(\mu\text{-}C_5Me_5)_2]$, 2483–92
CYCLOPROPENIUM
Annelation of ring-opened arylcyclopropenium ions to co-ordinated cyclo-octatetraene, and the *X*-ray crystal structure of $[Fe(CO)_3(\sigma,\eta^3\text{-}C_{11}H_9Ph_3)]$, 777–82
CYCLOPROPYLMETHYL
Reaction of transition-metal carbonylate anions and 1,1,1-tris(halogenomethyl)ethane. *X*-Ray crystal structures of tricarbonyl(η^5-cyclopentadienyl)(1-methylcyclopropylmethyl)tungsten(II), and tetraethylammonium enneacarbonyliododirhenate(0), 931–40
CYCLOTRIAZATHIENE
Reaction of trithiazyl trichloride, $(NSCl)_3$, with triphenylphosphine or triphenylphosphine metal complexes. *X*-Ray crystal structure of aminotriphenylphosphonium chloride–dichloromethane (1/1), $[Ph_3PNH_2]Cl\cdot CH_2Cl_2$, 1043–8
Thiazyl chloride complexes of ruthenium(II), 2449–52
CYCLOTRIPHOSPHAZATRIENE
Mass spectrometric studies on cyclo- and poly-phosphazenes. Part 2. Oligomerization of hexa(aryloxy)cyclotriphosphazatrienes, 1547–54
CYCLOTRIPHOSPHAZENE
Studies of phosphazenes. Part 21. Associative and dissociative pathways in the aminolysis reactions of halogenocyclotriphosphazenes, 285–90
Studies of phosphazenes. Part 22. High-field nuclear magnetic resonance investigation of novel isomeric oxophosphazadienes, 1431–4
Studies of phosphazenes. Part 25. Synthesis, nuclear magnetic resonance spectroscopy, and mode of formation of (aziridino)(triphenylphosphazenyl)cyclotriphosphazenes. *X*-ray crystal structure and enzyme-inhibiting activity of $N_3P_3(NPPh_3)(NC_2H_4)_5$, 1881–90
Studies of phosphazenes. Part 21. Associative and dissociative pathways in the aminolysis reactions of halogenocyclotriphosphazenes (1985, 285), 2459–60
CYMENE
Mononuclear η^6-*p*-cymeneosmium(II) complexes and their reactions with Al_2Me_6 and other methylating reagents, 573–8
CYSTEINE
On the effect of cyanide ion on the reaction of pentacyanonitrosylferrate(2−) with cysteine, 1191–4
A kinetic study of the complexation of cysteine and related compounds with aqueous vanadium(II) and vanadium(III) at approximately neutral pH; the mediating role of sulphur compounds in electron transfer, 2461–8
CYSTINE
On the effect of cyanide ion on the reaction of pentacyanonitrosylferrate(2−) with cysteine, 1191–4
DANSYLGLYCINATE
The effect of a dansyl group on the co-ordinative ability of *N*-protected amino acids. Part 1. Behaviour of the copper(II) ion–*N*-dansylglycinate system in aqueous and methanolic solution, 2363–8
DEALKYLATION
Kinetic studies of oxidative dealkylation of alkylcobalamins by hexachloroplatinate(IV), 1375–80
DECABORANE
Reactions of 6,6′-bis(*nido*-decaboranyl) oxide and 6-hydroxy-*nido*-decaborane with dihalogenobis(phosphine) complexes of nickel, palladium, and platinum, and some related chemistry; nuclear magnetic resonance investigations and the crystal and molecular structures of bis(dimethylphosphine)-di-μ-(2,3,4-η^3-*nido*-hexaboranyl)-diplatinum(*Pt–Pt*), $[Pt_2(\mu\text{-}\eta^3\text{-}B_6H_9)_2(PMe_2Ph)_2]$, and of 2,4-dichloro-1,1-bis(dimethylphenylphosphine)-*closo*-1-nickeladecaborane, $[(PhMe_2P)_2NiB_9H_7Cl_2]$, 953–72
Gold(I) complexes of 1,2-ditertiary phosphino-*o*-carborane ligands: two-, three-, and four-co-ordination, 1387–90
DECARBOXYLATION
Acidic and basic properties of the hydroxycarbonyl complex $[IrCl_2(CO_2H)(CO)(PMe_2Ph)_2]$, 857–60
DECOMPOSITION
Kinetic and structural investigations of $[Fe^{III}(edta)]^-$ [edta = ethylenediaminetetra-acetate(4−)] catalysed decomposition of hydrogen peroxide, 493–502
Some alkyldiplatinum(II) complexes and studies of the photochemical and thermal decomposition of the $[Pt_2Et(\mu\text{-}Ph_2PCH_2PPh_2)_2(Ph_2PCH_2PPh_2\text{-}P)]^+$ cation, 2421–6
Studies on singlet oxygen in aqueous solution. Part 4. The 'spontaneous' and catalysed decomposition of hydrogen peroxide, 2525–30
DEUTERIATION
Ligand reactivity in polypyridine complexes; the deuteriation of the bis(2,2′:6′,2″-terpyridine)ruthenium(II) cation, 2687–90
DIACETYLPYRIDINE
The template synthesis and crystal and molecular structure of a seven-co-ordinate manganese(II) complex with 2,6-diacetylpyridine mono(2-aminobenzoylhydrazone), 215–8
DIALKYL DISELENIDE
Pyramidal inversions and 1,2-metal shifts in pentacarbonyl-chromium, -molybdenum, and -tungsten derivatives of dialkyl disulphides and dialkyl diselenides. A nuclear magnetic resonance investigation, 1561–8
DIALKYL DISULPHIDE
Pyramidal inversions and 1,2-metal shifts in pentacarbonyl-chromium, -molybdenum, and -tungsten derivatives of dialkyl disulphides and dialkyl diselenides. A nuclear magnetic resonance investigation, 1561–8
DIAMINE
Synthesis of α-lithioarylmethanes of *m*-xylene and its α-trimethylsilyl derivatives; crystal structure of $[\{Li(Me_2NCH_2CH_2NMe_2)\}_2\{C_6H_4(CHSiMe_3)_2\text{-}m\}]$, 337–44
Reaction of some aliphatic diamines with four-co-ordinated unsymmetrical ketoenamine copper(II)and nickel(II) complexes, 803–6
Palladium(II,IV) mixed-valence complexes of 1,2-diaminoethane, 1,3-diaminopropane, and diethylenetriamine: syntheses, electronic, infrared, Raman, and resonance Raman spectra and *X*-ray studies, 815–20
DIARSENE
Purification and *X*-ray crystal structure of bis[tris(trimethylsilyl)methyl]diarsene, 383–6
DIASTEREOISOMERISM
Diastereoisomeric organophosphorus compounds. Part 4. Proton and phosphorus-31 nuclear magnetic resonance spectra of compounds of the type $[RR'P(X)]_2Y$ and $RR'P(X)\text{-}Y\text{-}(Z)PRR'$ (R = CH_3, R′ = t-C_4H_9) (1984, 2803), 871–2
DIAZATETRAPHOSPHACYCLO-OCTADECANE
Macrocyclic polyphosphane ligands. Iron(II), cobalt(II), and nickel(II) complexes of (4*RS*,7*RS*,13*SR*,16*SR*)-tetraphenyl-1,10-dipropyl-1,10-diaza-4,7,13,16-tetraphosphacyclo-octadecane: crystal structures of their tetraphenylborate derivatives, 479–86
DIAZENIDE
Complexes of molybdenum-(II) and -(IV) and tungsten(II) with sterically hindered thiolate ligands. Synthesis, reactivity, and *X*-ray crystal structures of $[PPh_4][Mo(SC_6H_2Pr^i_3\text{-}2,4,6)_3(CO)_2]$ and $[Mo(NNPh)(SC_6H_2Pr^i_3\text{-}2,4,6)_3(NCMe)]$, 2639–46
DIAZONIUM
The mechanisms of the reactions between hydrido-complexes and diazonium salts. Part 1. The reactions of *trans*-$[PtH(Cl)(PEt_3)_2]$, 2059–66
The mechanisms of the reactions between hydrido-complexes and diazonium salts. Part 2. The reactions of *cis,mer*-

DIAZONIUM (contd)
[$RhHCl_2(PEtPh_2)_3$], 2067–78
DIBENZODIOXADIAZACYCLOPENTADECINE
Interactions of [$Mo(CO)_6$] and [$Mo(CO)_3(C_6H_5CH_3)$] with O_2N_2- and O_2N_3-donor macrocycles and the *X*-ray crystal structure of tetracarbonyl(6,7,16,17-tetrahydro-15*H*-dibenzo[*e*,*n*][1,4,8,12]dioxadiazacyclopentadecine)molybdenum–toluene (2/1), 2561–4
DIBORANE
Contributions to the chemistry of boron. Part 160. A convenient synthesis of catecholatoborane and diborane, 1689–92
DICARBADODECABORANE
Metallaborane chemistry. Part 14.[1] Icosahedral η^6-arene carbametallaboranes of iron and ruthenium; molecular structures of *closo*-[1-(η^6-C_6H_5Me)-2,4-Me_2-1,2,4-$FeC_2B_9H_9$] and *closo*-[3-(η^6-C_6H_6)-3,1,2-$RuC_2B_9H_{11}$], 2343–8
DICARBAPALLADADODECABORANE
Carbaborane derivatives of the late- and post-transition elements. Part 3. Structural consequences of ligand substitution in palladadicarbadodecaboranes 3-L_2-3,1,2-$PdC_2B_9H_{11}$. The crystal and molecular structures of 3-[$Me_2N(CH_2)_2NMe_2$]-3,1,2-$PdC_2B_9H_{11}$ and 3-$(PMe_3)_2$-3,1,2-$PdC_2B_9H_{11}$, 761–70
DIELECTRIC
Beryllocene: A microwave dielectric loss study, 1761–6
DIENE
Organoruthenium(II) complexes formed by insertion reactions of some vinyl compounds and conjugated dienes into a hydrido–ruthenium bond, 873–8
Reactions of co-ordinated ligands. Part 34. Synthesis, structure, and reactivity of cationic dieneruthenium complexes; crystal structures of [$Ru(\eta^4$-$C_6H_8)(CO)(\eta$-$C_5H_5)$][BF_4] and [$Ru(\eta^3$-$C_6H_9)(CO)(\eta$-$C_5H_5)$], 2145–54
Co-ordinatively unsaturated diene complexes of tungsten(II) and their reactions with nucleophiles to give six- and seven-co-ordinate derivatives, 2231–8
DIETHYLAMINOETHANOLATE
Magnetic behaviour of tetrakis[(2-diethylaminoethanolato)isocyanatocopper(II)], a complex with an antiferromagnetic ground state; the crystal and molecular structure of the triclinic modification, 1909–14
DIETHYLDITHIOCARBAMATE
Zinc–sulphur bond enthalpy: its determination in bis(diethyldithiocarbamato)zinc(II), 369–72
Electrostatic solvent effect on the formation of the mixed-chelate complex (acetylacetonato)(diethyldithiocarbamato)-copper(II), 987–90
High-co-ordination-number compounds of niobium and tantalum: Reactions of niobium and tantalum halides and sulphido-halides with sodium diethyldithiocarbamate. The crystal structures of [$Nb(S_2CNEt_2)_4$]Br, $Nb(S_2CNEt_2)_3S$, and $Ta(S_2CNEt_2)_3(S_2)$, 1821–8
DIFLUOROPHOSPHATE
Difluorophosphate complexes of chromium, manganese, iron, cobalt, and nickel, 707–10
Crystal structures of the difluorophosphate complexes, $Co(O_2PF_2)_2$·2MeCN and $Cu(O_2PF_2)_2$, 2433–6
DIFLUOROPHOSPHINE SULPHIDE
Determination of the molecular structure of difluorophosphine sulphide by the combined analysis of data from electron diffraction, microwave spectroscopy, and liquid crystal nuclear magnetic resonance spectroscopy, 755–60
DIHYDROGEN
Hydrogen generation by hydrolysis of sodium tetrahydroborate: effects of acids and transition metals and their salts, 307–14
Heteropolytungstates as catalysts for the photochemical reduction of oxygen and water, 395–400
DI-IMINE
Application of photoelectron spectroscopy to molecular properties. Part 19. Electronic structure of tris(α-di-imino) complexes of ruthenium(0), 43–50
DIKETONATE
Indium derivatives of monothio-β-diketones and the *X*-ray structure of tris[benzoyl(thiobenzoyl)methanato-*O*,*S*]indium(III), 2623–8
DIMETHYLAMIDOGALLANE
The molecular structure of gaseous dimethylamidogallane: characterization of the dimer [$Me_2NGaH_2]_2$ by electron diffraction and vibrational spectroscopy, 807–14
DIMETHYLAMINOPROPANOLATE
Crystal and molecular structures and magnetic properties of bromo(3-dimethylaminopropan-1-olato)copper(II), chloro(3-dimethylaminopropan-1-olato)copper(II), and (3-diethylaminopropan-1-olato)isocyanatocopper(II), 913–20
DIMETHYLFORMAMIDE
Ligand substitution on (*N*,*N*-dimethylformamide)[2,2′,2″-tri(*N*,*N*-dimethylamino)triethylamine]cobalt(II), 413–6
DIMETHYLGLYOXIMATE
Routes of formation and crystal structure of an alkylperoxycobaloxime: bis[dimethylglyoximato(1−)](4-ethoxycarbonylbut-3-en-2-ylperoxo)(pyridine)cobalt(III), 1997–2000
DIMETHYLPHOSPHINOMETHYLENE
Trimethylphosphine as a reactive solvent: synthesis, crystal structures, and reactions of [$Ta(PMe_3)_3(\eta^2$-$CH_2PMe_2)(\eta^2$-$CHPMe_2)$] and [$W(PMe_3)_4(\eta^2$-$CH_2PMe_2)H$] and related studies, 2025–36
DIMETHYLPYRAZOLATE
Pyrazolate A-frame rhodium complexes. Crystal structures of [$Rh_2(\mu$-dmpz$)(CO)_2(\mu$-dppm$)_2$][ClO_4] and [$Rh_2(\mu$-dmpz$)I_2(CO)_2(\mu$-dppm$)_2$][ClO_4], 973–80
DIMETHYLSILYL
Determination of the molecular structures and conformations of methylbis(methylsilyl)amine and bis(dimethylsilyl)methylamine in the gas phase by electron diffraction, 191–8
DIMETHYL SULPHOXIDE
Reactions in mixed non-aqueous systems containing sulphur dioxide. Part 6. The reaction of metal oxides with dimethyl sulphoxide–sulphur dioxide, 99–100
Equilibrium and kinetic study of pyridine binding to phthalocyaninatoiron(II) in dimethyl sulphoxide, 1107–12
Equilibrium and kinetic study of the reaction between phthalocyaninatoiron(II) and carbon monoxide in dimethyl sulphoxide in the presence of pyridine. Evidence for the formation of a transient, 1113–8
DINITROGEN
The preparation and properties of some diphosphines $R_2PCH_2CH_2PR_2$ (R = alkyl or aryl) and of their rhenium(I) dinitrogen derivatives, 1131–6
Electron-transfer reactions in nitrogen fixation. Part 1. The electrosynthesis of dinitrogen, hydride, isocyanide, and carbonyl complexes of molybdenum: intermediates, mechanisms, and energetics, 1255–64
Preparation and properties of *cis*-[$W(N_2)_2(Ph_2PCH_2PPh_2)_2$] and *trans*-[$M(N_2)_2(Ph_2PCH{=}CHPPh_2)_2$] (M = Mo or W), and the crystal structure of *cis*-[$W(N_2)_2(PMe_2Ph)_4$], 1523–6
Preparation and properties of *mer*-[$ReCl(N_2)(CNR)\{P(OMe)_3\}_3$] (R = Me, Et, Bu^t, C_6H_4Me-4, or C_6H_4Cl-4) and [$ReCl(N_2)(CNMe)(PPh_3)\{P(OEt)_3\}_2$]. *X*-Ray crystal structure of *mer*-[$ReCl(N_2)(CNMe)\{P(OMe)_3\}_3$] and reductive cleavage of the isocyanide ligands to primary amines upon protonation, 2079–84
Nitrogen-15 nuclear magnetic resonance relaxation mechanisms in dinitrogen complexes of molybdenum, tungsten, rhenium, and osmium, 2473–8
DINITROGEN TETRAOXIDE
Electrochemical studies in HNO_3–N_2O_4 mixtures: corrosion of stainless steel in HNO_3–N_2O_4 mixtures and the effect of inhibitors, 2551–4
DIOXANE
Influence of decreasing solvent polarity (dioxane–water mixtures) on the stability and structure of binary and ternary complexes of adenosine 5′-triphosphate and uridine 5′-triphosphate, 2291–304
DIOXYGEN
Rate parameters for oxygen and carbon monoxide binding to a liposome-embedded heme under physiological conditions, 65–8
Phosphocholine-substituted 5,10,15,20-tetraphenylporphyrinatoiron(II): oxygen carrier under physiological conditions, 275–8
Complexes of the platinum metals. Part 24. The role of dioxygen in the reactions of trifluoroacetic acid with the rhodium and iridium nitrosyls [$M(NO)(PPh_3)_3$], 617–20
Studies on singlet oxygen in aqueous solution. Part 1. Formation of singlet oxygen from hydrogen peroxide with two-electron oxidants, 1141–6
Studies on singlet oxygen in aqueous solution. Part 2. Water-soluble square-planar nickel complexes as quenchers, 1147–50
Studies on singlet oxygen in aqueous solution. Part 3. The decomposition of peroxy-acids, 1151–4
Oxygenation studies. Part 7. Catalytic dioxygenation of cyclo-octa-

DIOXYGEN (contd)
1,5-diene at a rhodium centre, 1591–6
Studies on singlet oxygen in aqueous solution. Part 4. The 'spontaneous' and catalysed decomposition of hydrogen peroxide, 2525–30
DIPHOSPHANE
Diastereoisomeric organophosphorus compounds. Part 4. Proton and phosphorus-31 nuclear magnetic resonance spectra of compounds of the type $[RR'P(X)]_2Y$ and RR'P(X)-Y-(Z)PRR' (R = CH_3, R' = t-C_4H_9) (1984, 2803), 871–2
DIPHOSPHINE
Bimetallic systems. Part 7. Platinum and palladium dicyanides containing terminal or bridging $Ph_2PCH_2PPh_2$ and heterobimetallics with silver, gold, mercury, rhodium, iridium, or molybdenum, 279–84
The preparation and hydridic reduction of dicationic dicarbonyl complexes of osmium(II); the crystal and molecular structure of *trans*-bis[1,2-bis(diphenylphosphino)ethane-*PP'*]carbonylformylosmium(II) hexafluoroantimonate–dichloromethane (1/1), 387–94
Bimetallic systems. Part 8. Heterobimetallic di-isonitrile or isonitrile–carbonyl complexes of rhodium or iridium bridged to silver or gold by $Ph_2PCH_2PPh_2$, 511–6
Synthesis and characterisation of 1,2-bis(dimethylphosphino)ethane (dmpe) complexes of chromium-(0) and -(IV): *X*-ray crystal structures of *trans*-$Cr(N_2)_2(dmpe)_2$, *cis*-$Cr(CO)_2(dmpe)_2$, $Cr(C_2Ph_2)_2(dmpe)$, and $CrH_4(dmpe)_2$, 685–92
The addition of protic acids to $[Mn_2(CO)_5(Ph_2PCH_2PPh_2)_2]$ to give bridging hydrido-compounds, 743–8
Interconversion of 42- and 44-electron platinum *triangulo*-clusters using chelating tertiary phosphine ligands and the structural characterisation of [1,3-bis(diphenylphosphino)propane]-tris(μ-sulphur dioxide)bis(tricyclohexylphosphine)triplatinum–benzene (1/2), $[Pt_3(\mu\text{-}SO_2)_3\{P(C_6H_{11})_3\}_2(dppp)]\cdot 2C_6H_6$, 845–50
Alkyl, hydrido, and tetrahydroaluminato complexes of manganese with 1,2-bis(dimethylphosphino)ethane (dmpe). *X*-Ray crystal structures of $Mn_2(\mu\text{-}C_6H_{11})_2(C_6H_{11})_2(\mu\text{-}dmpe)$, $(dmpe)_2Mn(\mu\text{-}H)_2AlH(\mu\text{-}H)_2AlH(\mu\text{-}H)_2Mn(dmpe)_2$, and $Li_4\{MnH(C_2H_4)[CH_3(Me)PCH_2CH_2PMe_2]_2\}_2\cdot 2Et_2O$, 921–30
Pyrazolate A-frame rhodium complexes. Crystal structures of $[Rh_2(\mu\text{-}dmpz)(CO)_2(\mu\text{-}dppm)_2][ClO_4]$ and $[Rh_2(\mu\text{-}dmpz)I_2(CO)_2(\mu\text{-}dppm)_2][ClO_4]$, 973–80
Bimetallic systems. Part 9. The synthesis of and nuclear magnetic resonance studies on 10-membered ring complexes of type $[(OC)_4M^1(\mu\text{-}Ph_2PCH_2CH_2PPh_2)_2M^2(CO)_4]$ (M^1, M^2 = Cr, Mo, or W), 1009–14
Bimetallic systems. Part 10. Synthesis of complexes of type $[(RC{\equiv}C)Pt(\mu\text{-}dppm)_2Pt(C{\equiv}CR)]$ (dppm = $Ph_2PCH_2PPh_2$, R = Ph or *p*-tolyl) and their corresponding 'A frames' $[(RC{\equiv}C)Pt(\mu\text{-}dppm)_2(\mu\text{-}H)Pt(C{\equiv}CR)]Cl$ or $[(RC{\equiv}C)Pt(\mu\text{-}dppm)_2(\mu\text{-}X)Pt(C{\equiv}R)]$ with X = CS_2 or MeOOCC≡CCOOMe, 1015–8
Reduction–oxidation properties of organotransition-metal complexes. Part 21. Synthesis and *X*-ray structural characterisation of the redox-related pair of cyclohexadienyl complexes $[Mn(CO)(dppe)(\eta^5\text{-}C_6H_6Ph)]$ and $[Mn(CO)(dppe)(\eta^5\text{-}C_6H_6Ph)][PF_6]\cdot 0.5CH_2Cl_2$, 1019–26
The preparation and properties of some diphosphines $R_2PCH_2CH_2PR_2$ (R = alkyl or aryl) and of their rhenium(I) dinitrogen derivatives, 1131–6
A mechanistic study on complexes of type *mer*-$[Cr(CO)_3(\eta^2\text{-}L\text{–}L)(\sigma\text{-}L\text{–}L)]$ (where L–L = $Ph_2PCH_2PPh_2$, $Ph_2PNHPPh_2$, or $Ph_2PNMePPh_2$) using spectroscopic and convolutive electrochemical techniques, 1213–22
Electron-transfer reactions in nitrogen fixation. Part 1. The electrosynthesis of dinitrogen, hydride, isocyanide, and carbonyl complexes of molybdenum: intermediates, mechanisms, and energetics, 1255–64
Synthesis and properties of the divalent 1,2-bis(dimethylphosphino)ethane (dmpe) complexes $MCl_2(dmpe)_2$ and $MMe_2(dmpe)_2$ (M = Ti, V, Cr, Mn, or Fe). *X*-Ray crystal structures of $MCl_2(dmpe)_2$ (M = Ti, V, or Cr), $MnBr_2(dmpe)_2$, $TiMe_{1.3}Cl_{0.7}(dmpe)_2$, and $CrMe_2(dmpe)_2$, 1339–48
Gold(I) complexes of 1,2-ditertiary phosphino-*o*-carborane ligands: two-, three-, and four-co-ordination, 1387–90
Bimetallic cyano-bridged cations: preparation and hydride reduction of $[(\eta^5\text{-}C_5H_5)L_2Ru(\mu\text{-}CN)ML'_2(\eta^5\text{-}C_5H_5)]PF_6$ [L_2, L'_2 = $(PPh_3)_2$, $Ph_2PCH_2CH_2PPh_2$; M = Ru or Fe]. Formation of $[Ru(\eta^5\text{-}C_5H_5)(PPh_3)H_3]$ and *X*-ray crystal structure of $[(\eta^5\text{-}C_5H_5)(Ph_2PCH_2CH_2PPh_2)Ru(\mu\text{-}CN)Ru(PPh_3)_2(\eta^5\text{-}C_5H_5)]PF_6$, 1479–86
Synthetic and nuclear magnetic resonance studies on dialkyl- and diaryl-platinum complexes containing chelating, monodentate, or bridging $Ph_2PCH_2PPh_2$ ligands, 1501–6
The syntheses and characterisation of, and the determination of $^1J(^{103}Rh\text{–}^{103}Rh)$ in $[(\eta^5\text{-}C_5Me_5)_2Rh_2(\mu\text{-}CH_2)_2\{\mu\text{-}CH_2CR(CH_2CR{=}CH_2)CH_2\}]$ (R = H or Me) and $[(\eta^5\text{-}C_5Me_5)_2Rh_2(\mu\text{-}CH_2)_2\{\mu\text{-}Ph_2P(CH_2)_nPPh_2\}]^{2+}$ (*n* = 1 or 2), 1555–60
Synthesis of bis[μ-bis(diphenylphosphino)methane]-tri-μ-carbonyl-tricarbonyl-*triangulo*-trirhodium(1+) perchlorate, $[Rh_3(CO)_3(\mu\text{-}CO)_3(\mu\text{-}dppm)_2]ClO_4$. An unusual 46-electron cluster possessing the A-frame structure with a bridging $Rh(CO)_3$ fragment, 1577–84
Addition of proton and boron trifluoride at terminal and bridging cyano-ligands in dinuclear manganese compounds, 1609–12
Bimetallic systems. Part 11. Heterobimetallic and unsymmetrical diplatinum complexes from *cis*-$[PtR_2(dppm\text{-}P)_2]$ (dppm = $Ph_2PCH_2PPh_2$; R = Me, 1-naphthyl, or C_6H_4Me-*o*): crystal structure of $[(C_6H_4Me\text{-}o)_2Pt(\mu\text{-}dppm)_2PtMe_2]$, 1677–82
Reaction between $[Ru_3(CO)_{12}]$ and the tridentate phosphine ligand $HC(PPh_2)_3$; *X*-ray crystal structures of complexes $[Ru_3(CO)_9\{Ph_2PCHP(Ph)C_6H_4PPh\}]$, $[Ru_2H(CO)_4(Ph_2PCHPPh_2)\{PhPC_6H_4C(O)\}]$, and $[Ru_2(CO)_4Cl(PPh_2)(dppm)]$, 1835–42
Bimetallic systems. Part 12. Mixed rhodium(I)–platinum(II) acetylide complexes containing bridging $Ph_2PCH_2PPh_2$. Crystal structures of $[(MeC{\equiv}C)Pt(\mu\text{-}dppm)_2(\sigma,\eta\text{-}C{\equiv}CMe)Rh(CO)]PF_6$ and of $[ClPt(\mu\text{-}dppm)_2(\sigma,\eta\text{-}C{\equiv}CMe)Rh(CO)]PF_6$, 2121–30
Bimetallic systems. Part 13. Platinum–manganese carbonyl complexes containing bridging $Ph_2PCH_2PPh_2$(dppm) ligands: crystal structure of $[(OC)_3Mn(\mu\text{-}dppm)_2PtH(Br)]BF_4$, 2131–8
Some alkyldiplatinum(II) complexes and studies of the photochemical and thermal decomposition of the $[Pt_2Et(\mu\text{-}Ph_2PCH_2PPh_2)_2(Ph_2PCH_2PPh_2\text{-}P)]^+$ cation, 2421–6
DIPHOSPHINE OXIDE
Investigation into diphosphine oxides as ligands in diorganotin(IV) adducts. Part 3. Synthesis and crystal structure of two adducts of dinitratodiphenyltin(IV) with *cis*- and *trans*-1,2-bis(diphenylphosphoryl)ethylene, 487–92
DIPHOSPHONATE
Synthesis, spectroscopy, and structure of the mixed-valence complexes $[Pt^{II}(en)_2][Pt^{IV}(en)_2X_2][Pt^{III}_2(H_2P_2O_5)_4X_2]$ (en = 1,2-diaminoethane, X = Br or I), 579–86
DISELENACYCLOPENTANE
Dynamic nuclear magnetic resonance study of Group 6 metal pentacarbonyl complexes. Pyramidal inversion and 1,2-metal shifts in the complexes $[M(CO)_5(Me_2CCH_2EECH_2)]$ (M = Cr, Mo, or W; E = S or Se), 1569–76
DISELENOETHER
Synthesis, properties, and multinuclear magnetic resonance (^1H, ^{77}Se, and ^{195}Pt) studies on diselenoether complexes of palladium, platinum, and rhodium, 1265–70
DISMUTATION
Dismutation of superoxide ion in an aprotic solvent by 5,10,15,20-tetra-*p*-tolylporphyrinatocobalt(II), 1513–6
DI-T-BUTYLPHOSPHAZENE
A photoelectron spectroscopic study of di-t-butylphosphazene, 879–84
DITHIACYCLOPENTANE
Dynamic nuclear magnetic resonance study of Group 6 metal pentacarbonyl complexes. Pyramidal inversion and 1,2-metal shifts in the complexes $[M(CO)_5(Me_2CCH_2EECH_2)]$ (M = Cr, Mo, or W; E = S or Se), 1569–76
DITHIADIZAOLIUM
Preparation of 1,3,2-dithiazolium hexafluoroarsenate(V), preparation and crystal structures of 5-methyl-1,3,2,4-dithiadiazolium and 4-methyl-1,3,2-dithiazolium hexafluoroarsenate(V) and the reduction of these salts to stable free radicals, 1405–16
DITHIAZOLIUM
Preparation of 1,3,2-dithiazolium hexafluoroarsenate(V), preparation and crystal structures of 5-methyl-1,3,2,4-dithiadiazolium and 4-methyl-1,3,2-dithiazolium hexafluoroarsenate(V) and the reduction of these salts to stable free radicals, 1405–16

DITHIOCARBAMATE
Synthesis and characterization of several mononuclear cobalt(IV) dithiocarbamates, $[Co(S_2CNRR')_3]ClO_4$ [R = R' = Me, Et, or CH_2Ph; R = Me, R' = Ph; R,R' = $-(CH_2)_4-$], 597–600
The concentration and selective extraction of copper(II), rhodium(III), and iridium(III) using a copolymer functionalised with dithiocarbamate groups. Spectroscopic evidence for the nature of the binding sites, 1655–60
DITHIOCARBENE
Fragmentation of co-ordinated carbon disulphide and dithiocarbene ligands by nucleophilic attack at carbon: the crystal structure of $[(Ph_3P)IPt(\mu\text{-}SMe)(\mu\text{-}CSMe)PtI(PPh_3)]\cdot Me_2CO$, 2595–602
DITHIOLATE
Molecular structure and solid-state properties of the two-dimensional conducting mixed-valence complex $[NBu_4]_{0.29}[Ni(dmit)_2]$ and the neutral $[Ni(dmit)_2]$ (H_2dmit = 4,5-dimercapto-1,3-dithiole-2-thione); members of an electron-transfer series, 783–94
Copper(II) salts of metal dithiolates, 1731–2
DITHIOMALONAMIDE
N,N'-Disubstituted dithiomalonamide complexes of antimony(III). Crystal and molecular structure of $[SbCl_3\{C_2H_5NHC(S)CH_2C(S)NHC_2H_5\}]$ with lone-pair occupation of an antimony co-ordination site, 1073–6
ELECTROCHEMISTRY
Reactions of sulphoxide–thioether bidentate ligands with platinum(II). Determination of the mode of binding by cyclic voltammetry, 209–12
Influence of secondary ligands on the stability of metal–xanthosine complexes in solution, 239–42
Electrochemistry of clusters. Part 4. Redox behaviour of tetrametallic clusters $[M^1{}_2M^2{}_2(\eta^5\text{-}C_5H_5)_2(\mu_3\text{-}CO)_2(\mu\text{-}CO)_4(PR_3)_2]$ (M^1 = Pt or Pd; M^2 = Cr, Mo, or W; R = Me, Et, Bu^n, or Ph), 711–6
Electron-transfer reactions from *cis*-dialkylbis(2,2'-bipyridyl)cobalt(III) complexes to organic oxidants, 899–904
Electrochemically induced ligand substitutions on $[OsCl_3(PMe_2Ph)_3]$: rational pathways to osmium(II) complexes, 947–52
Reduction–oxidation properties of organotransition-metal complexes. Part 21. Synthesis and *X*-ray structural characterisation of the redox-related pair of cyclohexadienyl complexes $[Mn(CO)(dppe)(\eta^5\text{-}C_6H_6Ph)]$ and $[Mn(CO)(dppe)(\eta^5\text{-}C_6H_6Ph)][PF_6]\cdot 0.5CH_2Cl_2$, 1019–26
Reduction–oxidation properties of organotransition-metal complexes. Part 22. Stereospecific oxidative cyclopropane ring opening and reductive cyclobutane ring formation in polycyclic hydrocarbon complexes of iron; *X*-ray crystal structures of $[Fe_2(CO)_6(\eta^5\text{:}\eta^5\text{-}C_{16}H_{16})][PF_6]_2\cdot CH_3NO_2$ and $[Fe_2(CO)_6(\eta^4\text{:}\eta^4\text{-}C_{16}H_{16})]$, 1027–36
SUPERQUAD: an improved general program for computation of formation constants from potentiometric data, 1195–200
A mechanistic study on complexes of type *mer*-$[Cr(CO)_3(\eta^2\text{-}L\text{–}L)(\sigma\text{-}L\text{–}L)]$ (where L–L = $Ph_2PCH_2PPh_2$, $Ph_2PNHPPh_2$, or $Ph_2PNMePPh_2$) using spectroscopic and convolutive electrochemical techniques, 1213–22
Electrochemical and chemical properties of di-iodonitrosyl[tris(3,5-dimethylpyrazolyl)borato]molybdenum, $[Mo\{HB(Me_2pz)_3\}(NO)I_2]$, and related complexes, 1249–54
Potentiometric study of the complex-formation equilibria of manganese(II), cobalt(II), nickel(II), copper(II), and zinc(II) with ethylenediamine-*N*-acetic acid, 1605–8
Studies of 2,5;6,10;8,10-tri-μ-hydro-nonahydro-*nido*-nonaborate(1−), $[B_9H_{12}]^-$: preparation, crystal and molecular structure, nuclear magnetic resonance spectra, electrochemistry, and reactions, 1645–54
Axially asymmetric metal alkyls. Part 3. Chemical, electrochemical, and structural studies of group 5A $d^{0,1,2}$ metallepines $[M\{(2\text{-}CH_2C_6H_4)_2\}(\eta\text{-}C_5H_5)_2]^z$ (M = Nb or Ta; z = 1−, 0, or 1+); synthesis of $[Nb\{2\text{-}CH_2(C_6H_4)_2Me\text{-}2'\}X(\eta\text{-}C_5H_5)_2]$ (X = Cl or CO) and $[\{(\eta\text{-}C_5H_5)_2ClV\}_2\{(2\text{-}CH_2C_6H_4)_2\}]$, and electrochemistry of $[M(CH_2Ph)_{2-n}Cl_n(\eta\text{-}C_5H_5)_2]$ (n = 0, M = Nb, or Ta; n = 1, M = V), 1747–60
The hydrolysis of metal ions. Part 8. Aluminium(III), 1967–70
Crystal structure and spectroscopic and redox properties of the iron–sulphur cluster compound $[NEt_4]_2[Fe_4S_4(SC_6H_4NH_2\text{-}4)_4]$, 2161–6
Copper(II) complexes of *N,N'*-bis(2-carbamoylethyl)ethylenediamine in methanol–water, 2217–20
Linkage isomerism in penta-ammineruthenium(II),(III) complexes of benzotriazole, 2469–72
Selective conversion of CO into methanol at ordinary temperature. Part 4. Activation by iron(II), iron(III), and chromium(III) complexes, 2499–504
Electrochemical studies in HNO_3–N_2O_4 mixtures: corrosion of stainless steel in HNO_3–N_2O_4 mixtures and the effect of inhibitors, 2551–4
Equilibrium and structural studies of silicon(IV) and aluminium(III) in aqueous solution. Part 13. A potentiometric and ^{27}Al nuclear magnetic resonance study of speciation and equilibria in the aluminium(III)–oxalic acid–hydroxide system, 2665–70
ELECTRON DIFFRACTION
Determination of the molecular structures and conformations of methylbis(methylsilyl)amine and bis(dimethylsilyl)methylamine in the gas phase by electron diffraction, 191–8
Determination of the molecular structure of difluorophosphine sulphide by the combined analysis of data from electron diffraction, microwave spectroscopy, and liquid crystal nuclear magnetic resonance spectroscopy, 755–60
The molecular structure of gaseous dimethylamidogallane: characterization of the dimer $[Me_2NGaH_2]_2$ by electron diffraction and vibrational spectroscopy, 807–14
Determination of the molecular structures of tri(t-butyl)phosphine oxide and tri(t-butyl)phosphine imide in the gas phase by electron diffraction, 827–30
The molecular structure of gaseous bis(trifluoromethyl)selenium difluoride as determined by electron diffraction, 941–6
Molecular structure of gaseous copper(I) acetate as determined by electron diffraction, 2555–60
ELECTRONIC
Application of photoelectron spectroscopy to molecular properties. Part 19. Electronic structure of tris(α-di-imino) complexes of ruthenium(0), 43–50
Glycine complexation with uranyl ion: absorptiometric, luminescence, and *X*-ray structural studies of tetrakis-(glycine)dioxouranium(VI) nitrate, 517–22
Characterisation of chromium(VI) oxide tetrafluoride, $CrOF_4$, and caesium pentafluoro-oxochromate(VI) $Cs[CrOF_5]$, 529–34
Synthesis, spectroscopy, and structure of the mixed-valence complexes $[Pt^{II}(en)_2][Pt^{IV}(en)_2X_2][Pt^{III}{}_2(H_2P_2O_5)_4X_2]$ (en = 1,2-diaminoethane, X = Br or I), 579–86
Absorption and emission in tris(2,2'-bipyridyl)ruthenium(II); effects of excited-state asymmetry, 1081–6
Spectroscopic studies on the higher binary fluorides of chromium: CrF_4, CrF_5, and CrF_6, both in the solid state and isolated in inert gas matrices, 1443–50
ELECTRON TRANSFER
Photochemistry of manganese porphyrins. Part 9. Redox reactions photosensitised by diamagnetic metalloporphyrins, 503–10
Kinetics and mechanism of single electron oxidations of the tervalent uranium ion, U^{3+}(aq), by free radicals in aqueous solutions, 641–4
Electron-transfer reactions from *cis*-dialkylbis(2,2'-bipyridyl)cobalt(III) complexes to organic oxidants, 899–904
Crystal structures and magnetic properties of binuclear five-co-ordinate copper(II) complexes with a phenolate bridge and their catalytic functions in multielectron redox reactions, 1945–52
Preparation and characterisation of 2,2'-bipyridine-4,4'-disulphonic and -5-sulphonic acids and their ruthenium(II) complexes. Excited-state properties and excited-state electron-transfer reactions of ruthenium(II) complexes containing 2,2'-bipyridine-4,4'-disulphonic acid or 2,2'-bipyridine-4,4'-dicarboxylic acid, 2247–62
A kinetic study of the complexation of cysteine and related compounds with aqueous vanadium(II) and vanadium(III) at approximately neutral pH; the mediating role of sulphur compounds in electron transfer, 2461–8
Kinetics and mechanism of electron-transfer reactions of bismuth(V) in aqueous acidic perchlorate–fluoride media. Part 1. Oxidation of hypophosphorous acid, 2571–4
ELECTROSYNTHESIS
Electron-transfer reactions in nitrogen fixation. Part 1. The electrosynthesis of dinitrogen, hydride, isocyanide, and carbonyl complexes of molybdenum: intermediates, mechanisms, and energetics, 1255–64
ELIMINATION
The chemistry of vitamin B_{12}. Part 25. Mechanism of the β-elimination of olefins from alkylcorrinoids; evidence for an initial

ELIMINATION (contd)
homolytic fission of the Co–C bond, 1613–8
ENANTIOSELECTIVITY
Enantioselectivity of nickel(II) and copper(II) complexes of Schiff bases derived from amino acids and (*S*)-*o*-[(*N*-benzylprolyl)amino]acetophenone or (*S*)-*o*-[(*N*-benzylprolyl)amino]benzaldehyde. Crystal and molecular structures of [Ni{(*S*)-bap-(*S*)-Val}] and [Cu{(*S*)-bap-(*S*)-Val}], 17–26
ENZYME
Studies of phosphazenes. Part 25. Synthesis, nuclear magnetic resonance spectroscopy, and mode of formation of (aziridino)(triphenylphosphazenyl)cyclotriphosphazenes. *X*-ray crystal structure and enzyme-inhibiting activity of $N_3P_3(NPPh_3)(NC_2H_4)_5$, 1881–90
EQUILIBRIUM
Electrostatic solvent effect on the formation of the mixed-chelate complex (acetylacetonato)(diethyldithocarbamato)-copper(II), 987–90
Equilibrium and kinetic study of pyridine binding to phthalocyaninatoiron(II) in dimethyl sulphoxide, 1107–12
Equilibrium and kinetic study of the reaction between phthalocyaninatoiron(II) and carbon monoxide in dimethyl sulphoxide in the presence of pyridine. Evidence for the formation of a transient, 1113–8
Equilibria between mono- and bi-nuclear complexes in $Cu(O_2CMe)_2$–pyridine derivative–diluent systems. The influence of the amine ligand basicity, 1849–52
The preparation of trichloro(phenyl 2-pyridylmethyl sulphoxide)gold(III) and the kinetics and equilibria of its formation and subsequent reactions, 2091–4
Copper(II) complexes of *N*,*N*′-bis(2-carbamoylethyl)ethylenediamine in methanol–water, 2217–20
Influence of decreasing solvent polarity (dioxane–water mixtures) on the stability and structure of binary and ternary complexes of adenosine 5′-triphosphate and uridine 5′-triphosphate, 2291–304
Equilibrium and structural studies of silicon(IV) and aluminium(III) in aqueous solution. Part 13. A potentiometric and ^{27}Al nuclear magnetic resonance study of speciation and equilibria in the aluminium(III)–oxalic acid–hydroxide system, 2665–70
E.S.R.
Electron spin resonance spectra of manganese(II) ions in the double-stranded chain polymers $CdLX_2$ (L = pyridine or methylpyridine; X = Cl or Br), 75–80
Comparison between the bis(2-thiopyridine *N*-oxide) derivatives of Cu^{II} and Ni^{I}: an electron spin resonance study, 379–82
Tin–molybdenum oxides. A study by *X*-ray diffraction, Mössbauer spectroscopy, and electron spin resonance, 451–4
Synthesis and physico-chemical properties of cationic derivatives of phthalocyaninatocopper(II), 651–8
A new electron spin resonance spectrum for an exchange- and dipole–dipole coupled superoxocobalamin · · · free-radical pair occurring in adenosylcobalamin-containing systems, 891–4
Electron spin resonance studies of iron(III) complexes of ethylenediaminetetra-acetate and *N*-(2-hydroxyethyl)ethylenediamine-*NN*′*N*′-triacetate in co-ordinating solvents, 1077–80
Electron spin resonance parameters for some copper(II)–bis(amino-acid) complexes, 1717–8
Photochemical studies of the alkylammonium molybdates. Part 7. Octahedral sites for multi-electron reduction of $[Mo_8O_{26}(MoO_4)_2]^{8-}$, 2585–90
ETHANE
Reaction of transition-metal carbonylate anions and 1,1,1-tris(halogenomethyl)ethane. *X*-Ray crystal structures of tricarbonyl(η⁵-cyclopentadienyl)(1-methylcyclopropylmethyl)tungsten(II), and tetraethylammonium enneacarbonyliododirhenate(O), 931–40
ETHANOATE
Metal–phenoxyalkanoic acid interactions. Part 13. Copper(II)–(2-chlorophenoxy)ethanoic acid complexes. Crystal and molecular structures of *catena*-tetra-μ-[(2-chlorophenoxy)ethanoato-*O*,*O*′]-dicopper(II), *catena*-(2-aminopyrimidine-*N*,*N*′)-tetra-μ-[(2-chlorophenoxy)ethanoato-*O*,*O*′]-dicopper(II), and 1,2,2,2,2-penta-aqua-tetra-μ-[(2-chlorophenoxy)ethanoato-*O*,*O*′]-copper(II)calcium(II), 243–52
ETHANOL
Dinuclear mono-μ-chloro-pyridyldiaza rhodium(I) complexes derived from pyridyldi-imines *via* hydrogen transfer from ethanol, 1053–6
ETHANOLATE
Crystal and molecular structure and magnetic properties of linear trimeric copper(II) complexes with predominant ferromagnetic exchange interaction, 2609–14
ETHENYL
Metallation of 2-ethenylpyridine at triosmium clusters: *X*-ray crystal structures of the open trinuclear clusters $[Os_3H(CO)_9L(NC_5H_4CH{=}CH)]$ (L = CO or PMe_2Ph), 85–90
ETHYL
Photochemistry of dicarbonyl(η⁵-cyclopentadienyl)-methyl- and -ethyl-iron and -ruthenium complexes in solutions at −30 °C and in frozen gas matrices at 12 K, 1365–74
ETHYLENEDIAMINE
Synthesis, spectroscopy, and structure of the mixed-valence complexes $[Pt^{II}(en)_2][Pt^{IV}(en)_2X_2][Pt^{III}_2(H_2P_2O_5)_4X_2]$ (en = 1,2-diaminoethane, X = Br or I), 579–86
Synthesis, characterization and acid hydrolysis of [*N*,*N*-bis(2-aminoethyl)-1,2-ethanediamine]bis(solvent)cobalt(III) complexes, 1923–8
ETHYLENEDIAMINEACETIC ACID
Potentiometric study of the complex-formation equilibria of manganese(II), cobalt(II), nickel(II), copper(II), and zinc(II) with ethylenediamine-*N*-acetic acid, 1605–8
ETHYLENEDIAMINEDIACETATEDIPROPIONATE
Correlation between structure and circular dichroism. Structure and absolute configuration of the (−) isomer of lithium (ethylenediamine-*N*,*N*′-diacetato-*N*,*N*′-di-3-propionato)rhodate(III) pentahydrate, 861–4
ETHYLENEDIAMINETETRA-ACETATE
Kinetic and structural investigations of $[Fe^{III}(edta)]^-$ [edta = ethylenediaminetetra-acetate(4−)] catalysed decomposition of hydrogen peroxide, 493–502
Electron spin resonance studies of iron(III) complexes of ethylenediaminetetra-acetate and *N*-(2-hydroxyethyl)ethylenediamine-*NN*′*N*′-triacetate in co-ordinating solvents, 1077–80
ETHYLIDYNE
The $[Fe_3(\mu_3\text{-}CR)(CO)_{10}]^-$ cluster anions as building blocks for the synthesis of mixed-metal clusters. Part 1. Synthesis of mixed clusters $[MFe_3(\mu_3\text{-}CMe)(CO)_{10}(PPh_3)]$ (M = Cu or Au) and crystal structure of $[CuFe_3(\mu_3\text{-}CMe)(CO)_{10}(PPh_3)]$, 2521–4
ETHYNYL
Cluster chemistry. Part 32. Synthesis and *X*-ray crystal structure of $[Ru_5(\mu_5\text{-}\eta^2\text{-}C_2PPh_2\text{-}P)(\mu\text{-}PPh_2)(CO)_{13}]$, a complex containing an alkynyl ligand in extended interaction with an open Ru_5 cluster, 1229–34
EXCHANGE
Co-ordinative interactions in chelated complexes of silicon. Part 7. Dynamic nuclear magnetic resonance studies of fluorine exchange at five-co-ordinated silicon, 5–8
Intercalation of alkylamines into tin(IV) bis(hydrogenphosphate) monohydrate, 213–4
Two-dimensional nuclear magnetic resonance for the analysis of the carbon-13 spectra of carbonyl groups in metallocarbonyls, 225–8
Proton exchange and base hydrolysis of *syn,anti-cis*-dichloro(1,4,7,10-tetra-azacyclododecane)cobalt(III) cations, 475–8
Thermodynamic treatment of exchange of H^+ and K^+ in α-titanium phosphate, 1865–8
Oxygen exchange and protonation of polyanions: a multinuclear magnetic resonance study of tetradecavanadophosphate(9−) and decavanadate(6−), 1953–8
EXTENDED HÜCKEL
A theoretical study of three-centre M–H–C interactions, 1637–44
EXTRACTION
The extraction and transport of metal ions by 6,6′-diamino-2,2′-bipyridine derivatives, 373–8
The concentration and selective extraction of copper(II), rhodium(III), and iridium(III) using a copolymer functionalised with dithiocarbamate groups. Spectroscopic evidence for the nature of the binding sites, 1655–60
FERRITE
Formation of oxidized molybdenum-bearing ferrites, $Fe_{2.95-x}Mo_xO_4$ (x = 0.03—0.30) in aqueous suspensions by air oxidation and valence state of molybdenum ions in the lattice, 1713–6
FERROCENE
Synthesis and structural studies of some 1,1′-dichloroferrocene

FERROCENE (contd)
derivatives of platinum(II). Crystal and molecular structure of 2,2-μ-[(1—2,5—6-η-*cis,cis*-cyclo-octa-1,5-diene)platinio]-bis(1,1′-dichloroferrocene), 1527–30
FLUORIDE
Preparation and characterisation of adducts of bismuth pentafluoride and antimony pentafluoride by vibrational spectroscopy, *X*-ray powder diffraction, and single-crystal *X*-ray crystallography, 9–16
Partially oxidized group 3B fluorometallophthalocyanines, 269–74
Synthesis and structural assessment of ammonium and caesium difluorodioxoperoxouranates(VI), $A_2[UO_2(O_2)F_2]$ (A = NH_4 or Cs), and alkali-metal difluorodioxoperoxouranate(VI) monohydrates, $A_2[UO_2(O_2)F_2]\cdot H_2O$ (A = K or Rb), 409–12
Characterisation of chromium(VI) oxide tetrafluoride, $CrOF_4$, and caesium pentafluoro-oxochromate(VI) $Cs[CrOF_5]$, 529–34
The molecular structure of gaseous bis(trifluoromethyl)selenium difluoride as determined by electron diffraction, 941–6
A tin-119 Mössbauer study of tin(II) fluoride, 1275–6
Spectroscopic studies on the higher binary fluorides of chromium: CrF_4, CrF_5, and CrF_6, both in the solid state and isolated in inert gas matrices, 1443–50
The preparation and crystal structure of (β)6$SbF_5\cdot 5SbF_5$, 1623–32
FLUORINE-19
Co-ordinative interactions in chelated complexes of silicon. Part 7. Dynamic nuclear magnetic resonance studies of fluorine exchange at five-co-ordinated silicon, 5–8
FLUOROPHOSPHINE
Fluorophosphine complexes of rhodium(I) and iridium(I): towards the design of systems with extended metal–metal interactions. The crystal structure of $[\{IrCl(PF_3)_2\}_2]$, 1295–302
FLUOROSULPHURIC ACID
Methyltin hydrides in fluorosulphuric acid, 2671–6
FLUXIONALITY
Synthesis of $[Fe(\eta^4\text{-}C_8H_8)(CO)_2L]$ [L = MeNC, Pr^{i}NC, Bu^{t}NC, PhNC, 2,6-$Me_2C_6H_3NC$, or $P(OMe)_3$] and proof of the applicability of the Woodward–Hoffmann rules to the fluxionality of $[Fe(\eta^4\text{-}C_8H_8)(CO)_2(CNPr^i)]$, 693–8
Mono- and di-nuclear rhodium and palladium complexes of macrocyclic ligands containing the 2,6-di(thiomethyl)pyridine sub-unit, 1517–22
Rhenium nitrosyl complexes with simple and with sterically demanding aromatic thiolate ligands: *X*-ray crystal structures of $[PPh_4][Re_2(SC_6H_4Me\text{-}4)_7(NO)_2]\cdot CH_2Cl_2$ and $[Re(SC_6H_3Pr^i_2\text{-}2,6)_4(NO)]$, 1533–42
Pyramidal inversions and 1,2-metal shifts in pentacarbonyl-chromium, -molybdenum, and -tungsten derivatives of dialkyl disulphides and dialkyl diselenides. A nuclear magnetic resonance investigation, 1561–8
Dynamic nuclear magnetic resonance study of Group 6 metal pentacarbonyl complexes. Pyramidal inversion and 1,2-metal shifts in the complexes $[M(CO)_5(Me_2\overbrace{CCH_2EECH_2}^{})]$ (M = Cr, Mo, or W; E = S or Se), 1569–76
Bis(triphenylphosphine)copper(I) derivatives of substituted *arachno* nine-vertex borane anions, $Cu(PPh_3)_2(B_9H_{13}X)$ (X = H, NCS, NCSe, $NCBPh_3$, $NCBH_3$, or $NCBH_2NCBH_3$), 1903–8
FORMYL
The preparation and hydridic reduction of dicationic dicarbonyl complexes of osmium(II); the crystal and molecular structure of *trans*-bis[1,2-bis(diphenylphosphino)ethane-*PP′*]carbonylformylosmium(II) hexafluoroantimonate–dichloromethane (1/1), 387–94
Reactions of hydride reagents with alkylmolybdenum carbonyl complexes. Reaction of $LiBHEt_3$ with $[MoMe(CO)_3(\eta\text{-}C_5H_5)]$, formation of an anionic acetaldehyde complex, and a stoicheiometric cycle for the synthesis of acetaldehyde, 1815–20
FRAGMENTATION
Complexes of the platinum metals. Part 30. Fragmentation reactions of rhodium and iridium trichloro- and tribromo-acetates, 2113–20
FUEL CELL
Hydrogen generation by hydrolysis of sodium tetrahydroborate: effects of acids and transition metals and their salts, 307–14
FULVALENE
Stereochemical consequences of the two-electron oxidation of a dirhodium fulvalene complex: the *X*-ray crystal structures of *trans*-$[Rh_2(CO)_2(PPh_3)_2(\eta^5\text{:}\eta'^5\text{-}C_{10}H_8)]$ and *cis*-$[Rh_2(CO)_2(PPh_3)_2(\eta^5\text{:}\eta'^5\text{-}C_{10}H_8)][PF_6]_2$, 2283–90
FULVENE
Synthesis, structure, and bonding of fulvene complexes of titanium, molybdenum, and tungsten, 2037–50
FURYL
The chemistry of heteroarylphosphorus compounds. Part 16. Unusual substituent effects on selenium-77 nuclear magnetic resonance chemical shifts of heteroaryl- and aryl-phosphine selenides. *X*-Ray crystal structure of tri(2-furyl)phosphine selenide, 2505–8
GALLIUM
Partially oxidized group 3B fluorometallophthalocyanines, 269–74
The molecular structure of gaseous dimethylamidogallane: characterization of the dimer $[Me_2NGaH_2]_2$ by electron diffraction and vibrational spectroscopy, 807–14
Kinetics and mechanisms of complex formation of gallium(III) and indium(III). The reactions with 4-(2-pyridylazo)resorcinol in water and other mixed solvents, 2615–22
GERMANIUM
Subvalent group 4B metal alkyls and amides. Part 7. Transition-metal chemistry of metal(II) bis(trimethylsilyl)amides $M'(NR_2)_2$ (R = $SiMe_3$; M′ = Ge, Sn, or Pb), 51–8
GLASS
Magnetic circular dichroism spectra of tris-chelate complexes of 2,2′-bipyridyl and 1,10-phenanthroline with iron(II), ruthenium(II), and osmium(II) at 4.2 K, 1781–8
GLUTATHIONE
Synthesis and characterization of some chromium(III) complexes with glutathione, 2085–90
GLYCINE
Glycine complexation with uranyl ion: absorptiometric, luminescence, and *X*-ray structural studies of tetrakis-(glycine)dioxouranium(VI) nitrate, 517–22
GOLD
Bimetallic systems. Part 8. Heterobimetallic di-isonitrile or isonitrile–carbonyl complexes of rhodium or iridium bridged to silver or gold by $Ph_2PCH_2PPh_2$, 511–6
Displacement of pyridine-2-methanol from dichloro(pyridine-2-methanolato)gold(III) in acidic solution. Ring opening at oxygen, 731–6
Short hydrogen bonds: diadducts of substituted pyridine *N*-oxides: synthesis, spectroscopic studies, and *X*-ray structure, 749–54
Solvent extraction of gold and platinum-group metals using 2-nonylpyridine 1-oxide, and the crystal and molecular structure of bis(2-nonylpyridine 1-oxide)hydrogen(1+) tetrachloroaurate(III), 771–6
Co-ordination chemistry of pyridyl and *N*-methylimidazolyl ketones. Synthetic and *X*-ray structural studies of copper(II), nickel(II), and dimethylgold(III) complexes, 981–6
Synthesis of gold-(I) and -(III) complexes with carbonyl-stabilized phosphorus ylides. Crystal structure of $[\{Au(PPh_3)\}_2\{\mu\text{-}C(PPh_3)CO_2Et\}]ClO_4$, 1163–8
Co-ordination chemistry of dimethylgold(III). Synthesis, structural studies, and fluxional behaviour of complexes with polydentate ligands, 1183–90
Gold(I) complexes of 1,2-ditertiary phosphino-*o*-carborane ligands: two-, three-, and four-co-ordination, 1387–90
Studies of gold cluster compounds using high-resolution ^{31}P solid-state nuclear magnetic resonance spectroscopy, 1811–4
The preparation of trichloro(phenyl 2-pyridylmethyl sulphoxide)gold(III) and the kinetics and equilibria of its formation and subsequent reactions, 2091–4
Synthesis of trinuclear gold(I) and gold(III) complexes containing the tridentate bis(diphenylphosphino)methanide ligand. Crystal structure of $[Cl(C_6F_5)_2Au\{Ph_2PCH(AuNC_5H_5)PPh_2\}AuCl]$, 2417–20
The $[Fe_3(\mu_3\text{-}CR)(CO)_{10}]^-$ cluster anions as building blocks for the synthesis of mixed-metal clusters. Part 1. Synthesis of mixed clusters $[MFe_3(\mu_3\text{-}CMe)(CO)_{10}(PPh_3)]$ (M = Cu or Au) and crystal structure of $[CuFe_3(\mu_3\text{-}CMe)(CO)_{10}(PPh_3)]$, 2521–4
Novel ring compounds of bidentate phosphines with gold(I). Two-, three-, and four-co-ordination, 2655–60
HAEM
Rate parameters for oxygen and carbon monoxide binding to a liposome-embedded heme under physiological conditions, 65–8
HALIDE
Electron spin resonance spectra of manganese(II) ions in the double-stranded chain polymers $CdLX_2$ (L = pyridine or methylpyridine; X = Cl or Br), 75–80
Niobium(IV) sulphidohalides: preparation of $Nb_2X_4S_3$ and

HALIDE (contd)
$Nb_2X_4S\cdot nL$[X = Br or Cl; n = 4, L = NCMe, SMe_2, or tetrahydrothiophene (tht); n = 2, L = $PhSCH_2CH_2SPh$]. Crystal and molecular structure of $Nb_2Cl_4S_2\cdot 4tht$, 417–22
The synthesis of $NR_4[(Ta_6Cl_{12})(H_2O)_6]X_4$ (R = Me or Et, X = Cl or Br) by the spontaneous reduction of $[Ta_6Cl_{12}]^{4+}$ to $[Ta_6Cl_{12}]^{3+}$ in acidic media: *X*-ray structure analysis of $NMe_4[(Ta_6Cl_{12})(H_2O)_6]Br_4$, 455–8
Palladium(II,IV) mixed-valence complexes of 1,2-diaminoethane, 1,3-diaminopropane, and diethylenetriamine: syntheses, electronic, infrared, Raman, and resonance Raman spectra and *X*-ray studies, 815–20
Lewis-base adducts of group 1B metal(I) compounds. Part 16. Synthesis, structure, and solid-state phosphorus-31 nuclear magnetic resonance spectra of some novel $[Cu_4X_4L_4]$ (X = halogen, L = N,P base) 'cubane' clusters, 831–8
Thermochemical data for adducts of zinc, cadmium, and mercury halides with hexamethylphosphoramide, 1103–6
Mössbauer study of the *cis–trans* isomers of tin(IV) complexes. Some considerations about the sign of the electric-field gradient, 1281–4
Synthesis and properties of the divalent 1,2-bis(dimethylphosphino)ethane (dmpe) complexes $MCl_2(dmpe)_2$ and $MMe_2(dmpe)_2$ (M = Ti, V, Cr, Mn, or Fe). *X*-Ray crystal structures of $MCl_2(dmpe)_2$ (M = Ti, V, or Cr), $MnBr_2(dmpe)_2$, $TiMe_{1.3}Cl_{0.7}(dmpe)_2$, and $CrMe_2(dmpe)_2$, 1339–48
The synthesis of $NR_4[(Ta_6Cl_{12})(H_2O)_6]X_4$ (R = Me or Et, X = Cl or Br) by the spontaneous reduction of $[Ta_6Cl_{12}]^{4+}$ to $[Ta_6Cl_{12}]^{3+}$ in acidic media: *X*-ray structure analysis of $NMe_4[(Ta_6Cl_{12})(H_2O)_6]Br_4$ (1985, 455), 1531–2
Structure and solvation of mercury(II) iodide, bromide, and chloride in pyridine solution; refinement of the crystal structure of di-iodobis(pyridine)mercury(II), $[HgI_2(py)_2]$, 1597–604
Preparation and vibrational spectra of $[OsX_6]^{3-}$ (X = Cl, Br, or I) and of other platinum-group hexahalogeno-complexes, 1673–6
High-co-ordination-number compounds of niobium and tantalum: Reactions of niobium and tantalum halides and sulphido-halides with sodium diethyldithiocarbamate. The crystal structures of $[Nb(S_2CNEt_2)_4]Br$, $Nb(S_2CNEt_2)_3S$, and $Ta(S_2CNEt_2)_3(S_2)$, 1821–8
New carbonyl derivatives of niobium(I) and tantalum(I), 1989–96
On the synthesis of dimethylphenylphosphine complexes of manganese(II) halides, 2203–4
The co-ordination of small molecules by manganese(II) phosphine complexes. Part 3. The dependence on the nature of the halogen in $[MnX_2(PR_3)]$ (X = Cl, Br, or I; R_3 = $PhMe_2$, $PhEt_2$, $PhPr^n_2$, $PhBu^n_2$, $PhBu^i_2$, Pr^n_3, Bu^n_3, or Ph_3) on adduct formation with sulphur dioxide in the solid state and in tetrahydrofuran solution, 2661–4

HEPTANETRIONE
Reactions of metal ions with triketones. Part 2. Kinetics and mechanisms of the reactions of nickel(II) and cobalt(II) with heptane-2,4,6-trione in methanol–water (70:30 v/v), 2565–70

HETEROPOLYTUNGSTATE
Heteropolytungstates as catalysts for the photochemical reduction of oxygen and water, 395–400

HEXABORANE
A kinetic study of the gas-phase thermolysis of hexaborane(10), 541–8

HEXAFLUOROANTIMONATE
The preparation and hydridic reduction of dicationic dicarbonyl complexes of osmium(II); the crystal and molecular structure of *trans*-bis[1,2-bis(diphenylphosphino)ethane-*PP'*]carbonylformylosmium(II) hexafluoroantimonate–dichloromethane (1/1), 387–94

HEXAMETHYLPHOSPHORAMIDE
Thermochemical data for adducts of zinc, cadmium, and mercury halides with hexamethylphosphoramide, 1103–6

HYDRAZIDE
Molybdenum, rhenium, and tungsten complexes with bi- and tridentate phosphinothiolato-ligands; structures of $[Mo\{PhP(CH_2CH_2S)_2\}_2]$ and $[Mo(NNMe_2)\{PhP(CH_2CH_2S)_2\}_2]$, 2647–54

HYDRAZONE
The template synthesis and crystal and molecular structure of a seven-co-ordinate manganese(II) complex with 2,6-diacetylpyridine mono(2-aminobenzoylhydrazone), 215–8
Dynamic behaviour of carbon-metallated palladium hydrazone complexes. Crystal structures of $[\{Pd[CH_2CMe_2C(=N\text{–}NMePh)Me]Cl\}_2]$ and $[\{Pd[CH_2C(=N\text{–}NMePh)Bu^t]Cl\}_2]$, 1155–62
Investigation into aroylhydrazones as chelating agents. Part 7. Synthesis and spectroscopic characterization of complexes of Mn^{II}, Co^{II}, Ni^{II}, Cu^{II}, and Zn^{II} with 2,6-diacetylpyridine bis(2-aminobenzoylhydrazone) and *X*-ray structure of chloro[2,6-diacetylpyridine bis(2-aminobenzoylhydrazone)](methanol)manganese(II) chloride monohydrate, 2387–92

HYDRIDE
Chemistry of the unsaturated cluster compound $[Os_3Pt(\mu\text{-}H)_2(CO)_{10}\{P(cyclo\text{-}C_6H_{11})_3\}]$; *X*-ray crystal structures of $[Os_3Pt(\mu\text{-}H)_2(CO)_{11}\{P(cyclo\text{-}C_6H_{11})_3\}]$, $[Os_3Pt(\mu\text{-}H)_4(CO)_{10}\{P(cyclo\text{-}C_6H_{11})_3\}]$, and $[Os_3Pt(\mu\text{-}H)_2(\mu\text{-}CH_2)(CO)_{10}\{P(cyclo\text{-}C_6H_{11})_3\}]$ (two isomers), 177–90
Two-dimensional nuclear magnetic resonance for the analysis of the carbon-13 spectra of carbonyl groups in metallocarbonyls, 225–8
Systematic synthesis of tetranuclear osmium clusters by the reaction of trinuclear clusters with $[OsH_2(CO)_4]$; crystal structure of $[Os_4H_3Br(CO)_{13}]$, 555–64
Trimethylphosphine polyhydrides of tungsten and rhenium, 587–90
Synthesis, characterization, and structure of the complex $[FeH(H_2BH_2)\{CH_3C(CH_2PPh_2)_3\}]$, 605–10
Studies on transition-metal cyano-complexes. Part 4. Cyanide hydride complexes of Groups 6A and 8, 717–22
The addition of protic acids to $[Mn_2(CO)_5(Ph_2PCH_2PPh_2)_2]$ to give bridging hydrido-compounds, 743–8
Organoruthenium(II) complexes formed by insertion reactions of some vinyl compounds and conjugated dienes into a hydrido–ruthenium bond, 873–8
Alkyl, hydrido, and tetrahydroaluminato complexes of manganese with 1,2-bis(dimethylphosphino)ethane (dmpe). *X*-Ray crystal structures of $Mn_2(\mu\text{-}C_6H_{11})_2(C_6H_{11})_2(\mu\text{-dmpe})$, $(dmpe)_2Mn(\mu\text{-}H)_2AlH(\mu\text{-}H)_2AlH(\mu\text{-}H)_2Mn(dmpe)_2$, and $Li_4\{MnH(C_2H_4)[CH_2(Me)PCH_2CH_2PMe_2]_2\}_2\cdot 2Et_2O$, 921–30
Electron-transfer reactions in nitrogen fixation. Part 1. The electrosynthesis of dinitrogen, hydride, isocyanide, and carbonyl complexes of molybdenum: intermediates, mechanisms, and energetics, 1255–64
Homo- and hetero-dinuclear hydride-bridged complexes containing cyclo-octadiene: the crystal and molecular structure of $[(\eta^4\text{-}C_8H_{12})Ir(\mu\text{-}H)(\mu\text{-}Cl)IrH_2(PPh_3)_2]$, 1277–80
The reaction of $[Os_3(\mu\text{-}H)_2(CO)_9L]$ (L = CO or PEt_3) with dimethylcyanamide, Me_2NCN: *X*-Ray crystal structure of $[Os_3(\mu\text{-}H)(\mu\text{-}NCHNMe_2)(CO)_{10}]$ and the reactions of this complex with acids, 1355–60
Bimetallic cyano-bridged cations: preparation and hydride reduction of $[(\eta^5\text{-}C_5H_5)L_2Ru(\mu\text{-}CN)ML'_2(\eta^5\text{-}C_5H_5)]PF_6$ [L_2, L'_2 = $(PPh_3)_2$, $Ph_2PCH_2CH_2PPh_2$; M = Ru or Fe]. Formation of $[Ru(\eta^5\text{-}C_5H_5)(PPh_3)H_3]$ and *X*-ray crystal structure of $[(\eta^5\text{-}C_5H_5)(Ph_2PCH_2CH_2PPh_2)Ru(\mu\text{-}CN)Ru(PPh_3)_2(\eta^5\text{-}C_5H_5)]PF_6$, 1479–86
Reactivity of the unsaturated anion decacarbonyltetra-μ-hydrido-trirhenate(1−) toward phenols. Crystal and molecular structures of the tetraethylammonium salts of the triangular cluster anion $[Re_3(\mu\text{-}H)_3(\mu\text{-}OC_6F_5)(CO)_{10}]^-$ and of the binuclear anion $[Re_2(\mu\text{-}OC_6H_5)_3(CO)_6]^-$, 1507–12
Addition of proton and boron trifluoride at terminal and bridging cyano-ligands in dinuclear manganese compounds, 1609–12
A theoretical study of three-centre M–H–C interactions, 1637–44
Reactions of hydride reagents with alkylmolybdenum carbonyl complexes. Reaction of $LiBHEt_3$ with $[MoMe(CO)_3(\eta\text{-}C_5H_5)]$, formation of an anionic acetaldehyde complex, and a stoicheiometric cycle for the synthesis of acetaldehyde, 1815–20
The chemistry and catalytic properties of ruthenium and osmium complexes. Part 1. Homogeneous catalysis of organic reactions by bromo(carbonyl)hydridotris(triphenylphosphine)osmium(II), 1859–64
Carbon-13 nuclear magnetic resonance evidence of a relaxation process dominated by scalar coupling with a quadrupolar nucleus in $[Re_3(\mu\text{-}H)_4(CO)_{10}]^-$, 1899–902
The mechanisms of the reactions between hydrido-complexes and diazonium salts. Part 1. The reactions of *trans*-$[PtH(Cl)(PEt_3)_2]$, 2059–66
The mechanisms of the reactions between hydrido-complexes and diazonium salts. Part 2. The reactions of *cis,mer*-$[RhHCl_2(PEtPh_2)_3]$, 2067–78
Bimetallic systems. Part 13. Platinum–manganese carbonyl complexes containing bridging $Ph_2PCH_2PPh_2$(dppm) ligands:

HYDRIDE (contd)
crystal structure of $[(OC)_3Mn(\mu\text{-dppm})_2PtH(Br)]BF_4$, 2131–8
Synthesis of tetra- and penta-nuclear platinum–osmium carbido-cluster complexes from non-carbido-precursors: *X*-ray structural studies on $[Os_3Pt(\mu\text{-}H)_2(\mu_4\text{-}C)(CO)_{10}\{P(cyclo\text{-}C_6H_{11})_3\}]$, $[Os_3Pt_2(\mu\text{-}H)_2(\mu_5\text{-}C)(\mu\text{-}CO)(CO)_9\{P(cyclo\text{-}C_6H_{11})_3\}_2]$, and $[Os_3Pt_2(\mu\text{-}H)(\mu_5\text{-}C)(\mu\text{-}OMe)(\mu\text{-}CO)(CO)_9\{P(cyclo\text{-}C_6H_{11})_3\}_2]$, 2437–48
Acid-induced displacement of acetaldehyde from a μ-vinyloxy-triosmium cluster, 2479–82
Methyltin hydrides in fluorosulphuric acid, 2671–6
HYDROGEN-1
Aluminium-27 and hydrogen-1 nuclear magnetic resonance studies of solutions of aluminium salts in alcohol–chloroform mixtures, 591–6
HYDROGEN BONDING
Short hydrogen bonds: diadducts of substituted pyridine *N*-oxides: synthesis, spectroscopic studies, and *X*-ray structure, 749–54
HYDROGEN PEROXIDE
Coloured species formed from the titanium(IV)-4-(2′-pyridylazo)resorcinol reagent in the spectrophotometric determination of trace amounts of hydrogen peroxide, 81–4
Kinetic and structural investigations of $[Fe^{III}(edta)]^-$ [edta = ethylenediaminetetra-acetate(4−)] catalysed decomposition of hydrogen peroxide, 493–502
Studies on singlet oxygen in aqueous solution. Part 1. Formation of singlet oxygen from hydrogen peroxide with two-electron oxidants, 1141–6
Studies on singlet oxygen in aqueous solution. Part 3. The decomposition of peroxy-acids, 1151–4
Studies on singlet oxygen in aqueous solution. Part 4. The 'spontaneous' and catalysed decomposition of hydrogen peroxide, 2525–30
HYDROGENPHOSPHATE
Intercalation of alkylamines into tin(IV) bis(hydrogenphosphate) monohydrate, 213–4
HYDROGEN TRANSFER
Dinuclear mono-μ-chloro-pyridyldiaza rhodium(I) complexes derived from pyridyldi-imines *via* hydrogen transfer from ethanol, 1053–6
HYDROLYSIS
Hydrogen generation by hydrolysis of sodium tetrahydroborate: effects of acids and transition metals and their salts, 307–14
Proton exchange and base hydrolysis of *syn,anti-cis*-dichloro(1,4,7,10-tetra-azacyclododecane)cobalt(III) cations, 475–8
An equation for predicting the formation constants of hydroxo-metal complexes, 723–30
Synthesis, characterization and acid hydrolysis of [*N*,*N*-bis(2-aminoethyl)-1,2-ethanediamine]bis(solvent)cobalt(III) complexes, 1923–8
The hydrolysis of metal ions. Part 8. Aluminium(III), 1967–70
HYDROXAMATE
Mono- and bi-nuclear hydroxamates of bis(2-phenylazopyridine)ruthenium(II), 361–8
HYDROXIDE
Co-ordination chemistry of higher oxidation states. Part 13. Synthesis and properties of alkali-metal hydroxo-oxo-osmate(VIII) compounds and the molecular structure of $Cs[O_4Os(\mu\text{-}OH)OsO_4]$, 199–204
An equation for predicting the formation constants of hydroxo-metal complexes, 723–30
A disagreement on the explanation of short and long As–O bonds of the $(As\text{–}O)_4$ ring in $As_4(CF_3)_6O_6(OH)_2$ in terms of As^{III}–O and As^{V}–O bonds, 2221–2
Equilibrium and structural studies of silicon(IV) and aluminium(III) in aqueous solution. Part 13. A potentiometric and ^{27}Al nuclear magnetic resonance study of speciation and equilibria in the aluminium(III)–oxalic acid–hydroxide system, 2665–70
HYDROXIDE
Complexes of lanthanoid salts with macrocyclic ligands. Part 17. Synthesis and crystal and molecular structure of a hydroxide-bridged praseodymium trifluoroacetate complex with 15-crown-5-ether, $[Pr_2(CF_3CO_2)_3(OH)(C_{10}H_{20}O_5)_2]$-$[Pr_2(CF_3CO_2)_6]$, 885–90
HYDROXYCARBONYL
Acidic and basic properties of the hydroxycarbonyl complex $[IrCl_2(CO_2H)(CO)(PMe_2Ph)_2]$, 857–60
HYDROXYLAMINE
A new reaction of nitrosyl complexes; one-electron reduction of *trans*-$[MX(NO)L_4]^{2+}$ [M = Ru or Fe, X = Cl or Br, L = pyridine or *o*-phenylenebis(dimethylarsine)] with hydroxylamine, 1733–4
HYPOPHOSPHOROUS ACID
Kinetics and mechanism of electron-transfer reactions of bismuth(V) in aqueous acidic perchlorate–fluoride media. Part 1. Oxidation of hypophosphorous acid, 2571–4
IMIDAZOLE
A mechanistic study of the reaction of iron(III) porphyrins with imidazoles. Hydrogen bonding by the propionic acid side chains in hemin chloride, 2269–76
Studies on spin-equilibrium iron(III) complexes. Part 1. Syntheses and magnetic properties of a new family of spin cross-over iron(III) complexes with a unidentate ligand over a wide range of the spectrochemical series and a quinquedentate ligand derived from salicylaldehyde and di(3-aminopropyl)amine. *X*-Ray crystal structure of [4-azaheptamethylene-1,7-bis(salicylideneiminato)](4-methylpyridine)iron(III) tetraphenylborate, 2575–84
IMIDAZOLYL
Co-ordination chemistry of dimethylgold(III). Synthesis, structural studies, and fluxional behaviour of complexes with polydentate ligands, 1183–90
IMIDE
Complexes of organoaluminium compounds. Part 13. Preparation and nuclear magnetic resonance spectra of the arylamido-compounds $AlMe_2(NHR')$ (R′ = Ph, C_6H_4Me-*o*, C_6H_4Me-*p*, $C_6H_3Me_2$-2,6) and the imido-compounds $AlMe(NR')$. Crystal and molecular structures of $[\{AlMe_2(NHC_6H_4Me\text{-}o)\}_2]$ and $[\{AlMe(NPh)\}_6]$, 1929–34
Rhenium nitrido-, arylimido-, nitrile, and carbonyl complexes with sterically hindered thiolate ligands, 2305–10
IMINE
Dinuclear mono-μ-chloro-pyridyldiaza rhodium(I) complexes derived from pyridyldi-imines *via* hydrogen transfer from ethanol, 1053–6
INDIUM
Kinetics and mechanisms of complex formation of gallium(III) and indium(III). The reactions with 4-(2-pyridylazo)resorcinol in water and other mixed solvents, 2615–22
Indium derivatives of monothio-β-diketones and the *X*-ray structure of tris[benzoyl(thiobenzoyl)methanato-*O*,*S*]indium(III), 2623–8
INSERTION
Insertion of carbon dioxide, of CO_2-like molecules, and of other unsaturated compounds into the platinum–nitrogen bond of $[Pt(PPh_3)_2(PhNO)]$, 163–8
Organoruthenium(II) complexes formed by insertion reactions of some vinyl compounds and conjugated dienes into a hydrido–ruthenium bond, 873–8
INTERCALATION
Intercalation of alkylamines into tin(IV) bis(hydrogenphosphate) monohydrate, 213–4
INVERSION
Pyramidal inversion in configurational isomers of tetracarbonyl[1,1,2,2-tetrakis(methylthio)ethane]chromium(0): a two-dimensional nuclear magnetic resonance exchange study, 2195–202
IODATE
Oxidation of thiourea by iodate: a new type of oligo-oscillatory reaction, 1669–72
IODIDE
The peroxodisulphate–iodide reaction. Reactivity and ionic association and solvation in isodielectric water–solvent mixtures, 31–4
Iodine-127 Mössbauer spectroscopy of copper(I) iodide–phosphine adducts, 1727–30
Reactions of some decaosmium clusters with electrophilic and nucleophilic reagents: *X*-ray structure analyses of $[N(PPh_3)_2][Os_{10}C(CO)_{24}(\mu\text{-}I)]$, $[Os_{10}C(CO)_{24}(\mu\text{-}I)_2]$, $[N(PPh_3)_2]_2[Os_{10}C(CO)_{22}(NO)I]$, $[Os_{10}C(CO)_{23}\{P(OMe)_3\}(\mu\text{-}I)_2]$ and of two isomers of $[Os_{10}C(CO)_{21}\{P(OMe)_3\}_4]$, 1795–810
IODINE
Thermochemistry of polyhalides. Part 5. Standard enthalpies of formation of tetramethylammonium and tetraethylammonium tetrachloroiodates, 997–1000
IODINE-127
Iodine-127 Mössbauer spectroscopy of copper(I) iodide–phosphine

IODINE-127 (contd)
adducts, 1727–30
ION EXCHANGE
Crystalline zirconium(IV) hydrogenarsenate hydrogenphosphate monohydrate: synthesis, ion-exchange properties, and thermal behaviour, 1737–42
Ion exchange of $K_4Nb_6O_{17}\cdot 3H_2O$, 2349–52
IONIC STRENGTH
On the possibility of determining the thermodynamic parameters for the formation of weak complexes using a simple model for the dependence on ionic strength of activity coefficients: Na^+, K^+, and Ca^{2+} complexes of low molecular weight ligands in aqueous solution, 2353–62
I.R.
Characterisation of chromium(VI) oxide tetrafluoride, $CrOF_4$, and caesium pentafluoro-oxochromate(VI) $Cs[CrOF_5]$, 529–34
Synthesis, spectroscopy, and structure of the mixed-valence complexes $[Pt^{II}(en)_2][Pt^{IV}(en)_2X_2][Pt^{III}_2(H_2P_2O_5)_4X_2]$ (en = 1,2-diaminoethane, X = Br or I), 579–86
The molecular structure of gaseous dimethylamidogallane: characterization of the dimer $[Me_2NGaH_2]_2$ by electron diffraction and vibrational spectroscopy, 807–14
Isolated CH stretching frequencies, methyl group geometry, and methyl CH^- bond lengths and strengths in tricarbonyl(η^5-cyclopentadienyl)methylchromium(II), -molybdenum(II), and -tungsten(II), 1207–12
Spectroscopic studies on the higher binary fluorides of chromium: CrF_4, CrF_5, and CrF_6, both in the solid state and isolated in inert gas matrices, 1443–50
Preparation and vibrational spectra of $[OsX_6]^{3-}$ (X = Cl, Br, or I) and of other platinum-group hexahalogeno-complexes, 1673–6
IRIDATETRABORANE
The crystal and molecular structure of [1,1,1,1-(CO)H$(PPh_3)_2$-*arachno*-1-IrB_3H_7] and some bonding considerations in *arachno*-type four-vertex metal 'π-allyl' and 'borallyl' clusters, 1843–8
IRIDIUM
Bimetallic systems. Part 8. Heterobimetallic di-isonitrile or isonitrile–carbonyl complexes of rhodium or iridium bridged to silver or gold by $Ph_2PCH_2PPh_2$, 511–6
Complexes of the platinum metals. Part 23. Synthesis of the nitrosyl carboxylate complexes $[M(O_2CR)_2(NO)(PPh_3)_2]$ (M = Rh or Ir; R = CF_3, C_2F_5, or C_6F_5): crystal and molecular structures of the trifluoroacetate derivatives $[M(O_2CCF_3)_2(NO)(PPh_3)_2]$, 611–6
Complexes of the platinum metals. Part 24. The role of dioxygen in the reactions of trifluoroacetic acid with the rhodium and iridium nitrosyls $[M(NO)(PPh_3)_3]$, 617–20
Studies on transition-metal cyano-complexes. Part 4. Cyanide hydride complexes of Groups 6A and 8, 717–22
Acidic and basic properties of the hydroxycarbonyl complex $[IrCl_2(CO_2H)(CO)(PMe_2Ph)_2]$, 857–60
Homo- and hetero-dinuclear hydride-bridged complexes containing cyclo-octadiene: the crystal and molecular structure of $[(\eta^4\text{-}C_8H_{12})Ir(\mu\text{-}H)(\mu\text{-}Cl)IrH_2(PPh_3)_2]$, 1277–80
Fluorophosphine complexes of rhodium(I) and iridium(I): towards the design of systems with extended metal–metal interactions. The crystal structure of $[\{IrCl(PF_3)_2\}_2]$, 1295–302
The concentration and selective extraction of copper(II), rhodium(III), and iridium(III) using a copolymer functionalised with dithiocarbamate groups. Spectroscopic evidence for the nature of the binding sites, 1655–60
Kinetics and mechanism of the oxidation of substituted malonic acids by the hexachloroiridate(IV) ion, 1683–8
The crystal and molecular structure of [1,1,1,1-(CO)H$(PPh_3)_2$-*arachno*-1-IrB_3H_7] and some bonding considerations in *arachno*-type four-vertex metal 'π-allyl' and 'borallyl' clusters, 1843–8
1*H*-pyrrolo[2,3-*b*]pyridine (HL) ligands in rhodium(I) and iridium(I) chemistry. Crystal and molecular structures of $[Rh_2(\mu\text{-}L)_2(nbd)_2]$ and $[Rh_4(\mu\text{-}Cl)_2(\mu\text{-}L)_2(\mu\text{-}CO)_2(CO)_2(nbd)_2]$, 1891–8
Complexes of the platinum metals. Part 30. Fragmentation reactions of rhodium and iridium trichloro- and tribromo-acetates, 2113–20
Reactions of co-ordinated ligands. Part 35. Evidence for carbon–carbon double-bond cleavage of cyclopropenes in their reaction with dinuclear cobalt, rhodium, and iridium complexes; crystal structure and protolysis of $[Rh_2(\mu\text{-}CO)(\mu\text{-}COCHCMe_2CH)(\mu\text{-}C_5Me_5)_2]$, 2483–92
IRON
Rate parameters for oxygen and carbon monoxide binding to a liposome-embedded heme under physiological conditions, 65–8
Phosphocholine-substituted 5,10,15,20-tetraphenylporphyrinatoiron(II): oxygen carrier under physiological conditions, 275–8
Macrocyclic polyphosphane ligands. Iron(II), cobalt(II), and nickel(II) complexes of (4*RS*,7*RS*,13*SR*,16*SR*)-tetraphenyl-1,10-dipropyl-1,10-diaza-4,7,13,16-tetraphosphacyclo-octadecane: crystal structures of their tetraphenylborate derivatives, 479–86
Kinetic and structural investigations of $[Fe^{III}(edta)]^-$ [edta = ethylenediaminetetra-acetate(4−)] catalysed decomposition of hydrogen peroxide, 493–502
Synthesis, characterization, and structure of the complex $[FeH(H_2BH_2)\{CH_3C(CH_2PPh_2)_3\}]$, 605–10
Synthesis of $[Fe(\eta^4\text{-}C_8H_8)(CO)_2L]$ [L = MeNC, Pr^{i}NC, Bu^{t}NC, PhNC, 2,6-$Me_2C_6H_3NC$, or $P(OMe)_3$] and proof of the applicability of the Woodward–Hoffmann rules to the fluxionality of $[Fe(\eta^4\text{-}C_8H_8)(CO)_3(CNPr^i)]$, 693–8
Reduction–oxidation properties of organotransition-metal complexes. Part 20. Oxidative and thermolytic cyclopropane ring-opening reactions; *X*-ray crystal structure of $[Fe_2(CO)_6(\eta^4:\eta'^4\text{-}C_{16}H_{18})]$, 699–706
Difluorophosphate complexes of chromium, manganese, iron, cobalt, and nickel, 707–10
Annelation of ring-opened arylcyclopropenium ions to co-ordinated cyclo-octatetraene, and the *X*-ray crystal structure of $[Fe(CO)_3(\sigma,\eta^3\text{-}C_{11}H_9Ph_3)]$, 777–82
Transfer chemical potentials for Fe^{2+} from water into aqueous methanol: a comment, 867–6
Transfer chemical potentials for Fe^{2+} from water into aqueous methanol: a reply, 867–70
The red form of bis(1,10-phenanthroline)dithiocyanatoiron(II), 991–6
Reduction–oxidation properties of organotransition-metal complexes. Part 22. Stereospecific oxidative cyclopropane ring opening and reductive cyclobutane ring formation in polycyclic hydrocarbon complexes of iron; *X*-ray crystal structures of $[Fe_2(CO)_6(\eta^5:\eta'^5\text{-}C_{16}H_{16})][PF_6]_2\cdot CH_3NO_2$ and $[Fe_2(CO)_6(\eta^4:\eta'^4\text{-}C_{16}H_{16})]$, 1027–36
Electron spin resonance studies of iron(III) complexes of ethylenediaminetetra-acetate and *N*-(2-hydroxyethyl)ethylenediamine-*NN'N'*-triacetate in co-ordinating solvents, 1077–80
Carbon–carbon formation at di-iron centres. Part 2. Reactivity of $[Fe_2(CO)_6(\mu\text{-}COEt)\{\mu\text{-}C(R)C(R)H\}]$ complexes toward MeOC(O)C≡CC(O)OMe (R = Ph) and $CF_3C{\equiv}CCF_3$ (R = Ph or H); *X*-ray crystal structures of $[Fe_2(CO)_5\{\mu\text{-}C(OEt)C[C(O)OMe]C[C(O)OMe]\}\{\mu\text{-}C(Ph)C(Ph)H\}]\cdot H_2O$ and $[Fe_2(CO)_6\{\mu\text{-}C(CF_3)C(CF_3)CHCHCH(OMe)\}]$, 1087–94
Equilibrium and kinetic study of pyridine binding to phthalocyaninatoiron(II) in dimethyl sulphoxide, 1107–12
Equilibrium and kinetic study of the reaction between phthalocyaninatoiron(II) and carbon monoxide in dimethyl sulphoxide in the presence of pyridine. Evidence for the formation of a transient, 1113–8
On the effect of cyanide ion on the reaction of pentacyanonitrosylferrate(2−) with cysteine, 1191–4
Synthesis, structures, and reactivities of some pentamethylcyclopentadienyl–sulphur compounds, 1303–8
Chemistry of di- and tri-metal complexes with bridging carbene or carbyne ligands. Part 34. Reactions of sulphur and selenium with the di-irontungsten complexes $[Fe_2W(\mu_3\text{-}CR)(\mu\text{-}CO)(CO)_8(\eta\text{-}C_5H_5)]$ (R = C_6H_4Me-4 or Me); crystal structures of $[Fe_2W(\mu\text{-}CC_6H_4Me\text{-}4)(\mu_3\text{-}S)(CO)_7(\eta\text{-}C_5H_5)]$ and $[Fe_2W(\mu_3\text{-}SCMe)(CO)_8(\eta\text{-}C_5H_5)]$, 1323–30
Photochemistry of dicarbonyl(η^5-cyclopentadienyl)-methyl- and -ethyl-iron and -ruthenium complexes in solutions at −30 °C and in frozen gas matrices at 12 K, 1365–74
Magnetic exchange interactions in perovskite solid solutions. Part 5. The unusual defect structure of $SrFeO_{3-y}$, 1455–70
Bimetallic cyano-bridged cations: preparation and hydride reduction of $[(\eta^5\text{-}C_5H_5)L_2Ru(\mu\text{-}CN)ML'_2(\eta^5\text{-}C_5H_5)]PF_6$ [L_2, L'_2 = $(PPh_3)_2$, $Ph_2PCH_2CH_2PPh_2$; M = Ru or Fe]. Formation of $[Ru(\eta^5\text{-}C_5H_5)(PPh_3)H_3]$ and *X*-ray crystal structure of $[(\eta^5\text{-}C_5H_5)(Ph_2PCH_2CH_2PPh_2)Ru(\mu\text{-}CN)Ru(PPh_3)_2(\eta^5\text{-}C_5H_5)]PF_6$, 1479–86
Synthesis and structural studies of some 1,1'-dichloroferrocene derivatives of platinum(II). Crystal and molecular structure of 2,2-μ-[(1—2,5—6-η-*cis*,*cis*-cyclo-octa-1,5-diene)platinio]-bis(1,1'-dichloroferrocene), 1527–30

IRON (contd)
Geometry-dependent carbon-13 chemical shifts in (η^6-[2.2]cyclophane)(η^5-cyclopentadienyl)iron(II) hexafluorophosphates, 1661–4
Formation of oxidized molybdenum-bearing ferrites, $Fe_{2.95-x}Mo_xO_4$ (x = 0.03—0.30) in aqueous suspensions by air oxidation and valence state of molybdenum ions in the lattice, 1713–6
Studies on the regioselective preparation of tricarbonyl(cyclohexadienyl)iron(1+) salts, 1723–6
A new reaction of nitrosyl complexes; one-electron reduction of *trans*-$[MX(NO)L_4]^{2+}$ [M = Ru or Fe, X = Cl or Br, L = pyridine or *o*-phenylenebis(dimethylarsine)] with hydroxylamine, 1733–4
Magnetic circular dichroism spectra of tris-chelate complexes of 2,2′-bipyridyl and 1,10-phenanthroline with iron(II), ruthenium(II), and osmium(II) at 4.2 K, 1781–8
Organic chemistry of dinuclear metal centres. Part 8. Organo-iron–ruthenium chemistry. *X*-Ray structure of *trans*-$[FeRu(CO)_2(\mu\text{-}CO)_2(\eta\text{-}C_5H_5)_2]$, 1935–44
The peroxodisulphate–hexacyanoferrate(II) reaction. Reactivity and ionic association in isodielectric water–co-solvent mixtures, 1975–6
Reactivity of $[PPh_4][Fe_2(CO)_6\{\mu\text{-}CPhCPhC(CF_3)C(CF_3))H\}]$ toward electrophiles. *X*-Ray structure of $[Fe_2(CO)_6\text{-}\{\mu\text{-}CPhCPhC(CF_3)CHC(OEt)_2\}]$, a product resulting from fluorine abstraction at a CF_3 group and subsequent fluorine substitution by ethoxy-groups, 1981–4
Crystal structure and spectroscopic and redox properties of the iron–sulphur cluster compound $[NEt_4]_2[Fe_4S_4(SC_6H_4NH_2\text{-}4)_4]$, 2161–6
A mechanistic study of the reaction of iron(III) porphyrins with imidazoles. Hydrogen bonding by the propionic acid side chains in hemin chloride, 2269–76
Metallaborane chemistry. Part 14.[1] Icosahedral η^6-arene carbametallaboranes of iron and ruthenium; molecular structures of *closo*-[1-(η^6-C_6H_5Me)-2,4-Me_2-1,2,4-$FeC_2B_9H_9$] and *closo*-[3-(η^6-C_6H_6)-3,1,2-$RuC_2B_9H_{11}$], 2343–8
Crystal structures and magnetism of binuclear iron(III) complexes with a linear oxo-bridge, $[Fe_2O(bbimae)_2X_2][NO_3]_2$ {bbimae = 2-[bis(benzimidazol-2-ylmethyl)amino]ethanol, X = Cl or NCS}, 2375–80
Selective conversion of CO into methanol at ordinary temperature. Part 4. Activation by iron(II), iron(III), and chromium(III) complexes, 2499–504
Magnetic and spectroscopic properties of some heterotrinuclear basic acetates of chromium(III), iron(III), and divalent metal ions, 2509–20
The $[Fe_3(\mu_3\text{-}CR)(CO)_{10}]^-$ cluster anions as building blocks for the synthesis of mixed-metal clusters. Part 1. Synthesis of mixed clusters $[MFe_3(\mu_3\text{-}CMe)(CO)_{10}(PPh_3)]$ (M = Cu or Au) and crystal structure of $[CuFe_3(\mu_3\text{-}CMe)(CO)_{10}(PPh_3)]$, 2521–4
Studies on spin-equilibrium iron(III) complexes. Part 1. Syntheses and magnetic properties of a new family of spin cross-over iron(III) complexes with a unidentate ligand over a wide range of the spectrochemical series and a quinquedentate ligand derived from salicylaldehyde and di(3-aminopropyl)amine. *X*-Ray crystal structure of [4-azaheptamethylene-1,7-bis(salicylideneiminato)](4-methylpyridine)iron(III) tetraphenylborate, 2575–84
Conformation analysis of compounds of the type $[Fe(\eta^5\text{-}C_5H_5)(CO)(PPh_3)(CH_2R)]$ (R = alkyl or aryl): a comment, 2691
Conformation analysis of compounds of the type $[Fe(\eta^5\text{-}C_5H_5)(CO)(PPh_3)(CH_2R)]$ (R = alkyl or aryl): a reply, 2691

ISOCYANATE
Magnetic behaviour of tetrakis[(2-diethylaminoethanolato)isocyanatocopper(II)], a complex with an antiferromagnetic ground state; the crystal and molecular structure of the triclinic modification, 1909–14

ISOCYANIDE
Bimetallic systems. Part 8. Heterobimetallic di-isonitrile or isonitrile–carbonyl complexes of rhodium or iridium bridged to silver or gold by $Ph_2PCH_2PPh_2$, 511–6
Synthesis and structural characterisation of some *triangulo*-platinum clusters containing isocyanide ligands, 1693–8
Preparation and properties of *mer*-$[ReCl(N_2)(CNR)\{P(OMe)_3\}_3]$ (R = Me, Et, Bu^t, C_6H_4Me-4, or C_6H_4Cl-4) and $[ReCl(N_2)(CNMe)(PPh_3)\{P(OEt)_3\}_2]$. *X*-Ray crystal structure of *mer*-$[ReCl(N_2)(CNMe)\{P(OMe)_3\}_3]$ and reductive cleavage of the isocyanide ligands to primary amines upon protonation, 2079–84

ISOMERISM
Studies of phosphazenes. Part 22. High-field nuclear magnetic resonance investigation of novel isomeric oxophosphazadienes, 1431–4
Linkage isomerism in penta-ammineruthenium(II),(III) complexes of benzotriazole, 2469–72

ISOMERIZATION
The red form of bis(1,10-phenanthroline)dithiocyanatoiron(II), 991–6
Crystal structures and interrelationships of the blue and green conformational isomers of tetrakis(trimethylarsine sulphide)cobalt(II) perchlorate, 1289–94

ISOPOLYANION
Ion exchange of $K_4Nb_6O_{17}\cdot 3H_2O$, 2349–52

ISOTHIOCYANATE
Crystal and molecular structures and magnetic properties of four new exchange-coupled copper(II) complexes derived from different 3-*N*,*N*-dialkylamino-1-propanols and pseudohalogens, 1243–8

KETONE
Co-ordination chemistry of pyridyl and *N*-methylimidazolyl ketones. Synthetic and *X*-ray structural studies of copper(II), nickel(II), and dimethylgold(III) complexes, 981–6
From diarylruthenium complexes to *ortho*-metallated ketones: a mechanistic and crystal structure study, 1235–42

KINETICS
The kinetics of the displacement, by chloride, of heterocyclic nitrogen bases (am) from *trans*-$[PtL(am)Cl_2]$ (L = C_2H_4, CO, or PMe_3). The effect of steric hindrance in the leaving group on the *trans* effect of L, 27–30
The peroxodisulphate–iodide reaction. Reactivity and ionic association and solvation in isodielectric water–solvent mixtures, 31–4
Rate parameters for oxygen and carbon monoxide binding to a liposome-embedded heme under physiological conditions, 65–8
Reaction of bis-μ-diethylphosphido-bis(tetracarbonylmetal) (*M*–*M*) (M = Cr or W) with tri-n-butylphosphine: kinetics and mechanism of a reaction involving seven-co-ordinate complexes, 91–8
Studies of phosphazenes. Part 21. Associative and dissociative pathways in the aminolysis reactions of halogenocyclotriphosphazenes, 285–90
Hydrogen generation by hydrolysis of sodium tetrahydroborate: effects of acids and transition metals and their salts, 307–14
Light-induced electron-transfer reactions. Part 3. Kinetics of the oxidation reaction of oxalate ion by peroxodisulphate ion, induced by irradiation with visible light of an aqueous solution containing tris(2,2′-bipyridine)ruthenium(II), 355–60
Proton exchange and base hydrolysis of *syn,anti-cis*-dichloro(1,4,7,10-tetra-azacyclododecane)cobalt(III) cations, 475–8
Kinetic and structural investigations of $[Fe^{III}(edta)]^-$ [edta = ethylenediaminetetra-acetate(4−)] catalysed decomposition of hydrogen peroxide, 493–502
A kinetic study of the gas-phase thermolysis of hexaborane(10), 541–8
Kinetics and mechanism of single electron oxidations of the tervalent uranium ion, U^{3+}(aq), by free radicals in aqueous solutions, 641–4
Displacement of pyridine-2-methanol from dichloro(pyridine-2-methanolato)gold(III) in acidic solution. Ring opening at oxygen, 731–6
Electron-transfer reactions from *cis*-dialkylbis(2,2′-bipyridyl)cobalt(III) complexes to organic oxidants, 899–904
Mechanism of addition of aryltrimethyl-silanes and -stannanes to tricarbonyl(cyclohexadienyl)ruthenium(II) cation, 1049–52
Reactions of cobalt(II) protoporphyrin IX dimethyl ester, [Co^{II}P], and [Co^{III}P(Cl)] in co-ordinating aliphatic alcohols, 1095–102
Equilibrium and kinetic study of pyridine binding to phthalocyaninatoiron(II) in dimethyl sulphoxide, 1107–12
Equilibrium and kinetic study of the reaction between phthalocyaninatoiron(II) and carbon monoxide in dimethyl sulphoxide in the presence of pyridine. Evidence for the formation of a transient, 1113–8
Studies on singlet oxygen in aqueous solution. Part 1. Formation of singlet oxygen from hydrogen peroxide with two-electron oxidants, 1141–6

KINETICS (contd)
Studies on singlet oxygen in aqueous solution. Part 2. Water-soluble square-planar nickel complexes as quenchers, 1147–50
Studies on singlet oxygen in aqueous solution. Part 3. The decomposition of peroxy-acids, 1151–4
On the effect of cyanide ion on the reaction of pentacyanonitrosylferrate(2−) with cysteine, 1191–4
Kinetic studies of oxidative dealkylation of alkylcobalamins by hexachloroplatinate(IV), 1375–80
Kinetics and mechanism of the reductions of tris(oxalato)cobaltate(III) ion by ruthenium(II) species in aqueous solution, 1665–8
Oxidation of thiourea by iodate: a new type of oligo-oscillatory reaction, 1669–72
Kinetics and mechanism of the oxidation of substituted malonic acids by the hexachloroiridate(IV) ion, 1683–8
The kinetics and stoicheiometry of silver(III) reduction by the octacyano-complexes of molybdenum(IV) and tungsten(IV), 1789–94
The peroxodisulphate–hexacyanoferrate(II) reaction. Reactivity and ionic association in isodielectric water–co-solvent mixtures, 1975–6
The mechanisms of the reactions between hydrido-complexes and diazonium salts. Part 1. The reactions of *trans*-$[PtH(Cl)(PEt_3)_2]$, 2059–66
The mechanisms of the reactions between hydrido-complexes and diazonium salts. Part 2. The reactions of *cis,mer*-$[RhHCl_2(PEtPh_2)_3]$, 2067–78
The preparation of trichloro(phenyl 2-pyridylmethyl sulphoxide)gold(III) and the kinetics and equilibria of its formation and subsequent reactions, 2091–4
A mechanistic study of the reaction of iron(III) porphyrins with imidazoles. Hydrogen bonding by the propionic acid side chains in hemin chloride, 2269–76
Studies of phosphazenes. Part 21. Associative and dissociative pathways in the aminolysis reactions of halogenocyclotriphosphazenes (1985, 285), 2459–60
A kinetic study of the complexation of cysteine and related compounds with aqueous vanadium(II) and vanadium(III) at approximately neutral pH; the mediating role of sulphur compounds in electron transfer, 2461–8
Linkage isomerism in penta-ammineruthenium(II),(III) complexes of benzotriazole, 2469–72
Kinetics and mechanism of the equilibration reaction between (2,2′,2″-nitrilotriethoxy)nitrosylvanadate(1−) and cyanide. Crystal structures of sodium (2,2′,2″-nitrilotriethoxy)nitrosylvanadate(I)–sodium perchlorate tetrahydrate and of barium cyano(2,2′,2″-nitrilotriethoxy)nitrosylvanadate(I) pentahydrate, 2493–8
Studies on singlet oxygen in aqueous solution. Part 4. The 'spontaneous' and catalysed decomposition of hydrogen peroxide, 2525–30
Reactions of metal ions with triketones. Part 2. Kinetics and mechanisms of the reactions of nickel(II) and cobalt(II) with heptane-2,4,6-trione in methanol–water (70:30 v/v), 2565–70
Kinetics and mechanism of electron-transfer reactions of bismuth(V) in aqueous acidic perchlorate–fluoride media. Part 1. Oxidation of hypophosphorous acid, 2571–4
Kinetics and mechanisms of complex formation of gallium(III) and indium(III). The reactions with 4-(2-pyridylazo)resorcinol in water and other mixed solvents, 2615–22
Kinetics and mechanism of ortho-palladation of ring-substituted *NN*-dimethylbenzylamines, 2629–38

LABELLING
Steric changes at labelled NH_2 sites during the base-catalysed ammoniation of the *trans*-azido(dimethyl sulphoxide)bis(ethylenediamine)cobalt(III) ion, 1495–500
Oxygenation studies. Part 7. Catalytic dioxygenation of cyclo-octa-1,5-diene at a rhodium centre, 1591–6
Steric changes at labelled NH_2 sites during the base-catalysed ammoniation of the *trans*-azido(dimethyl sulphoxide)bis(ethylenediamine)cobalt(III) ion, 2223–4

LANTHANIDE
Mixed-ligand complexes of trivalent lanthanides. Part 3. Complexes of heptafluorodimethyloctane-3,5-dione and pyrazine: syntheses and spectral studies, 2547–50

LEAD
Subvalent group 4B metal alkyls and amides. Part 7. Transition-metal chemistry of metal(II) bis(trimethylsilyl)amides $M'(NR_2)_2$ (R = $SiMe_3$; M′ = Ge, Sn, or Pb), 51–8

LIQUID CRYSTAL N.M.R.
Determination of the molecular structure of difluorophosphine sulphide by the combined analysis of data from electron diffraction, microwave spectroscopy, and liquid crystal nuclear magnetic resonance spectroscopy, 755–60

LITHIO
Synthesis of α-lithioarylmethanes of *m*-xylene and its α-trimethylsilyl derivatives; crystal structure of $[\{Li(Me_2NCH_2CH_2NMe_2)\}_2\{C_6H_4(CHSiMe_3)_2\text{-}m\}]$, 337–44

LITHIUM
Synthesis of α-lithioarylmethanes of *m*-xylene and its α-trimethylsilyl derivatives; crystal structure of $[\{Li(Me_2NCH_2CH_2NMe_2)\}_2\{C_6H_4(CHSiMe_3)_2\text{-}m\}]$, 337–44

LUMINESCENCE
Glycine complexation with uranyl ion: absorptiometric, luminescence, and *X*-ray structural studies of tetrakis-(glycine)dioxouranium(VI) nitrate, 517–22

MACROCYCLE
Preparation of 11-(2′-dimethylaminoethyl)-1,4,7-trimethyl-1,4,7,11-tetra-azacyclotetradecane, and characterisation of its nickel(II), copper(II), and zinc(II) complexes, 219–22
The preparation and co-ordination chemistry of 2,2′:6′,2″-terpyridine macrocycles. Part 4. Structural characterisation of an intermediate in a transient template reaction, 333–6
Macrocyclic polyphosphane ligands. Iron(II), cobalt(II), and nickel(II) complexes of (4*RS*,7*RS*,13*SR*,16*SR*)-tetraphenyl-1,10-dipropyl-1,10-diaza-4,7,13,16-tetraphosphacyclo-octadecane: crystal structures of their tetraphenylborate derivatives, 479–86
Reaction of some aliphatic diamines with four-co-ordinated unsymmetrical ketoenamine copper(II)and nickel(II) complexes, 803–6
Reactions of cobalt(II) protoporphyrin IX dimethyl ester, $[Co^{II}P]$, and $[Co^{III}P(Cl)]$ in co-ordinating aliphatic alcohols, 1095–102
Solution chemistry of macrocycles. Part 4. Thermodynamics of protonation and complexation of several N_2S_2 macrocycles, 1169–72
Macrocyclic polyphosphane ligands. Cobalt(II) and nickel(II) complexes of 5*RS*,8*RS*,16*RS*,19*RS*-tetraphenyl-5,8,16,19-tetraphospha-1,12-dithiacyclododecosane (δ-L^1) and the crystal structure of $[Co(\delta\text{-}L^1)][BF_4]_2\cdot 0.5H_2O$, 1179–82
Studies of pendant-arm macrocyclic ligands. Part 4. Two penta-aza macrocycles based on 1-(2′-dimethylaminoethyl)-1,5,9,13-tetra-azacyclohexadecane and its complexes with bivalent metal ions, 1361–4
Macrocyclic polyphosphane ligands. Cobalt(II) and nickel(II) complexes of the γ and ε diastereoisomers of 4,7,13,16-tetraphenyl-4,7,13,16-tetraphospha-1,10-dithiacyclo-octadecane (L^1) and the crystal structure of $[Co(\varepsilon\text{-}L^1)][BPh_4]_2\cdot$ EtOH, 1425–30
Dismutation of superoxide ion in an aprotic solvent by 5,10,15,20-tetra-*p*-tolylporphyrinatocobalt(II), 1513–6
Mono- and di-nuclear rhodium and palladium complexes of macrocyclic ligands containing the 2,6-di(thiomethyl)pyridine sub-unit, 1517–22
Studies in the flexibility of macrocycle ligands. Calculation of macrocycle cavity size by force-field methods. Crystal and molecular structures of $[CoLCl][ClO_4]_2$ and $[CuL][PF_6]_2$ {L = 2,13-dimethyl-3,6,9,12,18-penta-azabicyclo[12.3.1]octadeca-1(18),14,16-triene}, 1829–34
The stability of nickel(II) complexes of tetra-aza macrocycles, 1877–80
Reactions of tris[(2*S*)-2-(aminomethyl)pyrrolidine]nickel(II) ion with alk-3-en-2-ones or 4-hydroxyalkan-2-ones: formation of an optically active tetra-aza macrocycle, 2139–44
Synthesis and cryptate complexes of azathia macropolycyclic ligands based on 12-membered N_2S_2 and 15-membered N_2S_3 macrocyclic subunits, 2311–8
Interactions of $[Mo(CO)_6]$ and $[Mo(CO)_3(C_6H_3CH_3)]$ with O_2N_2- and O_2N_3-donor macrocycles and the *X*-ray crystal structure of tetracarbonyl(6,7,16,17-tetrahydro-15*H*-dibenzo[*e*,*n*][1,4,8,12]dioxadiazacyclopentadecine)molybdenum–toluene (2/1), 2561–4
Complexes of lanthanoid salts with macrocyclic ligands. Part 17. Synthesis and crystal and molecular structure of a hydroxide-bridged praseodymium trifluoroacetate complex with 15-crown-5-ether, $[Pr_2(CF_3CO_2)_3(OH)(C_{10}H_{20}O_5)_2]\text{-}[Pr_2(CF_3CO_2)_6]$, 885–90

MAGNESIUM
Oxoalkyls of rhenium-(V) and -(VI). *X*-Ray crystal structures of

MAGNESIUM (contd)
$(Me_4ReO)_2Mg(thf)_4$, $[(Me_3SiCH_2)_4ReO]_2Mg(thf)_2$, $Re_2O_3Me_6$, and $Re_2O_3(CH_2SiMe_3)_6$, 2167–76
MAGNETISM
Crystal and molecular structures and magnetic properties of tetrameric copper(II) complexes with 3-hydroxy-5-hydroxymethyl-4-(4′-hydroxy-4′-phenyl-2′-azabut-1′-en-1′-yl)-2-methylpyridine (H_2L^3), $[Cu_4L^3_4]$·9CH_3OH and 3-hydroxy-5-hydroxymethyl-4-(4′-hydroxy-3′-methyl-4′-phenyl-2′-azabut-1′-en-1′-yl)-2-methylpyridine (H_2L^1), $[Cu_4L^1_4]$·8CH_3CH_2OH: two complexes with ferromagnetic ground states, 315–20
Spectroscopic and magnetic properties of cobalt(II) and nickel(II) clusters obtained from 1-(hydroxymethyl)-3,5-dimethylpyrazole. *X*-Ray structure of tetrakis[chloro(μ_3-3,5-dimethyl-*N*-oxymethylpyrazolato-N^2,μ_3-*O*)(ethanol)nickel(II)] 737–42
Crystal and molecular structures and magnetic properties of bromo(3-dimethylaminopropan-1-olato)copper(II), chloro(3-dimethylaminopropan-1-olato)copper(II), and (3-diethylaminopropan-1-olato)isocyanatocopper(II), 913–20
Crystal and molecular structures and magnetic properties of four new exchange-coupled copper(II) complexes derived from different 3-*N*,*N*-dialkylamino-1-propanols and pseudohalogens, 1243–8
Magnetic behaviour of tetrakis[(2-diethylaminoethanolato)isocyanatocopper(II)], a complex with an antiferromagnetic ground state; the crystal and molecular structure of the triclinic modification, 1909–14
Crystal structures and magnetic properties of binuclear five-co-ordinate copper(II) complexes with a phenolate bridge and their catalytic functions in multielectron redox reactions, 1945–52
X-Ray crystal structures and magnetic properties of azide-bridged binuclear copper(II) complexes containing the Schiff-base ligand derived from 2-pyridinecarbaldehyde and histamine. Structure–magnetism relationship, 2095–100
Electrical and magnetic properties of $K_3Cu_8S_6$, 2319–20
Crystal structures and magnetism of binuclear iron(III) complexes with a linear oxo-bridge, $[Fe_2O(bbimae)_2X_2][NO_3]_2$ {bbimae = 2-[bis(benzimidazol-2-ylmethyl)amino]ethanol, X = Cl or NCS}, 2375–80
Magnetic and spectroscopic properties of some heterotrinuclear basic acetates of chromium(III), iron(III), and divalent metal ions, 2509–20
Studies on spin-equilibrium iron(III) complexes. Part 1. Syntheses and magnetic properties of a new family of spin cross-over iron(III) complexes with a unidentate ligand over a wide range of the spectrochemical series and a quinquedentate ligand derived from salicylaldehyde and di(3-aminopropyl)amine. *X*-Ray crystal structure of [4-azaheptamethylene-1,7-bis(salicylideneiminato)](4-methylpyridine)iron(III) tetraphenylborate, 2575–84
Crystal and molecular structure and magnetic properties of linear trimeric copper(II) complexes with predominant ferromagnetic exchange interaction, 2609–14
MALONIC ACID
Kinetics and mechanism of the oxidation of substituted malonic acids by the hexachloroiridate(IV) ion, 1683–8
MALONITRILE
The chemistry of vitamin B_{12}. Part 26. Co-ordination of the malonitrile anion by Co^{III} corrinoids: first experimental determination of equilibrium constants for the co-ordination of a tetrahedral carbanion by a transition-metal ion, 1619–22
MANGANESE
Binuclear metal complexes. Part 55. Dimanganese(II) and mixed-valence manganese(II,III) complexes of 'strati-bis' Schiff bases with two salen-like donating sites, 59–64
Electron spin resonance spectra of manganese(II) ions in the double-stranded chain polymers $CdLX_2$ (L = pyridine or methylpyridine; X = Cl or Br), 75–80
The co-ordination chemistry of manganese. Part 14. Synthesis of manganese(II) complexes of tertiary phosphine ligands containing 2-cyanoethyl groups, 135–40
The template synthesis and crystal and molecular structure of a seven-co-ordinate manganese(II) complex with 2,6-diacetylpyridine mono(2-aminobenzoylhydrazone), 215–8
Photochemistry of manganese porphyrins. Part 9. Redox reactions photosensitised by diamagnetic metalloporphyrins, 503–10
Difluorophosphate complexes of chromium, manganese, iron, cobalt, and nickel, 707–10
The addition of protic acids to $[Mn_2(CO)_8(Ph_2PCH_2PPh_2)_2]$ to give bridging hydrido-compounds, 743–8
Alkyl, hydrido, and tetrahydroaluminato complexes of manganese with 1,2-bis(dimethylphosphino)ethane (dmpe). *X*-Ray crystal structures of $Mn_2(\mu\text{-}C_6H_{11})_2(C_6H_{11})_2(\mu\text{-dmpe})$, $(dmpe)_2Mn(\mu\text{-}H)_2AlH(\mu\text{-}H)_2AlH(\mu\text{-}H)_2Mn(dmpe)_2$, and $Li_4\{MnH(C_2H_4)[CH_2(Me)PCH_2CH_2PMe_2]_2\}_2$·2$Et_2O$, 921–30
Reduction–oxidation properties of organotransition-metal complexes. Part 21. Synthesis and *X*-ray structural characterisation of the redox-related pair of cyclohexadienyl complexes $[Mn(CO)(dppe)(\eta^5\text{-}C_6H_6Ph)]$ and $[Mn(CO)(dppe)(\eta^5\text{-}C_6H_6Ph)][PF_6]$·0.5$CH_2Cl_2$, 1019–26
Water photolysis. Part 1. The photolysis of co-ordinated water in $[\{MnL(H_2O)\}_2][ClO_4]_2$ (L = dianion of tetradentate O_2N_2-donor Schiff bases). A model for the manganese site in photosystem II of green plant photosynthesis, 1391–8
Potentiometric study of the complex-formation equilibria of manganese(II), cobalt(II), nickel(II), copper(II), and zinc(II) with ethylenediamine-*N*-acetic acid, 1605–8
Addition of proton and boron trifluoride at terminal and bridging cyano-ligands in dinuclear manganese compounds, 1609–12
Sulphur-containing metal complexes. Part 14. Reactions of carbene anions with carbon disulphide or carbon diselenide, 1963–6
Bimetallic systems. Part 13. Platinum–manganese carbonyl complexes containing bridging $Ph_2PCH_2PPh_2$(dppm) ligands: crystal structure of $[(OC)_3Mn(\mu\text{-dppm})_2PtH(Br)]BF_4$, 2131–8
On the synthesis of dimethylphenylphosphine complexes of manganese(II) halides, 2203–4
Investigation into aroylhydrazones as chelating agents. Part 7. Synthesis and spectroscopic characterization of complexes of Mn^{II}, Co^{II}, Ni^{II}, Cu^{II}, and Zn^{II} with 2,6-diacetylpyridine bis(2-aminobenzoylhydrazone) and *X*-ray structure of chloro[2,6-diacetylpyridine bis(2-aminobenzoylhydrazone)](methanol)manganese(II) chloride monohydrate, 2387–92
The co-ordination of small molecules by manganese(II) phosphine complexes. Part 3. The dependence on the nature of the halogen in $[MnX_2(PR_3)]$ (X = Cl, Br, or I; R_3 = $PhMe_2$, $PhEt_2$, $PhPr^n_2$, $PhBu^n_2$, $PhBu^t_2$, Pr^n_3, Bu^n_3, or Ph_3) on adduct formation with sulphur dioxide in the solid state and in tetrahydrofuran solution, 2661–4
He I and He II photoelectron spectra of open-chain pentadienyl complexes of manganese and rhenium, 2677–82
MASS SPECTROMETRY
A kinetic study of the gas-phase thermolysis of hexaborane(10), 541–8
A photoelectron spectroscopic study of di-t-butylphosphazene, 879–84
Mass spectrometric studies on cyclo- and poly-phosphazenes. Part 2. Oligomerization of hexa(aryloxy)cyclotriphosphazatrienes, 1547–54
MATRIX
Characterisation of chromium(VI) oxide tetrafluoride, $CrOF_4$, and caesium pentafluoro-oxochromate(VI) $Cs[CrOF_5]$, 529–34
Photochemistry of dicarbonyl(η^5-cyclopentadienyl)-methyl- and -ethyl-iron and -ruthenium complexes in solutions at −30 °C and in frozen gas matrices at 12 K, 1365–74
Spectroscopic studies on the higher binary fluorides of chromium: CrF_4, CrF_5, and CrF_6, both in the solid state and isolated in inert gas matrices, 1443–50
M.C.D.
Magnetic circular dichroism spectra of tris-chelate complexes of 2,2′-bipyridyl and 1,10-phenanthroline with iron(II), ruthenium(II), and osmium(II) at 4.2 K, 1781–8
MECHANISM
Reaction of bis-μ-diethylphosphido-bis(tetracarbonylmetal) (*M*–*M*) (M = Cr or W) with tri-n-butylphosphine: kinetics and mechanism of a reaction involving seven-co-ordinate complexes, 91–8
Studies of phosphazenes. Part 21. Associative and dissociative pathways in the aminolysis reactions of halogenocyclotriphosphazenes, 285–90
Ligand substitution on (*N*,*N*-dimethylformamide)[2,2′,2″-tri(*N*,*N*-dimethylamino)triethylamine]cobalt(II), 413–6
Proton exchange and base hydrolysis of *syn,anti-cis*-dichloro(1,4,7,10-tetra-azacyclododecane)cobalt(III) cations, 475–8
Complexes of the platinum metals. Part 24. The role of dioxygen in the reactions of trifluoroacetic acid with the rhodium and iridium nitrosyls $[M(NO)(PPh_3)_3]$, 617–20
Kinetics and mechanism of single electron oxidations of the

MECHANISM (contd)
tervalent uranium ion, U^{3+}(aq), by free radicals in aqueous solutions, 641–4
Solvent dependence of the stereochemistry of base-catalysed solvolysis of *trans*-$[Co(NH_3)_4(^{15}NH_3)X]^{3+/2+}$ ions in dipolar aprotic solvents, 659–62
Displacement of pyridine-2-methanol from dichloro(pyridine-2-methanolato)gold(III) in acidic solution. Ring opening at oxygen, 731–6
Mechanism of addition of aryltrimethyl-silanes and -stannanes to tricarbonyl(cyclohexadienyl)ruthenium(II) cation, 1049–52
Reactions of cobalt(II) protoporphyrin IX dimethyl ester, $[Co^{II}P]$, and $[Co^{III}P(Cl)]$ in co-ordinating aliphatic alcohols, 1095–102
A mechanistic study on complexes of type *mer*-$[Cr(CO)_3(\eta^2\text{-}L\text{–}L)(\sigma\text{-}L\text{–}L)]$ (where L–L = $Ph_2PCH_2PPh_2$, $Ph_2PNHPPh_2$, or $Ph_2PNMePPh_2$) using spectroscopic and convolutive electrochemical techniques, 1213–22
From diarylruthenium complexes to *ortho*-metallated ketones: a mechanistic and crystal structure study, 1235–42
Electron-transfer reactions in nitrogen fixation. Part 1. The electrosynthesis of dinitrogen, hydride, isocyanide, and carbonyl complexes of molybdenum: intermediates, mechanisms, and energetics, 1255–64
Kinetic studies of oxidative dealkylation of alkylcobalamins by hexachloroplatinate(IV), 1375–80
Steric changes at labelled NH_2 sites during the base-catalysed ammoniation of the *trans*-azido(dimethyl sulphoxide)bis(ethylenediamine)cobalt(III) ion, 1495–500
The chemistry of vitamin B_{12}. Part 25. Mechanism of the β-elimination of olefins from alkylcorrinoids; evidence for an initial homolytic fission of the Co–C bond, 1613–8
Kinetics and mechanism of the reductions of tris(oxalato)cobaltate(III) ion by ruthenium(II) species in aqueous solution, 1665–8
Kinetics and mechanism of the oxidation of substituted malonic acids by the hexachloroiridate(IV) ion, 1683–8
The mechanisms of the reactions between hydrido-complexes and diazonium salts. Part 1. The reactions of *trans*-$[PtH(Cl)(PEt_3)_2]$, 2059–66
The mechanisms of the reactions between hydrido-complexes and diazonium salts. Part 2. The reactions of *cis,mer*-$[RhHCl_2(PEtPh_2)_3]$, 2067–78
Steric changes at labelled NH_2 sites during the base-catalysed ammoniation of the *trans*-azido(dimethyl sulphoxide)bis(ethylenediamine)cobalt(III) ion, 2223–4
Preparation and characterisation of 2,2′-bipyridine-4,4′-disulphonic and -5-sulphonic acids and their ruthenium(II) complexes. Excited-state properties and excited-state electron-transfer reactions of ruthenium(II) complexes containing 2,2′-bipyridine-4,4′-disulphonic acid or 2,2′-bipyridine-4,4′-dicarboxylic acid, 2247–62
A mechanistic study of the reaction of iron(III) porphyrins with imidazoles. Hydrogen bonding by the propionic acid side chains in hemin chloride, 2269–76
Studies of phosphazenes. Part 21. Associative and dissociative pathways in the aminolysis reactions of halogenocyclotriphosphazenes (1985, 285), 2459–60
Reactions of metal ions with triketones. Part 2. Kinetics and mechanisms of the reactions of nickel(II) and cobalt(II) with heptane-2,4,6-trione in methanol–water (70:30 v/v), 2565–70
Kinetics and mechanism of electron-transfer reactions of bismuth(V) in aqueous acidic perchlorate–fluoride media. Part 1. Oxidation of hypophosphorous acid, 2571–4
Kinetics and mechanisms of complex formation of gallium(III) and indium(III). The reactions with 4-(2-pyridylazo)resorcinol in water and other mixed solvents, 2615–22
Kinetics and mechanism of ortho-palladation of ring-substituted *NN*-dimethylbenzylamines, 2629–38

MERCURY
Synthesis, solution structure, and halide addition compounds of bis(dicyclohexylphosphinothioito)mercury(II), 223–4
The structure of pentachlorocyclopentadienylphenylmercury(II) in the solid state and in solution; the effect of *X*-ray absorption and crystal decay on molecular geometry parameters, 471–4
Spectrochemistry of solutions. Part 16. A Raman spectroscopic study of the complexation of mercury(II) by cyanide ligands in liquid ammonia at 293 K, 663–8
Mercurated and tellurated Schiff bases and phenylhydrazones, 821–6
Preparation and crystal structure of $Na_4[Hg_4(\mu\text{-}Cl)_4\{P(O)(OEt)_2\}_8]\cdot 6H_2O$, 1061–4
Thermochemical data for adducts of zinc, cadmium, and mercury halides with hexamethylphosphoramide, 1103–6
Structure and solvation of mercury(II) iodide, bromide, and chloride in pyridine solution; refinement of the crystal structure of di-iodobis(pyridine)mercury(II), $[HgI_2(py)_2]$, 1597–604
Crystal structure of tris(methylmercurio)sulphonium perchlorate, 2457–8

MERCURY-199
Synthesis, solution structure, and halide addition compounds of bis(dicyclohexylphosphinothioito)mercury(II), 223–4

METALLATION
Metallation of 2-ethenylpyridine at triosmium clusters: *X*-ray crystal structures of the open trinuclear clusters $[Os_3H(CO)_9L(NC_5H_4CH{=}CH)]$ (L = CO or PMe_2Ph), 85–90

METALLEPINE
Axially asymmetric metal alkyls. Part 3. Chemical, electrochemical, and structural studies of group 5A $d^{0,1,2}$ metallepines $[M\{(2\text{-}CH_2C_6H_4)_2\}(\eta\text{-}C_5H_5)_2]^z$ (M = Nb or Ta; $z = 1-$, 0, or 1+); synthesis of $[Nb\{2\text{-}CH_2(C_6H_4)_2Me\text{-}2'\}X(\eta\text{-}C_5H_5)_2]$ (X = Cl or CO) and $[\{(\eta\text{-}C_5H_5)_2ClV\}_2\{(2\text{-}CH_2C_6H_4)_2\}]$, and electrochemistry of $[M(CH_2Ph)_{2-n}Cl_n(\eta\text{-}C_5H_5)_2]$ (n = 0, M = Nb, or Ta; n = 1, M = V), 1747–60

METHANOL
Selective conversion of CO into methanol at ordinary temperature. Part 4. Activation by iron(II), iron(III), and chromium(III) complexes, 2499–504

METHYL
Determination of the molecular structures and conformations of methylbis(methylsilyl)amine and bis(dimethylsilyl)methylamine in the gas phase by electron diffraction, 191–8
The preparation and co-ordination chemistry of 2,2′:6′,2″-terpyridine macrocycles. Part 4. Structural characterisation of an intermediate in a transient template reaction, 333–6
Static and dynamic nuclear magnetic resonance studies of complexes of trimethylplatinum(IV) halides with olefinic thio-and seleno-ethers. *X*-Ray crystal structures of $[PtXMe_3(MeSeCH{=}CHSeMe)]$ (X = Cl or I), 345–54
X-Ray crystal structures and properties of tris(tetrathiafulvalenium) tetrachlorodimethylstannate(IV) and tetrathiafulvalenium trichlorodimethylstannate (IV), 465–70
Mononuclear η^6-*p*-cymeneosmium(II) complexes and their reactions with Al_2Me_6 and other methylating reagents, 573–8
Isolated CH stretching frequencies, methyl group geometry, and methyl CH bond lengths and strengths in tricarbonyl(η^5-cyclopentadienyl)methylchromium(II), -molybdenum(II), and -tungsten(II), 1207–12
The crystal structure of diaquadichlorodimethyltin(IV)–purine (1/4), 1271–4
Synthesis and properties of the divalent 1,2-bis(dimethylphosphino)ethane (dmpe) complexes $MCl_2(dmpe)_2$ and $MMe_2(dmpe)_2$ (M = Ti, V, Cr, Mn, or Fe). *X*-Ray crystal structures of $MCl_2(dmpe)_2$ (M = Ti, V, or Cr), $MnBr_2(dmpe)_2$, $TiMe_{1.3}Cl_{0.7}(dmpe)_2$, and $CrMe_2(dmpe)_2$, 1339–48
Photochemistry of dicarbonyl(η^5-cyclopentadienyl)-methyl- and -ethyl-iron and -ruthenium complexes in solutions at −30 °C and in frozen gas matrices at 12 K, 1365–74
Bimetallic systems. Part 11. Heterobimetallic and unsymmetrical diplatinum complexes from *cis*-$[PtR_2(dppm\text{-}P)_2]$ (dppm = $Ph_2PCH_2PPh_2$; R = Me, 1-naphthyl, or C_6H_4Me-*o*): crystal structure of $[(C_6H_4Me\text{-}o)_2Pt(\mu\text{-}dppm)_2PtMe_2]$, 1677–82
Complexes of organoaluminium compounds. Part 13. Preparation and nuclear magnetic resonance spectra of the arylamido-compounds $AlMe_2(NHR')$ (R′ = Ph, C_6H_4Me-*o*, C_6H_4Me-*p*, $C_6H_3Me_2$-2,6) and the imido-compounds $AlMe(NR')$. Crystal and molecular structures of $[\{AlMe_2(NHC_6H_4Me\text{-}o)\}_2]$ and $[\{AlMe(NPh)\}_6]$, 1929–34
Trimethylphosphine as a reactive solvent: synthesis, crystal structures, and reactions of $[Ta(PMe_3)_3(\eta^2\text{-}CH_2PMe_2)(\eta^2\text{-}CHPMe_2)]$ and $[W(PMe_3)_4(\eta^2\text{-}CH_2PMe_2)H]$ and related studies, 2025–36
Oxoalkyls of rhenium-(v) and -(vi). *X*-Ray crystal structures of $(Me_4ReO)_2Mg(thf)_4$, $[(Me_3SiCH_2)_4ReO]_2Mg(thf)_2$, $Re_2O_3Me_6$, and $Re_2O_3(CH_2SiMe_3)_6$, 2167–76
Methyltin hydrides in fluorosulphuric acid, 2671–6
A novel pentameric hydrolysis product of $SnMe_2Cl_2$: crystal and molecular structure of $[NHEt_3][(SnMe_2Cl)_5O_3]$, 2683–6

METHYLATION
Mononuclear η^6-*p*-cymeneosmium(II) complexes and their reactions with Al_2Me_6 and other methylating reagents, 573–8
METHYLENE
Chemistry of the unsaturated cluster compound $[Os_3Pt(\mu\text{-}H)_2(CO)_{10}\{P(cyclo\text{-}C_6H_{11})_3\}]$; *X*-ray crystal structures of $[Os_3Pt(\mu\text{-}H)_2(CO)_{11}\{P(cyclo\text{-}C_6H_{11})_3\}]$, $[Os_3Pt(\mu\text{-}H)_4(CO)_{10}\{P(cyclo\text{-}C_6H_{11})_3\}]$, and $[Os_3Pt(\mu\text{-}H)_2(\mu\text{-}CH_2)(CO)_{10}\{P(cyclo\text{-}C_6H_{11})_3\}]$ (two isomers), 177–90
The syntheses and characterisation of, and the determination of $^1J(^{103}Rh\text{–}^{103}Rh)$ in $[(\eta^5\text{-}C_5Me_5)_2Rh_2(\mu\text{-}CH_2)_2\{\mu\text{-}CH_2CR(CH_2CR{=}CH_2)CH_2\}]$ (R = H or Me) and $[(\eta^5\text{-}C_5Me_5)_2Rh_2(\mu\text{-}CH_2)_2\{\mu\text{-}Ph_2P(CH_2)_nPPh_2\}]^{2+}$ (n = 1 or 2), 1555–60
METHYLENEAMINO
The reaction of $[Os_3(\mu\text{-}H)_2(CO)_9L]$ (L = CO or PEt_3) with dimethylcyanamide, Me_2NCN: *X*-Ray crystal structure of $[Os_3(\mu\text{-}H)(\mu\text{-}NCHNMe_2)(CO)_{10}]$ and the reactions of this complex with acids, 1355–60
METHYLENEDI-IMINE
Preparation and crystal structure of $[AsPh_4]_2[(WCl_5)_2\{\mu\text{-}NC(CF_3)_2N\}]$, 2205–8
METHYLMETHYLIDYNE
Chemistry of di- and tri-metal complexes with bridging carbene or carbyne ligands. Part 33. Reactions of $[W(\equiv CMe)(CO)_2(\eta\text{-}C_5H_5)]$ with the dimetal compounds $[MRh(\mu\text{-}CO)_2(\eta\text{-}C_5Me_5)_2]$ (M = Co or Rh); *X*-ray crystal structure of $[Rh_2W(\mu\text{-}CO)(\mu_3\text{-}CMe)(CO)_2(\eta\text{-}C_5H_5)(\eta\text{-}C_5Me_5)_2]$, 1315–22
METHYLPYRIDINE
Lewis-base adducts of Group 1B metal(I) compounds. Part 11. Synthesis and crystal structure of adducts of silver(I) bromide with monomethyl-substituted pyridine bases, 111–6
Lewis-base adducts of Group 1B metal(I) compounds. Part 12. Structural studies of some bis(methyl-substituted pyridine)-copper(I) and -silver(I) nitrates and perchlorates, 117–24
METHYLSILYL
Determination of the molecular structures and conformations of methylbis(methylsilyl)amine and bis(dimethylsilyl)methylamine in the gas phase by electron diffraction, 191–8
MICROWAVE
Determination of the molecular structure of difluorophosphine sulphide by the combined analysis of data from electron diffraction, microwave spectroscopy, and liquid crystal nuclear magnetic resonance spectroscopy, 755–60
MIXED VALENCE
Binuclear metal complexes. Part 55. Dimanganese(II) and mixed-valence manganese(II,III) complexes of 'strati-bis' Schiff bases with two salen-like donating sites, 59–64
Sulphur ligand–metal complexes. Part 16. Copper complexes of thioethers and the single-crystal *X*-ray structure of the polymeric mixed-valence complex, penta-μ-chloro-tris-μ-tetrahydrothiophene-tetracopper(I,II), 151–8
Molecular structure and solid-state properties of the two-dimensional conducting mixed-valence complex $[NBu_4]_{0.29}[Ni(dmit)_2]$ and the neutral $[Ni(dmit)_2]$ (H_2dmit = 4,5-dimercapto-1,3-dithiole-2-thione); members of an electron-transfer series, 783–94
Palladium(II,IV) mixed-valence complexes of 1,2-diaminoethane, 1,3-diaminopropane, and diethylenetriamine: syntheses, electronic, infrared, Raman, and resonance Raman spectra and *X*-ray studies, 815–20
Synthesis, spectroscopy, and structure of the mixed-valence complexes $[Pt^{II}(en)_2][Pt^{IV}(en)_2X_2][Pt^{III}_2(H_2P_2O_5)_4X_2]$ (en = 1,2-diaminoethane, X = Br or I), 579–86
MNDO
Structure and relative energies of some nitrogen-containing radical cations by MNDO calculation, 1915–22
MOLYBDENUM
Hexakis(trimethylphosphine)molybdenum chemistry: dinitrogen, ethylene, butadiene, η-cyclopentadienyl, and related derivatives, 423–34
Reactions of co-ordinated ligands. Part 33. Mononuclear η^2-vinyl complexes: synthesis, structure, and reactivity, 435–50
Tin–molybdenum oxides. A study by *X*-ray diffraction, Mössbauer spectroscopy, and electron spin resonance, 451–4
Catalytic air oxidation of benzoin in the presence of dioxomolybdenum(VI) complexes with sulphur chelate ligands, 635–40
Studies on transition-metal cyano-complexes. Part 4. Cyanide hydride complexes of Groups 6A and 8, 717–22
Bimetallic systems. Part 9. The synthesis of and nuclear magnetic resonance studies on 10-membered ring complexes of type $[(OC)_4M^1(\mu\text{-}Ph_2PCH_2CH_2PPh_2)_2M^2(CO)_4]$ (M^1, M^2 = Cr, Mo, or W), 1009–14
Tetrachlorobis(tetrahydrofuran)molybdate(III), a new intermediate for the synthesis of molybdenum(III) complexes: preparation and crystal structure, 1069–72
Isolated CH stretching frequencies, methyl group geometry, and methyl CH bond lengths and strengths in tricarbonyl(η^5-cyclopentadienyl)methylchromium(II), -molybdenum(II), and -tungsten(II), 1207–12
Electrochemical and chemical properties of di-iodonitrosyl[tris(3,5-dimethylpyrazolyl)borato]molybdenum, $[Mo\{HB(Me_2pz)_3\}(NO)I_2]$, and related complexes, 1249–54
Electron-transfer reactions in nitrogen fixation. Part 1. The electrosynthesis of dinitrogen, hydride, isocyanide, and carbonyl complexes of molybdenum: intermediates, mechanisms, and energetics, 1255–64
Preparation and properties of *cis*-$[W(N_2)_2(Ph_2PCH{=}PPh_2)_2]$ and *trans*-$[M(N_2)_2(Ph_2PCH{=}CHPPh_2)_2]$ (M = Mo or W), and the crystal structure of *cis*-$[W(N_2)_2(PMe_2Ph)_4]$, 1523–6
Pyramidal inversions and 1,2-metal shifts in pentacarbonyl-chromium, -molybdenum, and -tungsten derivatives of dialkyl disulphides and dialkyl diselenides. A nuclear magnetic resonance investigation, 1561–8
Dynamic nuclear magnetic resonance study of Group 6 metal pentacarbonyl complexes. Pyramidal inversion and 1,2-metal shifts in the complexes $[M(CO)_5(Me_2\overline{CCH_2EECH_2})]$ (M = Cr, Mo, or W; E = S or Se), 1569–76
Studies on cyclic bis(η^5:σ-2-cyclopentadienylidene-ethyl)- and bis(η^5:σ-4-cyclopentadienylidenebutyl)-molybdenum compounds, 1585–90
Formation of oxidized molybdenum-bearing ferrites, $Fe_{2.95-x}Mo_xO_4$ (x = 0.03—0.30) in aqueous suspensions by air oxidation and valence state of molybdenum ions in the lattice, 1713–6
The kinetics and stoicheiometry of silver(III) reduction by the octacyano-complexes of molybdenum(IV) and tungsten(IV), 1789–94
Reactions of hydride reagents with alkylmolybdenum carbonyl complexes. Reaction of $LiBHEt_3$ with $[MoMe(CO)_3(\eta\text{-}C_5H_5)]$, formation of an anionic acetaldehyde complex, and a stoicheiometric cycle for the synthesis of acetaldehyde, 1815–20
Chemistry of di- and tri-metal complexes with bridging carbene or carbyne ligands. Part 36. Reactions of the dimetal compounds $[ReM(\equiv CC_6H_4Me\text{-}4)(CO)_9]$ (M = Cr, Mo, or W) with octacarbonyldicobalt; crystal structures of $[Co_2WRe(\mu_3\text{-}CC_6H_4Me\text{-}4)(CO)_{15}]$ and $[Co_2Re(\mu_3\text{-}CC_6H_4Me\text{-}4)(CO)_{10}]$, 2001–8
Trimethylphosphine as a reactive solvent: synthesis, crystal structures, and reactions of $[Ta(PMe_3)_3(\eta^2\text{-}CH_2PMe_2)(\eta^2\text{-}CHPMe_2)]$ and $[W(PMe_3)_4(\eta^2\text{-}CH_2PMe_2)H]$ and related studies, 2025–36
Synthesis, structure, and bonding of fulvene complexes of titanium, molybdenum, and tungsten, 2037–50
Reactivity of $[NBu_4]_2[Mo_2Br_6]$ with several uni- and poly-dentate phosphines. *X*-Ray structure of $[NBu_4][MoBr_4(Ph_2PCH_2CH_2PPh_2)]$, 2263–8
Nitrogen-15 nuclear magnetic resonance relaxation mechanisms in dinitrogen complexes of molybdenum, tungsten, rhenium, and osmium, 2473–8
Interactions of $[Mo(CO)_6]$ and $[Mo(CO)_3(C_6H_5CH_3)]$ with O_2N_2- and O_2N_3-donor macrocycles and the *X*-ray crystal structure of tetracarbonyl(6,7,16,17-tetrahydro-15*H*-dibenzo[*e,n*][1,4,8,12]dioxadiazacyclopentadecine)molybdenum–toluene (2/1), 2561–4
Photochemical studies of the alkylammonium molybdates. Part 7. Octahedral sites for multi-electron reduction of $[Mo_8O_{26}(MoO_4)_2]^{8-}$, 2585–90
Complexes of molybdenum-(II) and -(IV) and tungsten(II) with sterically hindered thiolate ligands. Synthesis, reactivity, and *X*-ray crystal structures of $[PPh_4][Mo(SC_6H_2Pr^i_3\text{-}2,4,6)_3(CO)_2]$ and $[Mo(NNPh)(SC_6H_2Pr^i_3\text{-}2,4,6)_3(NCMe)]$, 2639–46
Molybdenum, rhenium, and tungsten complexes with bi- and tri-dentate phosphinothiolato-ligands; structures of $[Mo\{PhP(CH_2CH_2S)_2\}_2]$ and $[Mo(NNMe_2)\{PhP(CH_2CH_2S)_2\}_2]$, 2647–54

MORPHOLINIO
Reaction of bis(morpholinothiocarbonyl) disulphide with iodine. Existence of a 1:1 charge-transfer precursory adduct in an oxidation reaction. Isolation and crystal structure of bis[3,5-di(*N*-morpholinio)-1,2,4-trithiolane] hexadecaiodide, 1349–54
MORPHOLINOTHIOCARBONYL
Reaction of bis(morpholinothiocarbonyl) disulphide with iodine. Existence of a 1:1 charge-transfer precursory adduct in an oxidation reaction. Isolation and crystal structure of bis[3,5-di(*N*-morpholinio)-1,2,4-trithiolane] hexadecaiodide, 1349–54
MÖSSBAUER
Tin-119 Mössbauer and nuclear magnetic resonance studies of organotin compounds. Part 1. Sterically crowded tetraorganotin derivatives, 169–76
Tin–molybdenum oxides. A study by *X*-ray diffraction, Mössbauer spectroscopy, and electron spin resonance, 451–4
The red form of bis(1,10-phenanthroline)dithiocyanatoiron(II), 991–6
A tin-119 Mössbauer study of tin(II) fluoride, 1275–6
Mössbauer study of the *cis–trans* isomers of tin(IV) complexes. Some considerations about the sign of the electric-field gradient, 1281–4
Organotin biocides. Part 2. Variable-temperature ^{119}Sn Mössbauer study of phenyl-and cyclohexyl-tin compounds, 1417–24
Magnetic exchange interactions in perovskite solid solutions. Part 5. The unusual defect structure of $SrFeO_{3-y}$, 1455–70
Iodine-127 Mössbauer spectroscopy of copper(I) iodide–phosphine adducts, 1727–30
Methyltin hydrides in fluorosulphuric acid, 2671–6
NEPTUNIUM
Actinide structural studies. Part 7. The crystal and molecular structures of (2,2′-bipyridyl)dinitratodioxo-uranium(VI) and -neptunium(VI), and diacetato(2,2′-bipyridyl)dioxo-uranium(VI) and -neptunium(VI), 1001–8
NICKEL
Enantioselectivity of nickel(II) and copper(II) complexes of Schiff bases derived from amino acids and (*S*)-*o*-[(*N*-benzylprolyl)amino]acetophenone or (*S*)-*o*-[(*N*-benzylprolyl)amino]benzaldehyde. Crystal and molecular structures of [Ni{(*S*)-bap-(*S*)-Val}] and [Cu{(*S*)-bap-(*S*)-Val}], 17–26
Preparation of 11-(2′-dimethylaminoethyl)-1,4,7-trimethyl-1,4,7,11-tetra-azacyclotetradecane, and characterisation of its nickel(II), copper(II), and zinc(II) complexes, 219–22
Chlorine trioxide, Cl_2O_6, a most efficient perchlorating reagent in new syntheses of anhydrous metal perchlorates, chloryl and nitryl perchloratometalates of cobalt(II), nickel(II), and copper(II). Reactivity of chlorine trioxide with anhydrous or hydrated chlorides and nitrates, 297–306
Comparison between the bis(2-thiopyridine *N*-oxide) derivatives of Cu^{II} and Ni^{I}: an electron spin resonance study, 379–82
Macrocyclic polyphosphane ligands. Iron(II), cobalt(II), and nickel(II) complexes of (4*RS*,7*RS*,13*SR*,16*SR*)-tetraphenyl-1,10-dipropyl-1,10-diaza-4,7,13,16-tetraphosphacyclo-octadecane: crystal structures of their tetraphenylborate derivatives, 479–86
Difluorophosphate complexes of chromium, manganese, iron, cobalt, and nickel, 707–10
Spectroscopic and magnetic properties of cobalt(II) and nickel(II) clusters obtained from 1-(hydroxymethyl)-3,5-dimethylpyrazole. *X*-Ray structure of tetrakis[chloro(μ_3-3,5-dimethyl-*N*-oxymethylpyrazolato-N^2,μ_3-*O*)(ethanol)nickel(II)] 737–42
Molecular structure and solid-state properties of the two-dimensional conducting mixed-valence complex $[NBu_4]_{0.29}[Ni(dmit)_2]$ and the neutral $[Ni(dmit)_2]$ (H_2dmit = 4,5-dimercapto-1,3-dithiole-2-thione); members of an electron-transfer series, 783–94
Reaction of some aliphatic diamines with four-co-ordinated unsymmetrical ketoenamine copper(II)and nickel(II) complexes, 803–6
Reactions of 6,6′-bis(*nido*-decaboranyl) oxide and 6-hydroxy-*nido*-decaborane with dihalogenobis(phosphine) complexes of nickel, palladium, and platinum, and some related chemistry; nuclear magnetic resonance investigations and the crystal and molecular structures of bis(dimethylphosphine)-di-μ-(2,3,4-η^3-*nido*-hexaboranyl)-diplatinum(*Pt–Pt*), $[Pt_2(\mu\text{-}\eta^3\text{-}B_6H_9)_2(PMe_2Ph)_2]$, and of 2,4-dichloro-1,1-bis(dimethylphenylphosphine)-*closo*-1-nickeladecaborane, $[(PhMe_2P)_2NiB_9H_7Cl_2]$, 953–72
Co-ordination chemistry of pyridyl and *N*-methylimidazolyl ketones. Synthetic and *X*-ray structural studies of copper(II), nickel(II), and dimethylgold(III) complexes, 981–6
Studies on singlet oxygen in aqueous solution. Part 2. Water-soluble square-planar nickel complexes as quenchers, 1147–50
Macrocyclic polyphosphane ligands. Cobalt(II) and nickel(II) complexes of 5*RS*,8*RS*,16*RS*,19*RS*-tetraphenyl-5,8,16,19-tetraphospha-1,12-dithiacyclodocosane (δ-L^1) and the crystal structure of $[Co(\delta\text{-}L^1)][BF_4]_2{\cdot}0.5H_2O$, 1179–82
Some water-soluble Schiff-base complexes of nickel(II): the role of water, 1451–4
Reactions of metal ions with triketones in solution. Part 1. Formation constants for the systems of heptane-2,4,6-trione, 1-phenylhexane-1,3,5-trione, 1,5-diphenylpentane-1,3,5-trione, and 2,2′-dihydroxybenzophenone with proton, nickel(II), and cobalt(II), 1543–6
Potentiometric study of the complex-formation equilibria of manganese(II), cobalt(II), nickel(II), copper(II), and zinc(II) with ethylenediamine-*N*-acetic acid, 1605–8
Copper(II) salts of metal dithiolates, 1731–2
The stability of nickel(II) complexes of tetra-aza macrocycles, 1877–80
Metal complexes of vitamin B_6 related compounds. Crystal and molecular structures of aqua(5′-phosphopyridoxylideneglycinato)copper(II) trihydrate and bis(pyridoxylideneglycinato)nickel(II) hexahydrate, 2051–8
Reactions of tris[(2*S*)-2-(aminomethyl)pyrrolidine]nickel(II) ion with alk-3-en-2-ones or 4-hydroxyalkan-2-ones: formation of an optically active tetra-aza macrocycle, 2139–44
Unidentate *versus* symmetrically and unsymmetrically bidentate nitrate co-ordination in pyrazole-containing chelates. The crystal and molecular structures of (nitrato-*O*)[tris(3,5-dimethylpyrazol-1-ylmethyl)amine]copper(II) nitrate, (nitrato-*O*,*O*′)[tris(3,5-dimethylpyrazol-1-ylmethyl)amine]nickel(II)nitrate, and (nitrato-*O*)(nitrato-*O*,*O*′)[tris(3,5-dimethylpyrazol-1-ylmethyl)amine]cadmium(II), 2177–84
Synthesis and *X*-ray crystal structure of the asymmetric trinuclear complex $[Ni_3(\mu_3\text{-}S)_2(H_2O)(PPh_3)_5][PF_6]_2$, 2209–12
One-electron reduction of nickel(IV) oxime complexes, 2213–6
Synthesis and cryptate complexes of azathia macropolycyclic ligands based on 12-membered N_2S_2 and 15-membered N_2S_3 macrocyclic subunits, 2311–8
Reactivity of $[NiR(R')L_2]$ compounds and the crystal structure of $[Ni(C_2Cl_3)(C_6H_2Me_3\text{-}2,4,6)(PMe_2Ph)_2]$, 2333–42
Investigation into aroylhydrazones as chelating agents. Part 7. Synthesis and spectroscopic characterization of complexes of Mn^{II}, Co^{II}, Ni^{II}, Cu^{II}, and Zn^{II} with 2,6-diacetylpyridine bis(2-aminobenzoylhydrazone) and *X*-ray structure of chloro[2,6-diacetylpyridine bis(2-aminobenzoylhydrazone)](methanol)manganese(II) chloride monohydrate, 2387–92
Reactions of metal ions with triketones. Part 2. Kinetics and mechanisms of the reactions of nickel(II) and cobalt(II) with heptane-2,4,6-trione in methanol–water (70:30 v/v), 2565–70
NIOBIUM
Niobium(IV) sulphidohalides: preparation of $Nb_2X_4S_3$ and $Nb_2X_4S{\cdot}nL$[X = Br or Cl; *n* = 4, L = NCMe, SMe_2, or tetrahydrothiophene (tht); *n* = 2, L = $PhSCH_2CH_2SPh$]. Crystal and molecular structure of $Nb_2Cl_4S_2{\cdot}4tht$, 417–22
Axially asymmetric metal alkyls. Part 3. Chemical, electrochemical, and structural studies of group 5A $d^{0,1,2}$ metallepines $[M\{(2\text{-}CH_2C_6H_4)_2\}(\eta\text{-}C_5H_5)_2]^z$ (M = Nb or Ta; *z* = 1−, 0, or 1+); synthesis of $[Nb\{2\text{-}CH_2(C_6H_4)_2Me\text{-}2'\}X(\eta\text{-}C_5H_5)_2]$ (X = Cl or CO) and $[\{(\eta\text{-}C_5H_5)_2ClV\}_2\{(2\text{-}CH_2C_6H_4)_2\}]$, and electrochemistry of $[M(CH_2Ph)_{2-n}Cl_n(\eta\text{-}C_5H_5)_2]$ (*n* = 0, M = Nb, or Ta; *n* = 1, M = V), 1747–60
High-co-ordination-number compounds of niobium and tantalum: Reactions of niobium and tantalum halides and sulphido-halides with sodium diethyldithiocarbamate. The crystal structures of $[Nb(S_2CNEt_2)_4]Br$, $Nb(S_2CNEt_2)_3S$, and $Ta(S_2CNEt_2)_3(S_2)$, 1821–8
New carbonyl derivatives of niobium(I) and tantalum(I), 1989–96
Ion exchange of $K_4Nb_6O_{17}{\cdot}3H_2O$, 2349–52
NITRATE
Lewis-base adducts of Group 1B metal(I) compounds. Part 12. Structural studies of some bis(methyl-substituted pyridine)-copper(I) and -silver(I) nitrates and perchlorates, 117–24
The crystal structure and electronic properties of the complex acetatobis(1,10-phenanthroline)copper(II) perchlorate dihydrate, acetatobis(1,10-phenanthroline)copper(II) nitrate dihydrate, and acetatobis(1,10-phenanthroline)zinc(II) tetrafluoroborate

NITRATE (contd)
dihydrate, 141–50
Chemical and structural aspects of silver–triphenylarsine complexes and silver–tin complex salts, 321–32
Investigation into diphosphine oxides as ligands in diorganotin(IV) adducts. Part 3. Synthesis and crystal structure of two adducts of dinitratodiphenyltin(IV) with *cis*- and *trans*-1,2-bis(diphenylphosphoryl)ethylene, 487–92
Synthesis and characterisation of uranium(IV) nitrate complexes with piperazines, 1985–8
Unidentate *versus* symmetrically and unsymmetrically bidentate nitrate co-ordination in pyrazole-containing chelates. The crystal and molecular structures of (nitrato-*O*)[tris(3,5-dimethylpyrazol-1-ylmethyl)amine]copper(II) nitrate, (nitrato-*O*,*O'*)[tris(3,5-dimethylpyrazol-1-ylmethyl)amine]nickel(II)nitrate, and (nitrato-*O*)(nitrato-*O*,*O'*)[tris(3,5-dimethylpyrazol-1-ylmethyl)amine]cadmium(II), 2177–84
NITRIC ACID
Electrochemical studies in HNO_3–N_2O_4 mixtures: corrosion of stainless steel in HNO_3–N_2O_4 mixtures and the effect of inhibitors, 2551–4
NITRIDE
Rhenium nitrido-, arylimido-, nitrile, and carbonyl complexes with sterically hindered thiolate ligands, 2305–10
NITRILOTRIETHOXY
Kinetics and mechanism of the equilibration reaction between (2,2',2''-nitrilotriethoxy)nitrosylvanadate(1–) and cyanide. Crystal structures of sodium (2,2',2''-nitrilotriethoxy)nitrosylvanadate(I)–sodium perchlorate tetrahydrate and of barium cyano(2,2',2''-nitrilotriethoxy)nitrosylvanadate(I) pentahydrate, 2493–8
NITROGEN
Studies of phosphazenes. Part 21. Associative and dissociative pathways in the aminolysis reactions of halogenocyclotriphosphazenes, 285–90
Structural and nuclear magnetic resonance studies of short selenium–nitrogen bonds, 565–72
Studies of phosphazenes. Part 22. High-field nuclear magnetic resonance investigation of novel isomeric oxophosphazadienes, 1431–4
Mass spectrometric studies on cyclo- and poly-phosphazenes. Part 2. Oligomerization of hexa(aryloxy)cyclotriphosphazatrienes, 1547–54
Studies of phosphazenes. Part 25. Synthesis, nuclear magnetic resonance spectroscopy, and mode of formation of (aziridino)(triphenylphosphazenyl)cyclotriphosphazenes. *X*-ray crystal structure and enzyme-inhibiting activity of $N_3P_3(NPPh_3)(NC_2H_4)_5$, 1881–90
Structure and relative energies of some nitrogen-containing radical cations by MNDO calculation, 1915–22
Studies of phosphazenes. Part 21. Associative and dissociative pathways in the aminolysis reactions of halogenocyclotriphosphazenes (1985, 285), 2459–60
NITROGEN-15
Addition of proton and boron trifluoride at terminal and bridging cyano-ligands in dinuclear manganese compounds, 1609–12
Nitrogen-15 nuclear magnetic resonance relaxation mechanisms in dinitrogen complexes of molybdenum, tungsten, rhenium, and osmium, 2473–8
NITROGEN MONOXIDE
Reactions of bis(cyclopentadienyl)vanadium derivatives with nitrogen mono-oxide and the crystal structure of an oxo-bridged nitrosyl complex of vanadium, 1435–42
NITROGEN OXIDE
Nitrosoarene complexes of ruthenium and platinum and their reactions with NO and NO^+, 401–4
Insertion of NO into transition metal–aryl bonds: formation of zirconium complexes containing the $[ON(R)NO]^-$ (R = Ph or *p*-MeC_6H_4) ligand, 405–8
NITROIMIDAZOLE
Platinum(II) complexes of nitroimidazoles: synthesis, characterisation, and *X*-ray crystal structures of *cis*-dichlorobis[1-(2'-hydroxyethyl)-2-hydroxymethyl-5-nitroimidazole)platinum(II) and *trans*-dichlorobis[1-(2'-hydroxy-3'-methoxypropyl)-2-nitroimidazole]platinum(II), 795–802
NITROSOARENE
Nitrosoarene complexes of ruthenium and platinum and their reactions with NO and NO^+, 401–4
NITROSOBENZENE
Insertion of carbon dioxide, of CO_2-like molecules, and of other unsaturated compounds into the platinum–nitrogen bond of $[Pt(PPh_3)_2(PhNO)]$, 163–8
NITROSOHYDROXYLAMIDE
Insertion of NO into transition metal–aryl bonds: formation of zirconium complexes containing the $[ON(R)NO]^-$ (R = Ph or *p*-MeC_6H_4) ligand, 405–8
NITROSYL
Nitrosoarene complexes of ruthenium and platinum and their reactions with NO and NO^+, 401–4
Complexes of the platinum metals. Part 23. Synthesis of the nitrosyl carboxylate complexes $[M(O_2CR)_2(NO)(PPh_3)_2]$ (M = Rh or Ir; R = CF_3, C_2F_5, or C_6F_5): crystal and molecular structures of the trifluoroacetate derivatives $[M(O_2CCF_3)_2(NO)(PPh_3)_2]$, 611–6
Complexes of the platinum metals. Part 24. The role of dioxygen in the reactions of trifluoroacetic acid with the rhodium and iridium nitrosyls $[M(NO)(PPh_3)_3]$, 617–20
Complexes of the platinum metals. Part 25. Reactions of ruthenium and osmium nitrosyls $[MH(NO)(PPh_3)_3]$ and $[M(NO)_2(PPh_3)_2]$ with perfluorocarboxylic acids: *X*-ray crystal structure determination of nitrosyl(trifluoroacetato)-(trifluoroacetohydroximato-*OO'*)bis(triphenylphosphine)osmium(II)–dichloromethane (1/1), 621–8
On the effect of cyanide ion on the reaction of pentacyanonitrosylferrate(2–) with cysteine, 1191–4
Electrochemical and chemical properties of di-iodonitrosyl[tris(3,5-dimethylpyrazolyl)borato]molybdenum, $[Mo\{HB(Me_2pz)_3\}(NO)I_2]$, and related complexes, 1249–54
Reactions of bis(cyclopentadienyl)vanadium derivatives with nitrogen mono-oxide and the crystal structure of an oxo-bridged nitrosyl complex of vanadium, 1435–42
Rhenium nitrosyl complexes with simple and with sterically demanding aromatic thiolate ligands: *X*-ray crystal structures of $[PPh_4][Re_2(SC_6H_4Me\text{-}4)_7(NO)_2]\cdot CH_2Cl_2$ and $[Re(SC_6H_3Pr^i_2\text{-}2,6)_4(NO)]$, 1533–42
A new reaction of nitrosyl complexes; one-electron reduction of *trans*-$[MX(NO)L_4]^{2+}$ [M = Ru or Fe, X = Cl or Br, L = pyridine or *o*-phenylenebis(dimethylarsine)] with hydroxylamine, 1733–4
Complexes of the platinum metals. Part 30. Fragmentation reactions of rhodium and iridium trichloro- and tribromo-acetates, 2113–20
Electrophilic nitrosyls: preparation, structure, and reactivity of *cis*-chloronitrosylbis(pyridine-2-carboxylato)ruthenium and related complexes, 2427–32
Kinetics and mechanism of the equilibration reaction between (2,2',2''-nitrilotriethoxy)nitrosylvanadate(1–) and cyanide. Crystal structures of sodium (2,2',2''-nitrilotriethoxy)nitrosylvanadate(I)–sodium perchlorate tetrahydrate and of barium cyano(2,2',2''-nitrilotriethoxy)nitrosylvanadate(I) pentahydrate, 2493–8
N.M.R.
Co-ordinative interactions in chelated complexes of silicon. Part 7. Dynamic nuclear magnetic resonance studies of fluorine exchange at five-co-ordinated silicon, 5–8
Tin-119 Mössbauer and nuclear magnetic resonance studies of organotin compounds. Part 1. Sterically crowded tetraorganotin derivatives, 169–76
Synthesis, solution structure, and halide addition compounds of bis(dicyclohexylphosphinothioito)mercury(II), 223–4
Static and dynamic nuclear magnetic resonance studies of complexes of trimethylplatinum(IV) halides with olefinic thio-and seleno-ethers. *X*-Ray crystal structures of $[PtXMe_3(MeSeCH{=}CHSeMe)]$ (X = Cl or I), 345–54
Structural and nuclear magnetic resonance studies of short selenium–nitrogen bonds, 565–72
Aluminium-27 and hydrogen-1 nuclear magnetic resonance studies of solutions of aluminium salts in alcohol–chloroform mixtures, 591–6
Synthesis, characterization, and structure of the complex $[FeH(H_2BH_2)\{CH_3C(CH_2PPh_2)_3\}]$, 605–10
Complexes of the platinum metals. Part 26. Multinuclear nuclear magnetic resonance studies on rhodium(II) carboxylate adducts, 629–34
Platinum(II) complexes of nitroimidazoles: synthesis, characterisation, and *X*-ray crystal structures of *cis*-dichlorobis[1-(2'-hydroxyethyl)-2-hydroxymethyl-5-nitroimidazole)platinum(II) and *trans*-dichlorobis[1-(2'-hydroxy-3'-methoxypropyl)-2-nitroimidazole]platinum(II), 795–802

N.M.R. (contd)
Mercurated and tellurated Schiff bases and phenylhydrazones, 821–6
Lewis-base adducts of group 1B metal(I) compounds. Part 16. Synthesis, structure, and solid-state phosphorus-31 nuclear magnetic resonance spectra of some novel $[Cu_xX_4L_4]$ (X = halogen, L = N,P base) 'cubane' clusters, 831–8
Multinuclear nuclear magnetic resonance study of the interaction of some phosphorus(V) compounds with inorganic acids. The protonating abilities of HNO_3, $MeSO_3H$, and HPO_2F_2 towards the phosphoryl group, 865–6
Diastereoisomeric organophosphorus compounds. Part 4. Proton and phosphorus-31 nuclear magnetic resonance spectra of compounds of the type [RR'P(X)]$_2$Y and RR'P(X)-Y-(Z)PRR' (R = CH_3, R' = t-C_4H_9) (1984, 2803), 871–2
Reactions of 6,6'-bis(*nido*-decaboranyl) oxide and 6-hydroxy-*nido*-decaborane with dihalogenobis(phosphine) complexes of nickel, palladium, and platinum, and some related chemistry; nuclear magnetic resonance investigations and the crystal and molecular structures of bis(dimethylphosphine)-di-μ-(2,3,4-η^3-*nido*-hexaboranyl)-diplatinum(*Pt–Pt*), $[Pt_2(\mu\text{-}\eta^3\text{-}B_6H_9)_2(PMe_2Ph)_2]$, and of 2,4-dichloro-1,1-bis(dimethylphenylphosphine)-*closo*-1-nickeladecaborane, $[(PhMe_2P)_2NiB_9H_7Cl_2]$, 953–72
Bimetallic systems. Part 9. The synthesis of and nuclear magnetic resonance studies on 10-membered ring complexes of type $[(OC)_4M^1(\mu\text{-}Ph_2PCH_2CH_2PPh_2)_2M^2(CO)_4]$ (M^1, M^2 = Cr, Mo, or W), 1009–14
Geometry-dependent complexation effects in carbon-13 nuclear magnetic resonance spectra of tricarbonyl(3—8-η-[2.2]paracyclophane)chromium and related complexes, 1065–8
Polyhedral rhenaborane chemistry: crystal and molecular structures of the *nido*-6-rhenadecaborane cluster compounds [6,6,6,6-$(PMe_2Ph)_3$H-*nido*-6-ReB_9H_{13}] and [2-(PMe_2Ph)-6,6,6,6-$(PMe_2Ph)_2$ClH-*nido*-6-ReB_9H_{12}]; nuclear magnetic resonance parameters of these and other related *nido*-rhenadecaborane cluster species, 1119–30
High-field vanadium-51 and oxygen-17 nuclear magnetic resonance study of peroxovanadates(V), 1173–8
Synthesis, properties, and multinuclear magnetic resonance (^{1}H, ^{77}Se, and ^{195}Pt) studies on diselenoether complexes of palladium, platinum, and rhodium, 1265–70
Studies of phosphazenes. Part 22. High-field nuclear magnetic resonance investigation of novel isomeric oxophosphazadienes, 1431–4
Steric changes at labelled NH_2 sites during the base-catalysed ammoniation of the *trans*-azido(dimethyl sulphoxide)bis(ethylenediamine)cobalt(III) ion, 1495–500
Synthetic and nuclear magnetic resonance studies on dialkyl- and diaryl-platinum complexes containing chelating, monodentate, or bridging $Ph_2PCH_2PPh_2$ ligands, 1501–6
Mono- and di-nuclear rhodium and palladium complexes of macrocyclic ligands containing the 2,6-di(thiomethyl)pyridine sub-unit, 1517–22
The syntheses and characterisation of, and the determination of $^1J(^{103}Rh–^{103}Rh)$ in$[(\eta^5\text{-}C_5Me_5)_2Rh_2(\mu\text{-}CH_2)_2\{\mu\text{-}CH_2CR(CH_2CR{=}CH_2)CH_2\}]$ (R = H or Me) and $[(\eta^5\text{-}C_5Me_5)_2Rh_2(\mu\text{-}CH_2)_2\{\mu\text{-}Ph_2P(CH_2)_nPPh_2\}]^{2+}$ (*n* = 1 or 2), 1555–60
Pyramidal inversions and 1,2-metal shifts in pentacarbonyl-chromium, -molybdenum, and -tungsten derivatives of dialkyl disulphides and dialkyl diselenides. A nuclear magnetic resonance investigation, 1561–8
Dynamic nuclear magnetic resonance study of Group 6 metal pentacarbonyl complexes. Pyramidal inversion and 1,2-metal shifts in the complexes $[M(CO)_5(Me_2\overline{CCH_2EECH_2})]$ (M = Cr, Mo, or W; E = S or Se), 1569–76
Addition of proton and boron trifluoride at terminal and bridging cyano-ligands in dinuclear manganese compounds, 1609–12
Studies of 2,5;6,10;8,10-tri-μ-hydro-nonahydro-*nido*-nonaborate(1−), $[B_9H_{12}]^-$: preparation, crystal and molecular structure, nuclear magnetic resonance spectra, electrochemistry, and reactions, 1645–54
Geometry-dependent carbon-13 chemical shifts in (η^6-[2.2]cyclophane)(η^5-cyclopentadienyl)iron(II) hexafluorophosphates, 1661–4
Intermediates in the photochemical reaction of tetraphosphorus trisulphide with organic disulphides; phosphorus-31 nuclear magnetic resonance parameters for 2,6-bis(alkylthio)- and 2,6-diiodo-3,5,7-trithia-1,2,4,6-tetraphosphabicyclo[2.2.1]heptanes and for 3,6-bis(alkylthio)-2,5,7-trithia-1,3,4,6-tetraphosphabicyclo[2.2.1]heptanes, 1707–12
Studies of gold cluster compounds using high-resolution ^{31}P solid-state nuclear magnetic resonance spectroscopy, 1811–4
Studies of phosphazenes. Part 25. Synthesis, nuclear magnetic resonance spectroscopy, and mode of formation of (aziridino)(triphenylphosphazenyl)cyclotriphosphazenes. *X*-ray crystal structure and enzyme-inhibiting activity of $N_3P_3(NPPh_3)(NC_2H_4)_5$, 1881–90
Carbon-13 nuclear magnetic resonance evidence of a relaxation process dominated by scalar coupling with a quadrupolar nucleus in $[Re_3(\mu\text{-}H)_4(CO)_{10}]^-$, 1899–902
Bis(triphenylphosphine)copper(I) derivatives of substituted *arachno* nine-vertex borane anions, $Cu(PPh_3)_2(B_9H_{13}X)$ (X = H, NCS, NCSe, $NCBPh_3$, $NCBH_3$, or $NCBH_2NCBH_3$), 1903–8
Complexes of organoaluminium compounds. Part 13. Preparation and nuclear magnetic resonance spectra of the arylamido-compounds $AlMe_2(NHR')$ (R' = Ph, C_6H_4Me-*o*, C_6H_4Me-*p*, $C_6H_3Me_2$-2,6) and the imido-compounds AlMe(NR')$_3$. Crystal and molecular structures of $[\{AlMe_2(NHC_6H_4Me\text{-}o)\}_2]$ and $[\{AlMe(NPh)\}_6]$, 1929–34
Oxygen exchange and protonation of polyanions: a multinuclear magnetic resonance study of tetradecavanadophosphate(9−) and decavanadate(6−), 1953–8
Synthesis, properties, and ^{77}Se nuclear magnetic resonance studies of platinum metal complexes of two isomeric tris(selenoethers), $MeC(CH_2SeMe)_3$ and $Se(CH_2CH_2CH_2SeMe)_2$, 2185–90
Pyramidal inversion in configurational isomers of tetracarbonyl[1,1,2,2-tetrakis(methylthio)ethane]chromium(0): a two-dimensional nuclear magnetic resonance exchange study, 2195–202
Steric changes at labelled NH_2 sites during the base-catalysed ammoniation of the *trans*-azido(dimethyl sulphoxide)bis(ethylenediamine)cobalt(III) ion, 2223–4
Synthesis and nuclear magnetic resonance studies of co-ordinatively unsaturated alkyne complexes of tungsten(II), 2239–46
Carbon-13 nuclear magnetic resonance study of the complexes formed between zinc(II) and triethylenetetramine, 2381–6
Polyhedral ruthenaborane chemistry: characterization of several new ruthenaboranes by nuclear magnetic resonance spectroscopy, and the crystal and molecular structure of [5,6,6-$(PPh_3)_3$-6-H-*nido*-6-RuB_9H_{12}], 2397–406
Two unusual *closo*-type ruthenaboranes: preparation, molecular structure, and nuclear magnetic resonance properties of [1,1,1-(PPh_3)HCl-1-RuB_9H_7-3,5-$(PPh_3)_2$] and [1,1-$(PPh_3)_2$-1-$RuB_{10}H_8$-2,5-$(OEt)_2$], 2407–16
Nitrogen-15 nuclear magnetic resonance relaxation mechanisms in dinitrogen complexes of molybdenum, tungsten, rhenium, and osmium, 2473–8
The chemistry of heteroarylphosphorus compounds. Part 16. Unusual substituent effects on selenium-77 nuclear magnetic resonance chemical shifts of heteroaryl- and aryl-phosphine selenides. *X*-Ray crystal structure of tri(2-furyl)phosphine selenide, 2505–8
Mixed-ligand complexes of trivalent lanthanides. Part 3. Complexes of heptafluorodimethyloctane-3,5-dione and pyrazine: syntheses and spectral studies, 2547–50
Conformation analysis of compounds of the type $[Fe(\eta^5\text{-}C_5H_5)(CO)(PPh_3)(CH_2R)]$ (R = alkyl or aryl): a comment, 2691
Conformation analysis of compounds of the type $[Fe(\eta^5\text{-}C_5H_5)(CO)(PPh_3)(CH_2R)]$ (R = alkyl or aryl): a reply, 2691

NONABORATE
Studies of 2,5;6,10;8,10-tri-μ-hydro-nonahydro-*nido*-nonaborate(1−), $[B_9H_{12}]^-$: preparation, crystal and molecular structure, nuclear magnetic resonance spectra, electrochemistry, and reactions, 1645–54
Bis(triphenylphosphine)copper(I) derivatives of substituted *arachno* nine-vertex borane anions, $Cu(PPh_3)_2(B_9H_{13}X)$ (X = H, NCS, NCSe, $NCBPh_3$, $NCBH_3$, or $NCBH_2NCBH_3$), 1903–8

NONYLPYRIDINE OXIDE
Solvent extraction of gold and platinum-group metals using 2-nonylpyridine 1-oxide, and the crystal and molecular structure of bis(2-nonylpyridine 1-oxide)hydrogen(1+) tetrachloroaurate(III), 771–6

NORBORNADIENE
1*H*-pyrrolo[2,3-*b*]pyridine (HL) ligands in rhodium(I) and iridium(I) chemistry. Crystal and molecular structures of $[Rh_2(\mu\text{-}L)_2(nbd)_2]$ and $[Rh_4(\mu\text{-}Cl)_2(\mu\text{-}L)_2(\mu\text{-}CO)_2(CO)_2(nbd)_2]$, 1891–8

N.Q.R.
Bromine nuclear quadrupole resonance studies of some hexabromostannates: *X*-Ray crystal structure of pyridinium hexabromostannate(IV), $[Hpy]_2[SnBr_6]$, 1399–404
OCTAMOLYBDATE
Photochemical studies of the alkylammonium molybdates. Part 7. Octahedral sites for multi-electron reduction of $[Mo_8O_{26}(MoO_4)_2]^{8-}$, 2585–90
OCTANEDIONATE
Mixed-ligand complexes of trivalent lanthanides. Part 3. Complexes of heptafluorodimethyloctane-3,5-dione and pyrazine: syntheses and spectral studies, 2547–50
OLEFIN
Carbene complexes. Part 18. Synthetic routes to electron-rich olefin-derived monocarbenerhodium(I) neutral and cationic complexes and their chemical and physical properties (1984, 2355), 229–30
OLIGOMERIZATION
Mass spectrometric studies on cyclo- and poly-phosphazenes. Part 2. Oligomerization of hexa(aryloxy)cyclotriphosphazatrienes, 1547–54
OSMIUM
Metallation of 2-ethenylpyridine at triosmium clusters: *X*-ray crystal structures of the open trinuclear clusters $[Os_3H(CO)_9L(NC_5H_4CH{=}CH)]$ (L = CO or PMe_2Ph), 85–90
Chemistry of the unsaturated cluster compound $[Os_3Pt(\mu\text{-}H)_2(CO)_{10}\{P(cyclo\text{-}C_6H_{11})_3\}]$; *X*-ray crystal structures of $[Os_3Pt(\mu\text{-}H)_2(CO)_{11}\{P(cyclo\text{-}C_6H_{11})_3\}]$, $[Os_3Pt(\mu\text{-}H)_4(CO)_{10}\{P(cyclo\text{-}C_6H_{11})_3\}]$, and $[Os_3Pt(\mu\text{-}H)_2(\mu\text{-}CH_2)(CO)_{10}\{P(cyclo\text{-}C_6H_{11})_3\}]$ (two isomers), 177–90
Co-ordination chemistry of higher oxidation states. Part 13. Synthesis and properties of alkali-metal hydroxo-oxo-osmate(VIII) compounds and the molecular structure of $Cs[O_4Os(\mu\text{-}OH)OsO_4]$, 199–204
Synthesis and properties of dioxo-osmium(VI) compounds of thio- and seleno-ethers, 205–8
Two-dimensional nuclear magnetic resonance for the analysis of the carbon-13 spectra of carbonyl groups in metallocarbonyls, 225–8
The preparation and hydridic reduction of dicationic dicarbonyl complexes of osmium(II); the crystal and molecular structure of *trans*-bis[1,2-bis(diphenylphosphino)ethane-*PP'*]carbonylformylosmium(II) hexafluoroantimonate–dichloromethane (1/1), 387–94
Systematic synthesis of tetranuclear osmium clusters by the reaction of trinuclear clusters with $[OsH_2(CO)_4]$; crystal structure of $[Os_4H_3Br(CO)_{13}]$, 555–64
Mononuclear η^6-*p*-cymeneosmium(II) complexes and their reactions with Al_2Me_6 and other methylating reagents, 573–8
Complexes of the platinum metals. Part 25. Reactions of ruthenium and osmium nitrosyls $[MH(NO)(PPh_3)_3]$ and $[M(NO)_2(PPh_3)_2]$ with perfluorocarboxylic acids: *X*-ray crystal structure determination of nitrosyl(trifluoroacetato)-(trifluoroacetohydroximato-*OO'*)bis(triphenylphosphine)osmium(II)–dichloromethane (1/1), 621–8
Electrochemically induced ligand substitutions on $[OsCl_3(PMe_2Ph)_3]$: rational pathways to osmium(II) complexes, 947–52
New isomers of $[Os_3(CO)_{10}(PMe_2Ph)_2]$ and $[Os_3(CO)_9(PMe_2Ph)_3]$, 1037–42
The reaction of $[Os_3(\mu\text{-}H)_2(CO)_9L]$ (L = CO or PEt_3) with dimethylcyanamide, Me_2NCN: *X*-Ray crystal structure of $[Os_3(\mu\text{-}H)(\mu\text{-}NCHNMe_2)(CO)_{10}]$ and the reactions of this complex with acids, 1355–60
Preparation and vibrational spectra of $[OsX_6]^{3-}$ (X = Cl, Br, or I) and of other platinum-group hexahalogeno-complexes, 1673–6
Synthesis and properties of alkali-metal tetraoxo-osmate(VII) compounds, 1735–6
Magnetic circular dichroism spectra of tris-chelate complexes of 2,2'-bipyridyl and 1,10-phenanthroline with iron(II), ruthenium(II), and osmium(II) at 4.2 K, 1781–8
Reactions of some decaosmium clusters with electrophilic and nucleophilic reagents: *X*-ray structure analyses of $[N(PPh_3)_2][Os_{10}C(CO)_{24}(\mu\text{-}I)]$, $[Os_{10}C(CO)_{24}(\mu\text{-}I)_2]$, $[N(PPh_3)_2]_2[Os_{10}C(CO)_{22}(NO)I]$, $[Os_{10}C(CO)_{23}\{P(OMe)_3\}(\mu\text{-}I)_2]$ and of two isomers of $[Os_{10}C(CO)_{21}\{P(OMe)_3\}_4]$, 1795–810
The chemistry and catalytic properties of ruthenium and osmium complexes. Part 1. Homogeneous catalysis of organic reactions by bromo(carbonyl)hydridotris(triphenylphosphine)osmium(II), 1859–64
Complexes of the platinum metals. Part 29. Pyridine-2-thiolate derivatives of ruthenium and osmium: *X*-ray crystal structures of $[Ru(C_5H_4NS)_2(CO)_2(PPh_3)]$ and $[Ru(C_5H_4NS)_2(CO)(PPh_3)]$, 2101–12
Electrophilic nitrosyls: preparation, structure, and reactivity of *cis*-chloronitrosylbis(pyridine-2-carboxylato)ruthenium and related complexes, 2427–32
Synthesis of tetra- and penta-nuclear platinum–osmium carbido-cluster complexes from non-carbido-precursors: *X*-ray structural studies on $[Os_3Pt(\mu\text{-}H)_2(\mu_4\text{-}C)(CO)_{10}\{P(cyclo\text{-}C_6H_{11})_3\}]$, $[Os_3Pt_2(\mu\text{-}H)_2(\mu_5\text{-}C)(\mu\text{-}CO)(CO)_9\{P(cyclo\text{-}C_6H_{11})_3\}_2]$, and $[Os_3Pt_2(\mu\text{-}H)(\mu_5\text{-}C)(\mu\text{-}OMe)(\mu\text{-}CO)(CO)_9\{P(cyclo\text{-}C_6H_{11})_3\}_2]$, 2437–48
Nitrogen-15 nuclear magnetic resonance relaxation mechanisms in dinitrogen complexes of molybdenum, tungsten, rhenium, and osmium, 2473–8
Acid-induced displacement of acetaldehyde from a μ-vinyloxy-triosmium cluster, 2479–82
OXALATE
Light-induced electron-transfer reactions. Part 3. Kinetics of the oxidation reaction of oxalate ion by peroxodisulphate ion, induced by irradiation with visible light of an aqueous solution containing tris(2,2'-bipyridine)ruthenium(II), 355–60
OXALIC ACID
Equilibrium and structural studies of silicon(IV) and aluminium(III) in aqueous solution. Part 13. A potentiometric and ^{27}Al nuclear magnetic resonance study of speciation and equilibria in the aluminium(III)–oxalic acid–hydroxide system, 2665–70
OXIDATION
Light-induced electron-transfer reactions. Part 3. Kinetics of the oxidation reaction of oxalate ion by peroxodisulphate ion, induced by irradiation with visible light of an aqueous solution containing tris(2,2'-bipyridine)ruthenium(II), 355–60
Catalytic air oxidation of benzoin in the presence of dioxomolybdenum(VI) complexes with sulphur chelate ligands, 635–40
Reduction–oxidation properties of organotransition-metal complexes. Part 20. Oxidative and thermolytic cyclopropane ring-opening reactions; *X*-ray crystal structure of $[Fe_2(CO)_6(\eta^4\text{:}\eta'^4\text{-}C_{16}H_{18})]$, 699–706
Studies on singlet oxygen in aqueous solution. Part 1. Formation of singlet oxygen from hydrogen peroxide with two-electron oxidants, 1141–6
Oxidation of thiourea by iodate: a new type of oligo-oscillatory reaction, 1669–72
Kinetics and mechanism of the oxidation of substituted malonic acids by the hexachloroiridate(IV) ion, 1683–8
Formation of oxidized molybdenum-bearing ferrites, $Fe_{2.95-x}Mo_xO_4$ (x = 0.03—0.30) in aqueous suspensions by air oxidation and valence state of molybdenum ions in the lattice, 1713–6
Stereochemical consequences of the two-electron oxidation of a dirhodium fulvalene complex: the *X*-ray crystal structures of *trans*-$[Rh_2(CO)_2(PPh_3)_2(\eta^5\text{:}\eta'^5\text{-}C_{10}H_8)]$ and *cis*-$[Rh_2(CO)_2(PPh_3)_2(\eta^5\text{:}\eta'^5\text{-}C_{10}H_8)][PF_6]_2$, 2283–90
OXIDE
Reactions in mixed non-aqueous systems containing sulphur dioxide. Part 6. The reaction of metal oxides with dimethyl sulphoxide–sulphur dioxide, 99–100
Co-ordination chemistry of higher oxidation states. Part 13. Synthesis and properties of alkali-metal hydroxo-oxo-osmate(VIII) compounds and the molecular structure of $Cs[O_4Os(\mu\text{-}OH)OsO_4]$, 199–204
Synthesis and properties of dioxo-osmium(VI) compounds of thio- and seleno-ethers, 205–8
Synthesis and structural assessment of ammonium and caesium difluorodioxoperoxouranates(VI), $A_2[UO_2(O_2)F_2]$ (A = NH_4 or Cs), and alkali-metal difluorodioxoperoxouranate(VI) monohydrates, $A_2[UO_2(O_2)F_2]\text{-}H_2O$ (A = K or Rb), 409–12
Tin–molybdenum oxides. A study by *X*-ray diffraction, Mössbauer spectroscopy, and electron spin resonance, 451–4
Characterisation of chromium(VI) oxide tetrafluoride, $CrOF_4$, and caesium pentafluoro-oxochromate(VI) $Cs[CrOF_5]$, 529–34
Actinide structural studies. Part 7. The crystal and molecular structures of (2,2'-bipyridyl)dinitratodioxo-uranium(VI) and -neptunium(VI), and diacetato(2,2'-bipyridyl)dioxo-uranium(VI) and -neptunium(VI), 1001–8

OXIDE (contd)
Reactions of bis(cyclopentadienyl)vanadium derivatives with nitrogen mono-oxide and the crystal structure of an oxo-bridged nitrosyl complex of vanadium, 1435–42
Magnetic exchange interactions in perovskite solid solutions. Part 5. The unusual defect structure of $SrFeO_{3-y}$, 1455–70
Synthesis and crystal structure of the layer compound Sb_3TeO_6Cl, 1633–6
Formation of oxidized molybdenum-bearing ferrites, $Fe_{2.95-x}Mo_xO_4$ (x = 0.03—0.30) in aqueous suspensions by air oxidation and valence state of molybdenum ions in the lattice, 1713–6
Oxoalkyls of rhenium-(v) and -(vi). *X*-Ray crystal structures of $(Me_4ReO)_2Mg(thf)_4$, $[(Me_3SiCH_2)_4ReO]_2Mg(thf)_2$, $Re_2O_3Me_6$, and $Re_2O_3(CH_2SiMe_3)_6$, 2167–76
A disagreement on the explanation of short and long As–O bonds of the $(As–O)_4$ ring in $As_4(CF_3)_6O_6(OH)_2$ in terms of As^{III}–O and As^{V}–O bonds, 2221–2
Synthesis and characterisation of new mixed oxides of antimony and tellurium, 2225–30
Crystal structures and magnetism of binuclear iron(III) complexes with a linear oxo-bridge, $[Fe_2O(bbimae)_2X_2][NO_3]_2$ {bbimae = 2-[bis(benzimidazol-2-ylmethyl)amino]ethanol, X = Cl or NCS}, 2375–80
A novel pentameric hydrolysis product of $SnMe_2Cl_2$: crystal and molecular structure of $[NHEt_3][(SnMe_2Cl)_5O_3]$, 2683–6
OXIMATE
One-electron reduction of nickel(IV) oxime complexes, 2213–6
OXOPHOSPHAZADIENE
Studies of phosphazenes. Part 22. High-field nuclear magnetic resonance investigation of novel isomeric oxophosphazadienes, 1431–4
OXYACID
The easy catalytic reduction of $HClO_4$, HNO_3, and H_5IO_6 by CO in the presence of $[Pd_3(O_2CMe)_6]$, 1971–4
OXYGEN-17
High-field vanadium-51 and oxygen-17 nuclear magnetic resonance study of peroxovanadates(v), 1173–8
Oxygen exchange and protonation of polyanions: a multinuclear magnetic resonance study of tetradecavanadophosphate(9−) and decavanadate(6−), 1953–8
OXYGEN-18
Oxygenation studies. Part 7. Catalytic dioxygenation of cyclo-octa-1,5-diene at a rhodium centre, 1591–6
OXYGENATION
Oxygenation studies. Part 7. Catalytic dioxygenation of cyclo-octa-1,5-diene at a rhodium centre, 1591–6
PALLADACYCLOBUTANONE
Chemistry of metallacyclobutanones. Part 2. Synthesis and ring inversion of some highly puckered metallacyclobutan-3-one (slipped oxodimethylenemethane) complexes of palladium; crystal structures of 2,4-bis(methoxycarbonyl)-1,1-bis(triphenylphosphine)palladacyclobutan-3-one, 2,4-bis(methoxycarbonyl)-1,1-bis(triphenylarsine)palladacyclobutan-3-one, and 1,1-(2′,2″-bipyridyl)-2,4-bis(methoxycarbonyl)palladacyclobutan-3-one, 259–68
PALLADIUM
Chemistry of metallacyclobutanones. Part 2. Synthesis and ring inversion of some highly puckered metallacyclobutan-3-one (slipped oxodimethylenemethane) complexes of palladium; crystal structures of 2,4-bis(methoxycarbonyl)-1,1-bis(triphenylphosphine)palladacyclobutan-3-one, 2,4-bis(methoxycarbonyl)-1,1-bis(triphenylarsine)palladacyclobutan-3-one, and 1,1-(2′,2″-bipyridyl)-2,4-bis(methoxycarbonyl)palladacyclobutan-3-one, 259–68
Bimetallic systems. Part 7. Platinum and palladium dicyanides containing terminal or bridging $Ph_2PCH_2PPh_2$ and heterobimetallics with silver, gold, mercury, rhodium, iridium, or molybdenum, 279–84
Carbaborane derivatives of the late- and post-transition elements. Part 3. Structural consequences of ligand substitution in palladadicarbadodecaboranes 3-L_2-3,1,2-$PdC_2B_9H_{11}$. The crystal and molecular structures of 3-$[Me_2N(CH_2)_2NMe_2]$-3,1,2-$PdC_2B_9H_{11}$ and 3-$(PMe_3)_2$-3,1,2-$PdC_2B_9H_{11}$, 761–70
Palladium(II,IV) mixed-valence complexes of 1,2-diaminoethane, 1,3-diaminopropane, and diethylenetriamine: syntheses, electronic, infrared, Raman, and resonance Raman spectra and *X*-ray studies, 815–20
Reactions of 6,6′-bis(*nido*-decaboranyl) oxide and 6-hydroxy-*nido*-decaborane with dihalogenobis(phosphine) complexes of nickel, palladium, and platinum, and some related chemistry; nuclear magnetic resonance investigations and the crystal and molecular structures of bis(dimethylphosphine)-di-μ-(2,3,4-η³-*nido*-hexaboranyl)-diplatinum(*Pt–Pt*), $[Pt_2(\mu\text{-}\eta^3\text{-}B_6H_9)_2(PMe_2Ph)_2]$, and of 2,4-dichloro-1,1-bis(dimethylphenylphosphine)-*closo*-1-nickeladecaborane, $[(PhMe_2P)_2NiB_9H_7Cl_2]$, 953–72
Dynamic behaviour of carbon-metallated palladium hydrazone complexes. Crystal structures of $[\{Pd[CH_2CMe_2C(=N\text{–}NMePh)Me]Cl\}_2]$ and $[\{Pd[CH_2C(=N\text{–}NMePh)Bu^t]Cl\}_2]$, 1155–62
Synthesis, properties, and multinuclear magnetic resonance (^1H, ^{77}Se, and ^{195}Pt) studies on diselenoether complexes of palladium, platinum, and rhodium, 1265–70
Mono- and di-nuclear rhodium and palladium complexes of macrocyclic ligands containing the 2,6-di(thiomethyl)pyridine sub-unit, 1517–22
Cyclopalladated derivatives of 2,4′-bipyridine, 1719–22
The easy catalytic reduction of $HClO_4$, HNO_3, and H_5IO_6 by CO in the presence of $[Pd_3(O_2CMe)_6]$, 1971–4
The palladium-catalysed reaction between $[Re_2(CO)_{10}]$ and phosphines and the crystal and molecular structure of diaxial $[Re_2(CO)_8(PMe_2Ph)_2]$, 2277–82
Preparation, characterization, and physical properties of the series MPd_3S_4 (M = rare earth), 2369–74
Kinetics and mechanism of ortho-palladation of ring-substituted *NN*-dimethylbenzylamines, 2629–38
PARACYCLOPHANE
Geometry-dependent complexation effects in carbon-13 nuclear magnetic resonance spectra of tricarbonyl(3—8-η-[2.2]paracyclophane)chromium and related complexes, 1065–8
PENDANT ARM
Studies of pendant-arm macrocyclic ligands. Part 4. Two penta-aza macrocycles based on 1-(2′-dimethylaminoethyl)-1,5,9,13-tetra-azacyclohexadecane and its complexes with bivalent metal ions, 1361–4
PENTA-AZABICYCLOOCTADECATRIENE
Studies in the flexibility of macrocycle ligands. Calculation of macrocycle cavity size by force-field methods. Crystal and molecular structures of $[CoLCl][ClO_4]_2$ and $[CuL][PF_6]_2$ {L = 2,13-dimethyl-3,6,9,12,18-penta-azabicyclo[12.3.1]octadeca-1(18),14,16-triene}, 1829–34
PENTACHLOROCYCLOPENTADIENYL
The structure of pentachlorocyclopentadienylphenylmercury(II) in the solid state and in solution; the effect of *X*-ray absorption and crystal decay on molecular geometry parameters, 471–4
PENTADIENYL
Reactivity of $[PPh_4][Fe_2(CO)_6\{\mu\text{-}CPhCPhC(CF_3)C(CF_3)H\}]$ toward electrophiles. *X*-Ray structure of $[Fe_2(CO)_6\text{-}\{\mu\text{-}CPhCPhC(CF_3)CHC(OEt)_2\}]$, a product resulting from fluorine abstraction at a CF_3 group and subsequent fluorine substitution by ethoxy-groups, 1981–4
He I and He II photoelectron spectra of open-chain pentadienyl complexes of manganese and rhenium, 2677–82
PENTADIENYLIDENE
Carbon–carbon formation at di-iron centres. Part 2. Reactivity of $[Fe_2(CO)_6(\mu\text{-}COEt)\{\mu\text{-}C(R)C(R)H\}]$ complexes toward MeOC(O)C≡CC(O)OMe (R = Ph) and $CF_3C{\equiv}CCF_3$ (R = Ph or H); *X*-ray crystal structures of $[Fe_2(CO)_5\{\mu\text{-}C(OEt)C[C(O)OMe]C[C(O)OMe]\}\{\mu\text{-}C(Ph)C(Ph)H\}]\text{-}H_2O$ and $[Fe_2(CO)_6\{\mu\text{-}C(CF_3)C(CF_3)CHCHCH(OMe)\}]$, 1087–94
PENTAMETHYLCYCLOPENTADIENYL
Synthesis, structures, and reactivities of some pentamethylcyclopentadienyl–sulphur compounds, 1303–8
The syntheses and characterisation of, and the determination of $^1J(^{103}Rh\text{–}^{103}Rh)$ in $[(\eta^5\text{-}C_5Me_5)_2Rh_2(\mu\text{-}CH_2)_2\{\mu\text{-}CH_2CR(CH_2CR{=}CH_2)CH_2\}]$ (R = H or Me) and $[(\eta^5\text{-}C_5Me_5)_2Rh_2(\mu\text{-}CH_2)_2\{\mu\text{-}Ph_2P(CH_2)_nPPh_2\}]^{2+}$ (n = 1 or 2), 1555–60
PEPTIDE
The L-proline residue as a 'break-point' in metal–peptide systems, 535–40
Specific binding of the tyrosine residue in copper(II) complexes of Tyr-Pro-Gly-Tyr and Tyr-Gly-Pro-Tyr, 1201–6
PERCHLORATE
Lewis-base adducts of Group 1B metal(I) compounds. Part 12. Structural studies of some bis(methyl-substituted pyridine)-copper(I) and -silver(I) nitrates and perchlorates, 117–24

PERCHLORATE (contd)
Lewis-base adducts of Group 1B metal(I) compounds. Part 13. Crystal structure determinations of tetrakis(triphenyl-phosphine)-copper(I) and -silver(I) perchlorates, bis(pyridine)bis(triphenylphosphine)copper(I) perchlorate, (2,2′-bipyridyl)bis(triphenylphosphine)copper(I) perchlorate, and tetrahydroboratobis(triphenylphosphine)copper(I)–pyridine (1/0.5), 125–34
The crystal structure and electronic properties of the complex acetatobis(1,10-phenanthroline)copper(II) perchlorate dihydrate, acetatobis(1,10-phenanthroline)copper(II) nitrate dihydrate, and acetatobis(1,10-phenanthroline)zinc(II) tetrafluoroborate dihydrate, 141–50
Chlorine trioxide, Cl_2O_6, a most efficient perchlorating reagent in new syntheses of anhydrous metal perchlorates, chloryl and nitryl perchloratometalates of cobalt(II), nickel(II), and copper(II). Reactivity of chlorine trioxide with anhydrous or hydrated chlorides and nitrates, 297–306
PERFLUOROCARBOXYLIC ACID
Complexes of the platinum metals. Part 25. Reactions of ruthenium and osmium nitrosyls $[MH(NO)(PPh_3)_3]$ and $[M(NO)_2(PPh_3)_2]$ with perfluorocarboxylic acids: *X*-ray crystal structure determination of nitrosyl(trifluoroacetato)-(trifluoroacetohydroximato-*OO′*)bis(triphenylphosphine)osmium(II)–dichloromethane (1/1), 621–8
PEROVSKITE
Magnetic exchange interactions in perovskite solid solutions. Part 5. The unusual defect structure of $SrFeO_{3-y}$, 1455–70
PEROXIDE
Synthesis and structural assessment of ammonium and caesium difluorodioxoperoxouranates(VI), $A_2[UO_2(O_2)F_2]$ (A = NH_4 or Cs), and alkali-metal difluorodioxoperoxouranate(VI) monohydrates, $A_2[UO_2(O_2)F_2]\cdot H_2O$ (A = K or Rb), 409–12
High-field vanadium-51 and oxygen-17 nuclear magnetic resonance study of peroxovanadates(V), 1173–8
PEROXODISULPHATE
The peroxodisulphate–iodide reaction. Reactivity and ionic association and solvation in isodielectric water–solvent mixtures, 31–4
Light-induced electron-transfer reactions. Part 3. Kinetics of the oxidation reaction of oxalate ion by peroxodisulphate ion, induced by irradiation with visible light of an aqueous solution containing tris(2,2′-bipyridine)ruthenium(II), 355–60
The peroxodisulphate–hexacyanoferrate(II) reaction. Reactivity and ionic association in isodielectric water–co-solvent mixtures, 1975–6
PEROXY-ACID
Studies on singlet oxygen in aqueous solution. Part 3. The decomposition of peroxy-acids, 1151–4
PHENANTHROLINE
Co-ordinative interactions in chelated complexes of silicon. Part 5. Chirality of five-co-ordinate silicon compounds: crystal and molecular structures of 1-(chlorodimethylsilyl)-and 1-(dichloromethylsilyl)-1,2,3,4-tetrahydro-1,10-phenanthroline, 1–4
Co-ordinative interactions in chelated complexes of silicon. Part 7. Dynamic nuclear magnetic resonance studies of fluorine exchange at five-co-ordinated silicon, 5–8
The crystal structure and electronic properties of the complex acetatobis(1,10-phenanthroline)copper(II) perchlorate dihydrate, acetatobis(1,10-phenanthroline)copper(II) nitrate dihydrate, and acetatobis(1,10-phenanthroline)zinc(II) tetrafluoroborate dihydrate, 141–50
The red form of bis(1,10-phenanthroline)dithiocyanatoiron(II), 991–6
Magnetic circular dichroism spectra of tris-chelate complexes of 2,2′-bipyridyl and 1,10-phenanthroline with iron(II), ruthenium(II), and osmium(II) at 4.2 K, 1781–8
New photoreduction catalysis by $[Cu(N\text{–}N)(PPh_3)_2]^+$ (N–N = 2,9-dimethyl-1,10-phenanthroline or 4,4′,6,6′-tetramethyl-2,2′-bipyridine) and its application to cobalt(III) complexes, 1959–62
Influence of decreasing solvent polarity (dioxane–water mixtures) on the stability and structure of binary and ternary complexes of adenosine 5′-triphosphate and uridine 5′-triphosphate, 2291–304
Lewis-base adducts of Group 1B metal(I) compounds. Part 18. Stereochemistries and structures of the 1:1 neutral complexes of Cu^IX with 1,10-phenanthroline (X = I) or 2,9-dimethyl-1,10-phenanthroline (X = I, Br, or Cl), 2531–40
Lewis-base adducts of Group 1B metal(I) compounds. Part 19. Crystal structures of bis(1,10-phenanthroline)copper(I) perchlorate and dibromocuprate(I), 2541–6
PHENOL
Reactivity of the unsaturated anion decacarbonyltetra-μ-hydrido-trirhenate(1−) toward phenols. Crystal and molecular structures of the tetraethylammonium salts of the triangular cluster anion $[Re_3(\mu\text{-}H)_3(\mu\text{-}OC_6F_5)(CO)_{10}]^-$ and of the binuclear anion $[Re_2(\mu\text{-}OC_6H_5)_3(CO)_6]^-$, 1507–12
PHENOLATE
Crystal structures and magnetic properties of binuclear five-co-ordinate copper(II) complexes with a phenolate bridge and their catalytic functions in multielectron redox reactions, 1945–52
PHENYL
Chemical and structural aspects of silver–triphenylarsine complexes and silver–tin complex salts, 321–32
Insertion of NO into transition metal–aryl bonds: formation of zirconium complexes containing the $[ON(R)NO]^-$ (R = Ph or *p*-MeC_6H_4) ligand, 405–8
The structure of pentachlorocyclopentadienylphenylmercury(II) in the solid state and in solution; the effect of *X*-ray absorption and crystal decay on molecular geometry parameters, 471–4
Investigation into diphosphine oxides as ligands in diorganotin(IV) adducts. Part 3. Synthesis and crystal structure of two adducts of dinitratodiphenyltin(IV) with *cis*- and *trans*-1,2-bis(diphenylphosphoryl)ethylene, 487–92
Organotin biocides. Part 2. Variable-temperature ^{119}Sn Mössbauer study of phenyl-and cyclohexyl-tin compounds, 1417–24
PHENYLAZOPYRIDINE
Mono- and bi-nuclear hydroxamates of bis(2-phenylazopyridine)ruthenium(II), 361–8
PHENYLHYDRAZONE
Mercurated and tellurated Schiff bases and phenylhydrazones, 821–6
PHOSPHAPENTANEDITHIOLATE
Molybdenum, rhenium, and tungsten complexes with bi- and tri-dentate phosphinothiolato-ligands; structures of $[Mo\{PhP(CH_2CH_2S)_2\}_2]$ and $[Mo(NNMe_2)\{PhP(CH_2CH_2S)_2\}_2]$, 2647–54
PHOSPHATE
Crystalline zirconium(IV) hydrogenarsenate hydrogenphosphate monohydrate: synthesis, ion-exchange properties, and thermal behaviour, 1737–42
Thermodynamic treatment of exchange of H^+ and K^+ in α-titanium phosphate, 1865–8
PHOSPHIDE
Cluster chemistry. Part 32. Synthesis and *X*-ray crystal structure of $[Ru_3(\mu_3\text{-}\eta^2\text{-}C_2PPh_2\text{-}P)(\mu\text{-}PPh_2)(CO)_{13}]$, a complex containing an alkynyl ligand in extended interaction with an open Ru_5 cluster, 1229–34
Reaction between $[Ru_3(CO)_{12}]$ and the tridentate phosphine ligand $HC(PPh_2)_3$; *X*-ray crystal structures of complexes $[Ru_3(CO)_9\{Ph_2PCHP(Ph)C_6H_4PPh\}]$, $[Ru_2H(CO)_4(Ph_2PCHPPh_2)\{PhPC_6H_4C(O)\}]$, and $[Ru_2(CO)_4Cl(PPh_2)(dppm)]$, 1835–42
PHOSPHINE
Reaction of bis-μ-diethylphosphido-bis(tetracarbonylmetal) (*M*–*M*) (M = Cr or W) with tri-n-butylphosphine: kinetics and mechanism of a reaction involving seven-co-ordinate complexes, 91–8
Lewis-base adducts of Group 1B metal(I) compounds. Part 13. Crystal structure determinations of tetrakis(triphenyl-phosphine)-copper(I) and -silver(I) perchlorates, bis(pyridine)bis(triphenylphosphine)copper(I) perchlorate, (2,2′-bipyridyl)bis(triphenylphosphine)copper(I) perchlorate, and tetrahydroboratobis(triphenylphosphine)copper(I)–pyridine (1/0.5), 125–34
The co-ordination chemistry of manganese. Part 14. Synthesis of manganese(II) complexes of tertiary phosphine ligands containing 2-cyanoethyl groups, 135–40
Insertion of carbon dioxide, of CO_2-like molecules, and of other unsaturated compounds into the platinum–nitrogen bond of $[Pt(PPh_3)_2(PhNO)]$, 163–8
Hexakis(trimethylphosphine)molybdenum chemistry: dinitrogen, ethylene, butadiene, η-cyclopentadienyl, and related derivatives, 423–34
Trimethylphosphine polyhydrides of tungsten and rhenium, 587–90
Electrochemically induced ligand substitutions on $[OsCl_3(PMe_2Ph)_3]$: rational pathways to osmium(II) complexes, 947–52

PHOSPHINE (contd)
New isomers of $[Os_3(CO)_{10}(PMe_2Ph)_2]$ and $[Os_3(CO)_9(PMe_2Ph)_3]$, 1037–42
Reaction of trithiazyl trichloride, $(NSCl)_3$, with triphenylphosphine or triphenylphosphine metal complexes. *X*-Ray crystal structure of aminotriphenylphosphonium chloride–dichloromethane (1/1), $[Ph_3PNH_2]Cl{\cdot}CH_2Cl_2$, 1043–8
Fluorophosphine complexes of rhodium(I) and iridium(I): towards the design of systems with extended metal–metal interactions. The crystal structure of $[\{IrCl(PF_3)_2\}_2]$, 1295–302
Preparation and properties of *cis*-$[W(N_2)_2(Ph_2PCH_2PPh_2)_2]$ and *trans*-$[M(N_2)_2(Ph_2PCH{=}CHPPh_2)_2]$ (M = Mo or W), and the crystal structure of *cis*-$[W(N_2)_2(PMe_2Ph)_4]$, 1523–6
Synthesis and structural characterisation of some *triangulo* platinum clusters containing isocyanide ligands, 1693–8
Iodine-127 Mössbauer spectroscopy of copper(I) iodide–phosphine adducts, 1727–30
Studies of gold cluster compounds using high-resolution ^{31}P solid-state nuclear magnetic resonance spectroscopy, 1811–4
The chemistry and catalytic properties of ruthenium and osmium complexes. Part 1. Homogeneous catalysis of organic reactions by bromo(carbonyl)hydridotris(triphenylphosphine)osmium(II), 1859–64
Isolation and crystal structure of $[Rh(PPh_3)_3(MeCN)][BF_4]$, acetonitriletris(triphenylphosphine)rhodium(I) tetrafluoroborate, 1977–80
Trimethylphosphine as a reactive solvent: synthesis, crystal structures, and reactions of $[Ta(PMe_3)_3(\eta^2\text{-}CH_2PMe_2)(\eta^2\text{-}CHPMe_2)]$ and $[W(PMe_3)_4(\eta^2\text{-}CH_2PMe_2)H]$ and related studies, 2025–36
Complexes of the platinum metals. Part 30. Fragmentation reactions of rhodium and iridium trichloro- and tribromo-acetates, 2113–20
On the synthesis of dimethylphenylphosphine complexes of manganese(II) halides, 2203–4
Synthesis and *X*-ray crystal structure of the asymmetric trinuclear complex $[Ni_3(\mu_3\text{-}S)_2(H_2O)(PPh_3)_5][PF_6]_2$, 2209–12
Co-ordinatively unsaturated diene complexes of tungsten(II) and their reactions with nucleophiles to give six- and seven-co-ordinate derivatives, 2231–8
Reactivity of $[NBu_4]_2[Mo_2Br_6]$ with several uni- and poly-dentate phosphines. *X*-Ray structure of $[NBu_4][MoBr_4(Ph_2PCH_2CH_2PPh_2)]$, 2263–8
The palladium-catalysed reaction between $[Re_2(CO)_{10}]$ and phosphines and the crystal and molecular structure of diaxial $[Re_2(CO)_8(PMe_2Ph)_2]$, 2277–82
Reactivity of $[NiR(R')L_2]$ compounds and the crystal structure of $[Ni(C_2Cl_3)(C_6H_2Me_3\text{-}2,4,6)(PMe_2Ph)_2]$, 2333–42
The chemistry of heteroarylphosphorus compounds. Part 16. Unusual substituent effects on selenium-77 nuclear magnetic resonance chemical shifts of heteroaryl- and aryl-phosphine selenides. *X*-Ray crystal structure of tri(2-furyl)phosphine selenide, 2505–8
Cationic complexes of ruthenium-(II) and -(III) with uni- and poly-dentate ligands, 2603–8
Novel ring compounds of bidentate phosphines with gold(I). Two-, three-, and four-co-ordination, 2655–60
The co-ordination of small molecules by manganese(II) phosphine complexes. Part 3. The dependence on the nature of the halogen in $[MnX_2(PR_3)]$ (X = Cl, Br, or I; R_3 = $PhMe_2$, $PhEt_2$, $PhPr^n_2$, $PhBu^n_2$, $PhBu^t_2$, Pr^n_3, Bu^n_3, or Ph_3) on adduct formation with sulphur dioxide in the solid state and in tetrahydrofuran solution, 2661–4

PHOSPHINE IMIDE
Determination of the molecular structures of tri(t-butyl)phosphine oxide and tri(t-butyl)phosphine imide in the gas phase by electron diffraction, 827–30

PHOSPHINE OXIDE
Determination of the molecular structures of tri(t-butyl)phosphine oxide and tri(t-butyl)phosphine imide in the gas phase by electron diffraction, 827–30

PHOSPHINETRIYLTRIPROPIONATE
Trigonal bipyramidal penta-aquazinc(II): crystal structure of penta-aquazinc(II) bis(3,3′,3″-phosphinetriyltripropionato)dizincate(II,II) heptahydrate, 2393–6

PHOSPHINOTHIOITE
Synthesis, solution structure, and halide addition compounds of bis(dicyclohexylphosphinothioito)mercury(II), 223–4

PHOSPHITE
Reactions of co-ordinated ligands. Part 33. Mononuclear η^2-vinyl complexes: synthesis, structure, and reactivity, 435–50
Complexes of the platinum metals. Part 26. Multinuclear nuclear magnetic resonance studies on rhodium(II) carboxylate adducts, 629–34
Reactions of some decaosmium clusters with electrophilic and nucleophilic reagents: *X*-ray structure analyses of $[N(PPh_3)_2][Os_{10}C(CO)_{24}(\mu\text{-}I)]$, $[Os_{10}C(CO)_{24}(\mu\text{-}I)_2]$, $[N(PPh_3)_2]_2[Os_{10}C(CO)_{22}(NO)I]$, $[Os_{10}C(CO)_{23}\{P(OMe)_3\}(\mu\text{-}I)_2]$ and of two isomers of $[Os_{10}C(CO)_{21}\{P(OMe)_3\}_4]$, 1795–810

PHOSPHOCHOLINE
Phosphocholine-substituted 5,10,15,20-tetraphenylporphyrinatoiron(II): oxygen carrier under physiological conditions, 275–8

PHOSPHONATE
Preparation and crystal structure of $Na_4[Hg_4(\mu\text{-}Cl)_4\{P(O)(OEt)_2\}_8]{\cdot}6H_2O$, 1061–4

PHOSPHONIUM
Reaction of trithiazyl trichloride, $(NSCl)_3$, with triphenylphosphine or triphenylphosphine metal complexes. *X*-Ray crystal structure of aminotriphenylphosphonium chloride–dichloromethane (1/1), $[Ph_3PNH_2]Cl{\cdot}CH_2Cl_2$, 1043–8

PHOSPHORUS
Studies of phosphazenes. Part 21. Associative and dissociative pathways in the aminolysis reactions of halogenocyclotriphosphazenes, 285–90
Phosphorus–phosphorus bond cleavage in the cage molecule P_4S_3: synthesis and crystal structure of the trinuclear platinum complex $[\{Pt(\mu\text{-}P_4S_3)(PPh_3)\}_3]{\cdot}C_6H_6$, 291–6
Determination of the molecular structure of difluorophosphine sulphide by the combined analysis of data from electron diffraction, microwave spectroscopy, and liquid crystal nuclear magnetic resonance spectroscopy, 755–60
Determination of the molecular structures of tri(t-butyl)phosphine oxide and tri(t-butyl)phosphine imide in the gas phase by electron diffraction, 827–30
Multinuclear nuclear magnetic resonance study of the interaction of some phosphorus(V) compounds with inorganic acids. The protonating abilities of HNO_3, $MeSO_3H$, and HPO_2F_2 towards the phosphoryl group, 865–6
Diastereoisomeric organophosphorus compounds. Part 4. Proton and phosphorus-31 nuclear magnetic resonance spectra of compounds of the type $[RR'P(X)]_2Y$ and $RR'P(X)\text{-}Y\text{-}(Z)PRR'$ (R = CH_3, R′ = t-C_4H_9) (1984, 2803), 871–2
A photoelectron spectroscopic study of di-t-butylphosphazene, 879–84
Studies of phosphazenes. Part 22. High-field nuclear magnetic resonance investigation of novel isomeric oxophosphazadienes, 1431–4
Mass spectrometric studies on cyclo- and poly-phosphazenes. Part 2. Oligomerization of hexa(aryloxy)cyclotriphosphazatrienes, 1547–54
Intermediates in the photochemical reaction of tetraphosphorus trisulphide with organic disulphides; phosphorus-31 nuclear magnetic resonance parameters for 2,6-bis(alkylthio)- and 2,6-diiodo-3,5,7-trithia-1,2,4,6-tetraphosphabicyclo[2.2.1]heptanes and for 3,6-bis(alkylthio)-2,5,7-trithia-1,3,4,6-tetraphosphabicyclo[2.2.1]heptanes, 1707–12
Studies of phosphazenes. Part 25. Synthesis, nuclear magnetic resonance spectroscopy, and mode of formation of (aziridino)(triphenylphosphazenyl)cyclotriphosphazenes. *X*-ray crystal structure and enzyme-inhibiting activity of $N_3P_3(NPPh_3)(NC_2H_4)_5$, 1881–90
Studies of phosphazenes. Part 21. Associative and dissociative pathways in the aminolysis reactions of halogenocyclotriphosphazenes (1985, 285), 2459–60

PHOSPHORUS-31
Synthesis, solution structure, and halide addition compounds of bis(dicyclohexylphosphinothioito)mercury(II), 223–4
Lewis-base adducts of group 1B metal(I) compounds. Part 16. Synthesis, structure, and solid-state phosphorus-31 nuclear magnetic resonance spectra of some novel $[Cu_4X_4L_4]$ (X = halogen, L = N,P base) 'cubane' clusters, 831–8
Diastereoisomeric organophosphorus compounds. Part 4. Proton and phosphorus-31 nuclear magnetic resonance spectra of compounds of the type $[RR'P(X)]_2Y$ and $RR'P(X)\text{-}Y\text{-}(Z)PRR'$ (R = CH_3, R′ = t-C_4H_9) (1984, 2803), 871–2
Intermediates in the photochemical reaction of tetraphosphorus

PHOSPHORUS-31 (contd)
trisulphide with organic disulphides; phosphorus-31 nuclear magnetic resonance parameters for 2,6-bis(alkylthio)- and 2,6-di-iodo-3,5,7-trithia-1,2,4,6-tetraphosphabicyclo[2.2.1]heptanes and for 3,6-bis(alkylthio)-2,5,7-trithia-1,3,4,6-tetraphosphabicyclo[2.2.1]heptanes, 1707–12
Studies of gold cluster compounds using high-resolution ^{31}P solid-state nuclear magnetic resonance spectroscopy, 1811–4
PHOSPHORYL
Multinuclear nuclear magnetic resonance study of the interaction of some phosphorus(v) compounds with inorganic acids. The protonating abilities of HNO_3, $MeSO_3H$, and HPO_2F_2 towards the phosphoryl group, 865–6
PHOTOCHEMISTRY
Light-induced electron-transfer reactions. Part 3. Kinetics of the oxidation reaction of oxalate ion by peroxodisulphate ion, induced by irradiation with visible light of an aqueous solution containing tris(2,2′-bipyridine)ruthenium(II), 355–60
Heteropolytungstates as catalysts for the photochemical reduction of oxygen and water, 395–400
Photochemistry of manganese porphyrins. Part 9. Redox reactions photosensitised by diamagnetic metalloporphyrins, 503–10
Absorption and emission in tris(2,2′-bipyridyl)ruthenium(II); effects of excited-state asymmetry, 1081–6
Photochemistry of dicarbonyl(η^5-cyclopentadienyl)-methyl- and -ethyl-iron and -ruthenium complexes in solutions at −30 °C and in frozen gas matrices at 12 K, 1365–74
Water photolysis. Part 1. The photolysis of co-ordinated water in $[\{MnL(H_2O)\}_2][ClO_4]_2$ (L = dianion of tetradentate O_2N_2-donor Schiff bases). A model for the manganese site in photosystem II of green plant photosynthesis, 1391–8
Intermediates in the photochemical reaction of tetraphosphorus trisulphide with organic disulphides; phosphorus-31 nuclear magnetic resonance parameters for 2,6-bis(alkylthio)- and 2,6-di-iodo-3,5,7-trithia-1,2,4,6-tetraphosphabicyclo[2.2.1]heptanes and for 3,6-bis(alkylthio)-2,5,7-trithia-1,3,4,6-tetraphosphabicyclo[2.2.1]heptanes, 1707–12
New photoreduction catalysis by $[Cu(N–N)(PPh_3)_2]^+$ (N–N = 2,9-dimethyl-1,10-phenanthroline or 4,4′,6,6′-tetramethyl-2,2′-bipyridine) and its application to cobalt(III) complexes, 1959–62
Preparation and characterisation of 2,2′-bipyridine-4,4′-disulphonic and -5-sulphonic acids and their ruthenium(II) complexes. Excited-state properties and excited-state electron-transfer reactions of ruthenium(II) complexes containing 2,2′-bipyridine-4,4′-disulphonic acid or 2,2′-bipyridine-4,4′-dicarboxylic acid, 2247–62
Some alkyldiplatinum(I) complexes and studies of the photochemical and thermal decomposition of the $[Pt_2Et(\mu\text{-}Ph_2PCH_2PPh_2)_2(Ph_2PCH_2PPh_2\text{-}P)]^+$ cation, 2421–6
Photochemical studies of the alkylammonium molybdates. Part 7. Octahedral sites for multi-electron reduction of $[Mo_8O_{26}(MoO_4)_2]^{8-}$, 2585–90
PHOTOELECTRON
Application of photoelectron spectroscopy to molecular properties. Part 19. Electronic structure of tris(α-di-imino) complexes of ruthenium(0), 43–50
Mono-η-cycloheptatrienyltitanium chemistry: synthesis, molecular and electronic structures, and reactivity of the complexes $[Ti(\eta\text{-}C_7H_7)L_2X]$ (L = tertiary phosphine, O- or N-donor ligand; X = Cl or alkyl), 669–84
A photoelectron spectroscopic study of di-t-butylphosphazene, 879–84
Formation of oxidized molybdenum-bearing ferrites, $Fe_{2.95-x}Mo_xO_4$ (x = 0.03—0.30) in aqueous suspensions by air oxidation and valence state of molybdenum ions in the lattice, 1713–16
Photoelectron spectra of the aminodifluoroboranes NH_2BF_2, $NHMeBF_2$, and NMe_2BF_2, 1767–70
He I and He II photoelectron spectra of open-chain pentadienyl complexes of manganese and rhenium, 2677–82
PHOTOSYNTHESIS
Water photolysis. Part 1. The photolysis of co-ordinated water in $[\{MnL(H_2O)\}_2][ClO_4]_2$ (L = dianion of tetradentate O_2N_2-donor Schiff bases). A model for the manganese site in photosystem II of green plant photosynthesis, 1391–8
PHOTOSYSTEM II
Water photolysis. Part 1. The photolysis of co-ordinated water in $[\{MnL(H_2O)\}_2][ClO_4]_2$ (L = dianion of tetradentate O_2N_2-donor Schiff bases). A model for the manganese site in photosystem II of green plant photosynthesis, 1391–8
PHTHALOCYANINATE
Synthesis and physico-chemical properties of cationic derivatives of phthalocyaninatocopper(II), 651–8
Equilibrium and kinetic study of pyridine binding to phthalocyaninatoiron(II) in dimethyl sulphoxide, 1107–12
Equilibrium and kinetic study of the reaction between phthalocyaninatoiron(II) and carbon monoxide in dimethyl sulphoxide in the presence of pyridine. Evidence for the formation of a transient, 1113–8
PHTHALOCYANINE
Partially oxidized group 3B fluorometallophthalocyanines, 269–74
PIPERAZINE
Synthesis and characterisation of uranium(IV) nitrate complexes with piperazines, 1985–8
PLATINACYCLOBUTANONE
Chemistry of metallacyclobutanones. Part 3. Reactions of heptane-2,4,6-trione and 1,5-diphenylpentane-1,3,5-trione with some carbonate complexes of platinum(II); X-ray crystal structures of $[Pt\{CH(COMe)COCH(COMe)\}(PPh_3)_2]$ and $[Pt\{OC(CHCOPh)CHC(Ph)O\}(PPh_3)_2]$, 549–54
PLATINUM
The kinetics of the displacement, by chloride, of heterocyclic nitrogen bases (am) from *trans*-$[PtL(am)Cl_2]$ (L = C_2H_4, CO, or PMe_3). The effect of steric hindrance in the leaving group on the *trans* effect of L, 27–30
Insertion of carbon dioxide, of CO_2-like molecules, and of other unsaturated compounds into the platinum–nitrogen bond of $[Pt(PPh_3)_2(PhNO)]$, 163–8
Chemistry of the unsaturated cluster compound $[Os_3Pt(\mu\text{-}H)_2(CO)_{10}\{P(cyclo\text{-}C_6H_{11})_3\}]$; X-ray crystal structures of $[Os_3Pt(\mu\text{-}H)_2(CO)_{11}\{P(cyclo\text{-}C_6H_{11})_3\}]$, $[Os_3Pt(\mu\text{-}H)_4(CO)_{10}\{P(cyclo\text{-}C_6H_{11})_3\}]$, and $[Os_3Pt(\mu\text{-}H)_2(\mu\text{-}CH_2)(CO)_{10}\{P(cyclo\text{-}C_6H_{11})_3\}]$ (two isomers), 177–90
Reactions of sulphoxide–thioether bidentate ligands with platinum(II). Determination of the mode of binding by cyclic voltammetry, 209–12
Optically active co-ordination compounds. Part 43. Polysulphide complexes of platinum(IV), 253–8
Bimetallic systems. Part 7. Platinum and palladium dicyanides containing terminal or bridging $Ph_2PCH_2PPh_2$ and heterobimetallics with silver, gold, mercury, rhodium, iridium, or molybdenum, 279–84
Phosphorus–phosphorus bond cleavage in the cage molecule P_4S_3: synthesis and crystal structure of the trinuclear platinum complex $[\{Pt(\mu\text{-}P_4S_3)(PPh_3)\}_3]\cdot C_6H_6$, 291–6
Static and dynamic nuclear magnetic resonance studies of complexes of trimethylplatinum(IV) halides with olefinic thio-and seleno-ethers. X-Ray crystal structures of $[PtXMe_3(MeSeCH{=}CHSeMe)]$ (X = Cl or I), 345–54
Nitrosoarene complexes of ruthenium and platinum and their reactions with NO and NO^+, 401–4
Chemistry of metallacyclobutanones. Part 3. Reactions of heptane-2,4,6-trione and 1,5-diphenylpentane-1,3,5-trione with some carbonate complexes of platinum(II); X-ray crystal structures of $[Pt\{CH(COMe)COCH(COMe)\}(PPh_3)_2]$ and $[Pt\{OC(CHCOPh)CHC(Ph)O\}(PPh_3)_2]$, 549–54
Synthesis, spectroscopy, and structure of the mixed-valence complexes $[Pt^{II}(en)_2][Pt^{IV}(en)_2X_2][Pt^{III}_2(H_2P_2O_5)_4X_2]$ (en = 1,2-diaminoethane, X = Br or I), 579–86
Electrochemistry of clusters. Part 4. Redox behaviour of tetrametallic clusters $[M^1_2M^2_2(\eta^5\text{-}C_5H_5)_2(\mu_3\text{-}CO)_2(\mu\text{-}CO)_4(PR_3)_2]$ (M^1 = Pt or Pd; M^2 = Cr, Mo, or W; R = Me, Et, Bu^n, or Ph), 711–6
Solvent extraction of gold and platinum-group metals using 2-nonylpyridine 1-oxide, and the crystal and molecular structure of bis(2-nonylpyridine 1-oxide)hydrogen(1+) tetrachloroaurate(III), 771–6
Platinum(II) complexes of nitroimidazoles: synthesis, characterisation, and X-ray crystal structures of *cis*-dichlorobis[1-(2′-hydroxyethyl)-2-hydroxymethyl-5-nitroimidazole]platinum(II) and *trans*-dichlorobis[1-(2′-hydroxy-3′-methoxypropyl)-2-nitroimidazole]platinum(II), 795–802
Interconversion of 42- and 44-electron platinum *triangulo*-clusters using chelating tertiary phosphine ligands and the structural characterisation of [1,3-bis(diphenylphosphino)propane]-tris(μ-sulphur dioxide)bis(tricyclohexylphosphine)triplatinum–benzene

PLATINUM (contd)
(1/2), $[Pt_3(\mu\text{-}SO_2)_3\{P(C_6H_{11})_3\}_2(dppp)]\cdot 2C_6H_6$, 845–50
Chemistry of platinum sulphido-complexes. Part 5. Synthesis and crystal and molecular structure of 3-(η-cyclo-octa-1,5-diene)bis(μ_3-sulphido)-1,1,2,2-tetrakis(triphenylphosphine)diplatinum(II)rhodium(I) hexafluorophosphate–dichloromethane (1/1), $[Pt_2Rh(\mu_3\text{-}S)_2(PPh_3)_4(\eta\text{-}C_8H_{12})]PF_6\cdot CH_2Cl_2$, 851–6
Reactions of 6,6′-bis(*nido*-decaboranyl) oxide and 6-hydroxy-*nido*-decaborane with dihalogenobis(phosphine) complexes of nickel, palladium, and platinum, and some related chemistry; nuclear magnetic resonance investigations and the crystal and molecular structures of bis(dimethylphosphine)-di-μ-(2,3,4-η^3-*nido*-hexaboranyl)-diplatinum(*Pt–Pt*), $[Pt_2(\mu\text{-}\eta^3\text{-}B_6H_9)_2(PMe_2Ph)_2]$, and of 2,4-dichloro-1,1-bis(dimethylphenylphosphine)-*closo*-1-nickeladecaborane, $[(PhMe_2P)_2NiB_9H_7Cl_2]$, 953–72
Bimetallic systems. Part 10. Synthesis of complexes of type $[(RC{\equiv}C)P^{\cdot}(\mu\text{-}dppm)_2Pt(C{\equiv}CR)]$ (dppm = $Ph_2PCH_2PPh_2$, R = Ph or *p*-tolyl) and their corresponding 'A frames' $[(RC{\equiv}C)Pt(\mu\text{-}dppm)_2(\mu\text{-}H)Pt(C{\equiv}CR)]Cl$ or $[(RC{\equiv}C)Pt(\mu\text{-}dppm)_2(\mu\text{-}X)Pt(C{\equiv}R)]$ with X = CS_2 or $MeOOCC{\equiv}CCOOMe$, 1015–8
Platinum(II) complexes containing a cationic amine ligand: crystal structure of [(2-aminoethyl)ammonium]trichloroplatinum(II), 1057–60
Synthesis, properties, and multinuclear magnetic resonance (^1H, ^{77}Se, and ^{195}Pt) studies on diselenoether complexes of palladium, platinum, and rhodium, 1265–70
Kinetic studies of oxidative dealkylation of alkylcobalamins by hexachloroplatinate(IV), 1375–80
Synthetic and nuclear magnetic resonance studies on dialkyl- and diaryl-platinum complexes containing chelating, monodentate, or bridging $Ph_2PCH_2PPh_2$ ligands, 1501–6
Synthesis and structural studies of some 1,1′-dichloroferrocene derivatives of platinum(II). Crystal and molecular structure of 2,2-μ-[(1—2,5—6-η-*cis*,*cis*-cyclo-octa-1,5-diene)platinio]-bis(1,1′-dichloroferrocene), 1527–30
Bimetallic systems. Part 11. Heterobimetallic and unsymmetrical diplatinum complexes from *cis*-$[PtR_2(dppm\text{-}P)_2]$ (dppm = $Ph_2PCH_2PPh_2$; R = Me, 1-naphthyl, or C_6H_4Me-*o*): crystal structure of $[(C_6H_4Me\text{-}o)_2Pt(\mu\text{-}dppm)_2PtMe_2]$, 1677–82
Synthesis and structural characterisation of some *triangulo*-platinum clusters containing isocyanide ligands, 1693–8
Copper(II) salts of metal dithiolates, 1731–2
Chemistry of di- and tri-metal complexes with bridging carbene or carbyne ligands. Part 37. Methylene group transfer to carbon–metal multiple bonds; crystal structures of $[TiW\{\mu\text{-}C(C_6H_4Me\text{-}4){=}CH_2\}(\mu\text{-}CO)(CO)(\eta\text{-}C_5H_5)_3]$ and $[PtW\{\mu\text{-}C(C_6H_4Me\text{-}4){=}CH_2\}(CO)_2(PMe_3)_2(\eta\text{-}C_5H_5)]$, 2009–16
The mechanisms of the reactions between hydrido-complexes and diazonium salts. Part 1. The reactions of *trans*-$[PtH(Cl)(PEt_3)_2]$, 2059–66
Bimetallic systems. Part 12. Mixed rhodium(I)–platinum(II) acetylide complexes containing bridging $Ph_2PCH_2PPh_2$. Crystal structures of $[(MeC{\equiv}C)Pt(\mu\text{-}dppm)_2(\sigma,\eta\text{-}C{\equiv}CMe)Rh(CO)]PF_6$ and of $[ClPt(\mu\text{-}dppm)_2(\sigma,\eta\text{-}C{\equiv}CMe)Rh(CO)]PF_6$, 2121–30
Bimetallic systems. Part 13. Platinum–manganese carbonyl complexes containing bridging $Ph_2PCH_2PPh_2$(dppm) ligands: crystal structure of $[(OC)_3Mn(\mu\text{-}dppm)_2PtH(Br)]BF_4$, 2131–8
Synthesis, properties, and ^{77}Se nuclear magnetic resonance studies of platinum metal complexes of two isomeric tris(selenoethers), $MeC(CH_2SeMe)_3$ and $Se(CH_2CH_2CH_2SeMe)_2$, 2185–90
Some alkyldiplatinum(I) complexes and studies of the photochemical and thermal decomposition of the $[Pt_2Et(\mu\text{-}Ph_2PCH_2PPh_2)_2(Ph_2PCH_2PPh_2\text{-}P)]^+$ cation, 2421–6
Synthesis of tetra- and penta-nuclear platinum–osmium carbido-cluster complexes from non-carbido-precursors: *X*-ray structural studies on $[Os_3Pt(\mu\text{-}H)_2(\mu_4\text{-}C)(CO)_{10}\{P(cyclo\text{-}C_6H_{11})_3\}]$, $[Os_3Pt_2(\mu\text{-}H)_2(\mu_5\text{-}C)(\mu\text{-}CO)(CO)_9\{P(cyclo\text{-}C_6H_{11})_3\}_2]$, and $[Os_3Pt_2(\mu\text{-}H)(\mu_5\text{-}C)(\mu\text{-}OMe)(\mu\text{-}CO)(CO)_9\{P(cyclo\text{-}C_6H_{11})_3\}_2]$, 2437–48
Fragmentation of co-ordinated carbon disulphide and dithiocarbene ligands by nucleophilic attack at carbon: the crystal structure of $[(Ph_3P)IPt(\mu\text{-}SMe)(\mu\text{-}CSMe)PtI(PPh_3)]\cdot Me_2CO$, 2595–602
PLATINUM-195
Platinum(II) complexes of nitroimidazoles: synthesis, characterisation, and *X*-ray crystal structures of *cis*-dichlorobis[1-(2′-hydroxyethyl)-2-hydroxymethyl-5-nitroimidazole)platinum(II) and *trans*-dichlorobis[1-(2′-hydroxy-3′-methoxypropyl)-2-nitroimidazole]platinum(II), 795–802
POLYETHER
Co-ordination of alkali metals by open-chain polyethers in transition metal complexes. Part 4. Variation in alkali-metal ion selectivity in cobalt and zinc complexes of 1-(*o*-carboxymethoxyphenoxy)-2-(*o*-hydroxyphenoxy)ethane (HL) and the *X*-ray and molecular structure of $[Zn(NH_4L)_2]$, 459–64
POLYIODIDE
Reaction of bis(morpholinothiocarbonyl) disulphide with iodine. Existence of a 1:1 charge-transfer precursory adduct in an oxidation reaction. Isolation and crystal structure of bis[3,5-di(*N*-morpholinio)-1,2,4-trithiolane] hexadecaiodide, 1349–54
POLYMER
Electron spin resonance spectra of manganese(II) ions in the double-stranded chain polymers $CdLX_2$ (L = pyridine or methylpyridine; X = Cl or Br), 75–80
Sulphur ligand–metal complexes. Part 16. Copper complexes of thioethers and the single-crystal *X*-ray structure of the polymeric mixed-valence complex, penta-μ-chloro-tris-μ-tetrahydrothiophene-tetracopper(I,II), 151–8
Characterization of the adducts formed by Cu(CN) and Cu(NCS) with biquinoline. The crystal structure of the polymeric cyano-compound containing both linear and tetrahedrally co-ordinated copper(I), $[\{Cu_3(bq)_2(CN)_3\}_n]$, 1285–8
The concentration and selective extraction of copper(II), rhodium(III), and iridium(III) using a copolymer functionalised with dithiocarbamate groups. Spectroscopic evidence for the nature of the binding sites, 1655–60
POLYPHOSPHANE
Macrocyclic polyphosphane ligands. Cobalt(II) and nickel(II) complexes of the γ and ε diastereoisomers of 4,7,13,16-tetraphenyl-4,7,13,16-tetraphospha-1,10-dithiacyclo-octadecane (L^1) and the crystal structure of $[Co(\varepsilon\text{-}L^1)][BPh_4]_2\cdot EtOH$, 1425–30
POLYSULPHIDE
Optically active co-ordination compounds. Part 43. Polysulphide complexes of platinum(IV), 253–8
PORPHYRIN
Rate parameters for oxygen and carbon monoxide binding to a liposome-embedded heme under physiological conditions, 65–8
Phosphocholine-substituted 5,10,15,20-tetraphenylporphyrinatoiron(II): oxygen carrier under physiological conditions, 275–8
Photochemistry of manganese porphyrins. Part 9. Redox reactions photosensitised by diamagnetic metalloporphyrins, 503–10
Reactions of cobalt(II) protoporphyrin IX dimethyl ester, $[Co^{II}P]$, and $[Co^{III}P(Cl)]$ in co-ordinating aliphatic alcohols, 1095–102
A mechanistic study of the reaction of iron(III) porphyrins with imidazoles. Hydrogen bonding by the propionic acid side chains in hemin chloride, 2269–76
PORPHYRINATE
Dismutation of superoxide ion in an aprotic solvent by 5,10,15,20-tetra-*p*-tolylporphyrinatocobalt(II), 1513–6
PROCYCLIDINE
Sila-pharmaca. Part 32. Crystal and molecular structures of the (*R*)-enantiomer and the racemate of the antimuscarinic agent (cyclohexyl)phenyl[2-(pyrrolidin-1-yl)ethyl]silanol (sila-procyclidine), 1743–6
PROGRAM
SUPERQUAD: an improved general program for computation of formation constants from potentiometric data, 1195–200
PROLINE
The L-proline residue as a 'break-point' in metal–peptide systems, 535–40
Specific binding of the tyrosine residue in copper(II) complexes of Tyr-Pro-Gly-Tyr and Tyr-Gly-Pro-Tyr, 1201–6
PROPANOLATE
Crystal and molecular structures and magnetic properties of four new exchange-coupled copper(II) complexes derived from different 3-*N*,*N*-dialkylamino-1-propanols and pseudohalogens, 1243–8
PROPENYLYLIDENE
Carbon–carbon formation at di-iron centres. Part 2. Reactivity of $[Fe_2(CO)_6(\mu\text{-}COEt)\{\mu\text{-}C(R)C(R)H\}]$ complexes toward $MeOC(O)C{\equiv}CC(O)OMe$ (R = Ph) and $CF_3C{\equiv}CCF_3$ (R = Ph or H); *X*-ray crystal structures of $[Fe_2(CO)_5\{\mu\text{-}C(OEt)C[C(O)OMe]C[C(O)OMe]\}\{\mu\text{-}C(Ph)C(Ph)H\}]\cdot H_2O$ and $[Fe_2(CO)_6\{\mu\text{-}C(CF_3)C(CF_3)CHCHCH(OMe)\}]$, 1087–94

PROTONATION
Multinuclear nuclear magnetic resonance study of the interaction of some phosphorus(v) compounds with inorganic acids. The protonating abilities of HNO_3, $MeSO_3H$, and HPO_2F_2 towards the phosphoryl group, 865–6
The reaction of $[Os_3(\mu\text{-}H)_2(CO)_9L]$ (L = CO or PEt_3) with dimethylcyanamide, Me_2NCN: *X*-Ray crystal structure of $[Os_3(\mu\text{-}H)(\mu\text{-}NCHNMe_2)(CO)_{10}]$ and the reactions of this complex with acids, 1355–60
PULSE RADIOLYSIS
One-electron reduction of nickel(IV) oxime complexes, 2213–6
PURINE
The crystal structure of diaquadichlorodimethyltin(IV)–purine (1/4), 1271–4
PYRAZINE
Mixed-ligand complexes of trivalent lanthanides. Part 3. Complexes of heptafluorodimethyloctane-3,5-dione and pyrazine: syntheses and spectral studies, 2547–50
PYRAZOLATE
Spectroscopic and magnetic properties of cobalt(II) and nickel(II) clusters obtained from 1-(hydroxymethyl)-3,5-dimethylpyrazole. *X*-Ray structure of tetrakis[chloro(μ_3-3,5-dimethyl-*N*-oxymethylpyrazolato-N^2,μ_3-*O*)(ethanol)nickel(II)] 737–42
PYRAZOLE
Transition-metal co-ordination compounds of a novel aniline-based pyrazole derivative. *X*-Ray crystal structures of [*NN*-bis(3,5-dimethylpyrazol-1-ylmethyl)aminobenzene]-dichlorocobalt(II) and -dibromocopper(II), 1699–706
PYRAZOLYL
Co-ordination chemistry of dimethylgold(III). Synthesis, structural studies, and fluxional behaviour of complexes with polydentate ligands, 1183–90
Electrochemical and chemical properties of di-iodonitrosyl[tris(3,5-dimethylpyrazolyl)borato]molybdenum, $[Mo\{HB(Me_2pz)_3\}(NO)I_2]$, and related complexes, 1249–54
Unidentate *versus* symmetrically and unsymmetrically bidentate nitrate co-ordination in pyrazole-containing chelates. The crystal and molecular structures of (nitrato-*O*)[tris(3,5-dimethylpyrazol-1-ylmethyl)amine]copper(II) nitrate, (nitrato-*O*,*O'*)[tris(3,5-dimethylpyrazol-1-ylmethyl)amine]nickel(II)nitrate, and (nitrato-*O*)(nitrato-*O*,*O'*)[tris(3,5-dimethylpyrazol-1-ylmethyl)amine]cadmium(II), 2177–84
Copper complexes with quadridentate bis(pyrazolyl)thioether amine and tris(pyrazolyl)amine ligands. Structural characterization of the complexes $[Cu(NCS)(tpea)][Cu(NCS)_2]$ and $[CuCl(bdma)]Cl\cdot 2H_2O$, 2327–32
PYRIDINE
The kinetics of the displacement, by chloride, of heterocyclic nitrogen bases (am) from *trans*-$[PtL(am)Cl_2]$ (L = C_2H_4, CO, or PMe_3). The effect of steric hindrance in the leaving group on the *trans* effect of L, 27–30
Electron spin resonance spectra of manganese(II) ions in the double-stranded chain polymers $CdLX_2$ (L = pyridine or methylpyridine; X = Cl or Br), 75–80
Lewis-base adducts of Group 1B metal(I) compounds. Part 13. Crystal structure determinations of tetrakis(triphenyl-phosphine)-copper(I) and -silver(I) perchlorates, bis(pyridine)bis(triphenylphosphine)copper(I) perchlorate, (2,2'-bipyridyl)bis(triphenylphosphine)copper(I) perchlorate, and tetrahydroboratobis(triphenylphosphine)copper(I)–pyridine (1/0.5), 125–34
Dinuclear mono-μ-chloro-pyridyldiaza rhodium(I) complexes derived from pyridyldi-imines *via* hydrogen transfer from ethanol, 1053–6
Equilibrium and kinetic study of pyridine binding to phthalocyaninatoiron(II) in dimethyl sulphoxide, 1107–12
Structure and solvation of mercury(II) iodide, bromide, and chloride in pyridine solution; refinement of the crystal structure of di-iodobis(pyridine)mercury(II), $[HgI_2(py)_2]$, 1597–604
Equilibria between mono- and bi-nuclear complexes in $Cu(O_2CMe)_2$–pyridine derivative–diluent systems. The influence of the amine ligand basicity, 1849–52
Studies on spin-equilibrium iron(III) complexes. Part 1. Syntheses and magnetic properties of a new family of spin cross-over iron(III) complexes with a unidentate ligand over a wide range of the spectrochemical series and a quinquedentate ligand derived from salicylaldehyde and di(3-aminopropyl)amine. *X*-Ray crystal structure of [4-azaheptamethylene-1,7-bis(salicylideneiminato)](4-methylpyridine)iron(III) tetraphenylborate, 2575–84
PYRIDINECARBOXYLATE
Electrophilic nitrosyls: preparation, structure, and reactivity of *cis*-chloronitrosylbis(pyridine-2-carboxylato)ruthenium and related complexes, 2427–32
PYRIDINEMETHANOL
Displacement of pyridine-2-methanol from dichloro(pyridine-2-methanolato)gold(III) in acidic solution. Ring opening at oxygen, 731–6
PYRIDINE OXIDE
Short hydrogen bonds: diadducts of substituted pyridine *N*-oxides: synthesis, spectroscopic studies, and *X*-ray structure, 749–54
PYRIDINETHIOLATE
Complexes of the platinum metals. Part 29. Pyridine-2-thiolate derivatives of ruthenium and osmium: *X*-ray crystal structures of $[Ru(C_5H_4NS)_2(CO)_2(PPh_3)]$ and $[Ru(C_5H_4NS)_2(CO)(PPh_3)]$, 2101–12
PYRIDOXYLIDENEGLYCINATE
Metal complexes of vitamin B_6 related compounds. Crystal and molecular structures of aqua(5'-phosphopyridoxylideneglycinato)copper(II) trihydrate and bis(pyridoxylideneglycinato)nickel(II) hexahydrate, 2051–8
PYRIDYL
Co-ordination chemistry of dimethylgold(III). Synthesis, structural studies, and fluxional behaviour of complexes with polydentate ligands, 1183–90
PYRIDYLAZORESORCINOL
Kinetics and mechanisms of complex formation of gallium(III) and indium(III). The reactions with 4-(2-pyridylazo)resorcinol in water and other mixed solvents, 2615–22
PYRROLIDINE
Reactions of tris[(2*S*)-2-(aminomethyl)pyrrolidine]nickel(II) ion with alk-3-en-2-ones or 4-hydroxyalkan-2-ones: formation of an optically active tetra-aza macrocycle, 2139–44
PYRROLOPYRIDINATE
1*H*-pyrrolo[2,3-*b*]pyridine (HL) ligands in rhodium(I) and iridium(I) chemistry. Crystal and molecular structures of $[Rh_2(\mu\text{-}L)_2(nbd)_2]$ and $[Rh_4(\mu\text{-}Cl)_2(\mu\text{-}L)_2(\mu\text{-}CO)_2(CO)_2(nbd)_2]$, 1891–8
RADICAL CATION
Structure and relative energies of some nitrogen-containing radical cations by MNDO calculation, 1915–22
RADIOLYSIS
Kinetics and mechanism of single electron oxidations of the tervalent uranium ion, U^{3+}(aq), by free radicals in aqueous solutions, 641–4
RAMAN
Synthesis, spectroscopy, and structure of the mixed-valence complexes $[Pt^{II}(en)_2][Pt^{IV}(en)_2X_2][Pt^{III}_2(H_2P_2O_5)_4X_2]$ (en = 1,2-diaminoethane, X = Br or I), 579–86
Spectrochemistry of solutions. Part 16. A Raman spectroscopic study of the complexation of mercury(II) by cyanide ligands in liquid ammonia at 293 K, 663–8
Palladium(II,IV) mixed-valence complexes of 1,2-diaminoethane, 1,3-diaminopropane, and diethylenetriamine: syntheses, electronic, infrared, Raman, and resonance Raman spectra and *X*-ray studies, 815–20
Preparation and vibrational spectra of $[OsX_6]^{3-}$ (X = Cl, Br, or I) and of other platinum-group hexahalogeno-complexes, 1673–6
Fundamental vibrations of six- and seven-membered selenium sulphide ring molecules, 1869–76
RARE EARTH
Preparation, characterization, and physical properties of the series MPd_3S_4 (M = rare earth), 2369–74
REDOX POTENTIAL
Solvent effects on the redox potential of the uranium(VI)–uranium(V) couple, 601–4
REDUCTION
Heteropolytungstates as catalysts for the photochemical reduction of oxygen and water, 395–400
The synthesis of $NR_4[(Ta_6Cl_{12})(H_2O)_6]X_4$ (R = Me or Et, X = Cl or Br) by the spontaneous reduction of $[Ta_6Cl_{12}]^{4+}$ to $[Ta_6Cl_{12}]^{3+}$ in acidic media: *X*-ray structure analysis of $NMe_4[(Ta_6Cl_{12})(H_2O)_6]Br_4$, 455–8
Preparation of 1,3,2-dithiazolium hexafluoroarsenate(v), preparation and crystal structures of 5-methyl-1,3,2,4-dithiadiazolium and 4-methyl-1,3,2-dithiazolium hexafluoroarsenate(v) and the reduction of these salts to stable free radicals, 1405–16
Bimetallic cyano-bridged cations: preparation and hydride reduction of $[(\eta^5\text{-}C_5H_5)L_2Ru(\mu\text{-}CN)ML'_2(\eta^5\text{-}C_5H_5)]PF_6$ [L_2, L'_2

REDUCTION
= $(PPh_3)_2$, $Ph_2PCH_2CH_2PPh_2$; M = Ru or Fe]. Formation of $[Ru(\eta^5\text{-}C_5H_5)(PPh_3)H_3]$ and *X*-ray crystal structure of $[(\eta^5\text{-}C_5H_5)(Ph_2PCH_2CH_2PPh_2)Ru(\mu\text{-}CN)Ru(PPh_3)_2(\eta^5\text{-}C_5H_5)]PF_6$, 1479–86
The synthesis of $NR_4[(Ta_6Cl_{12})(H_2O)_6]X_4$ (R = Me or Et, X = Cl or Br) by the spontaneous reduction of $[Ta_6Cl_{12}]^{4+}$ to $[Ta_6Cl_{12}]^{3+}$ in acidic media: *X*-ray structure analysis of $NMe_4[(Ta_6Cl_{12})(H_2O)_6]Br_4$ (1985, 455), 1531–2
Kinetics and mechanism of the reductions of tris(oxalato)cobaltate(III) ion by ruthenium(II) species in aqueous solution, 1665–8
A new reaction of nitrosyl complexes; one-electron reduction of *trans*-$[MX(NO)L_4]^{2+}$ [M = Ru or Fe, X = Cl or Br, L = pyridine or *o*-phenylenebis(dimethylarsine)] with hydroxylamine, 1733–4
The kinetics and stoicheiometry of silver(III) reduction by the octacyano-complexes of molybdenum(IV) and tungsten(IV), 1789–94
New photoreduction catalysis by $[Cu(N\text{–}N)(PPh_3)_2]^+$ (N–N = 2,9-dimethyl-1,10-phenanthroline or 4,4′,6,6′-tetramethyl-2,2′-bipyridine) and its application to cobalt(III) complexes, 1959–62
The easy catalytic reduction of $HClO_4$, HNO_3, and H_5IO_6 by CO in the presence of $[Pd_3(O_2CMe)_6]$, 1971–4
One-electron reduction of nickel(IV) oxime complexes, 2213–6

RELAXATION
Carbon-13 nuclear magnetic resonance evidence of a relaxation process dominated by scalar coupling with a quadrupolar nucleus in $[Re_3(\mu\text{-}H)_4(CO)_{10}]^-$, 1899–902

RESOLUTION
Optical resolution of DL-2,2′-bipiperidine through its cobalt(III) complex, 895–8

RESORCINOL
Coloured species formed from the titanium(IV)-4-(2′-pyridylazo)resorcinol reagent in the spectrophotometric determination of trace amounts of hydrogen peroxide, 81–4

RHENADECABORANE
Polyhedral rhenaborane chemistry: crystal and molecular structures of the *nido*-6-rhenadecaborane cluster compounds [6,6,6,6-$(PMe_2Ph)_3$H-*nido*-6-ReB_9H_{13}] and [2-(PMe_2Ph)-6,6,6,6-$(PMe_2Ph)_2$ClH-*nido*-6-ReB_9H_{12}]; nuclear magnetic resonance parameters of these and other related *nido*-rhenadecaborane cluster species, 1119–30

RHENIUM
Trimethylphosphine polyhydrides of tungsten and rhenium, 587–90
Reaction of transition-metal carbonylate anions and 1,1,1-tris(halogenomethyl)ethane. *X*-Ray crystal structures of tricarbonyl(η^5-cyclopentadienyl)(1-methylcyclopropylmethyl)tungsten(II), and tetraethylammonium enneacarbonyliodidorhenate(O), 931–40
The preparation and properties of some diphosphines $R_2PCH_2CH_2PR_2$ (R = alkyl or aryl) and of their rhenium(I) dinitrogen derivatives, 1131–6
Reactivity of the unsaturated anion decacarbonyltetra-μ-hydrido-trirhenate(1−) toward phenols. Crystal and molecular structures of the tetraethylammonium salts of the triangular cluster anion $[Re_3(\mu\text{-}H)_3(\mu\text{-}OC_6F_5)(CO)_{10}]^-$ and of the binuclear anion $[Re_2(\mu\text{-}OC_6H_5)_3(CO)_6]^-$, 1507–12
Rhenium nitrosyl complexes with simple and with sterically demanding aromatic thiolate ligands: *X*-ray crystal structures of $[PPh_4][Re_2(SC_6H_4Me\text{-}4)_7(NO)_2]\cdot CH_2Cl_2$ and $[Re(SC_6H_3Pr^i_2\text{-}2,6)_4(NO)]$, 1533–42
Chemistry of di- and tri-metal complexes with bridging carbene or carbyne ligands. Part 36. Reactions of the dimetal compounds $[ReM(\equiv CC_6H_4Me\text{-}4)(CO)_9]$ (M = Cr, Mo, or W) with octacarbonyldicobalt; crystal structures of $[Co_2WRe(\mu_3\text{-}CC_6H_4Me\text{-}4)(CO)_{15}]$ and $[Co_2Re(\mu_3\text{-}CC_6H_4Me\text{-}4)(CO)_{10}]$, 2001–8
Trimethylphosphine as a reactive solvent: synthesis, crystal structures, and reactions of $[Ta(PMe_3)_3(\eta^2\text{-}CH_2PMe_2)(\eta^2\text{-}CHPMe_2)]$ and $[W(PMe_3)_4(\eta^2\text{-}CH_2PMe_2)H]$ and related studies, 2025–36
Preparation and properties of *mer*-$[ReCl(N_2)(CNR)\{P(OMe)_3\}_3]$ (R = Me, Et, Bu^t, C_6H_4Me-4, or C_6H_4Cl-4) and $[ReCl(N_2)(CNMe)(PPh_3)\{P(OEt)_3\}_2]$. *X*-Ray crystal structure of *mer*-$[ReCl(N_2)(CNMe)\{P(OMe)_3\}_3]$ and reductive cleavage of the isocyanide ligands to primary amines upon protonation, 2079–84
Oxoalkyls of rhenium-(V) and -(VI). *X*-Ray crystal structures of $(Me_4ReO)_2Mg(thf)_4$, $[(Me_3SiCH_2)_4ReO]_2Mg(thf)_2$, $Re_2O_3Me_6$, and $Re_2O_3(CH_2SiMe_3)_6$, 2167–76
The palladium-catalysed reaction between $[Re_2(CO)_{10}]$ and phosphines and the crystal and molecular structure of diaxial $[Re_2(CO)_8(PMe_2Ph)_2]$, 2277–82
Rhenium nitrido-, arylimido-, nitrile, and carbonyl complexes with sterically hindered thiolate ligands, 2305–10
Nitrogen-15 nuclear magnetic resonance relaxation mechanisms in dinitrogen complexes of molybdenum, tungsten, rhenium, and osmium, 2473–8
Molybdenum, rhenium, and tungsten complexes with bi- and tridentate phosphinothiolato-ligands; structures of $[Mo\{PhP(CH_2CH_2S)_2\}_2]$ and $[Mo(NNMe_2)\{PhP(CH_2CH_2S)_2\}_2]$, 2647–54
He I and He II photoelectron spectra of open-chain pentadienyl complexes of manganese and rhenium, 2677–82

RHODIUM
Carbene complexes. Part 18. Synthetic routes to electron-rich olefin-derived monocarbenerhodium(I) neutral and cationic complexes and their chemical and physical properties (1984, 2355), 229–30
Bimetallic systems. Part 8. Heterobimetallic di-isonitrile or isonitrile–carbonyl complexes of rhodium or iridium bridged to silver or gold by $Ph_2PCH_2PPh_2$, 511–6
Complexes of the platinum metals. Part 23. Synthesis of the nitrosyl carboxylate complexes $[M(O_2CR)_2(NO)(PPh_3)_2]$ (M = Rh or Ir; R = CF_3, C_2F_5, or C_6F_5): crystal and molecular structures of the trifluoroacetate derivatives $[M(O_2CCF_3)_2(NO)(PPh_3)_2]$, 611–6
Complexes of the platinum metals. Part 24. The role of dioxygen in the reactions of trifluoroacetic acid with the rhodium and iridium nitrosyls $[M(NO)(PPh_3)_3]$, 617–20
Complexes of the platinum metals. Part 26. Multinuclear nuclear magnetic resonance studies on rhodium(II) carboxylate adducts, 629–34
Synthesis and *X*-ray structural studies on the cluster compounds $[RuRh_3(\mu_3\text{-}CO)_2(CO)_3(\eta\text{-}C_5Me_5)_3]$ and $[RuRh_2(\mu\text{-}CO)(\mu_3\text{-}CO)(CO)_2(\eta^4\text{-}C_8H_{10})(\eta\text{-}C_5Me_5)_2]$, 645–50
Studies on transition-metal cyano-complexes. Part 4. Cyanide hydride complexes of Groups 6A and 8, 717–22
Chemistry of platinum sulphido-complexes. Part 5. Synthesis and crystal and molecular structure of 3-(η-cyclo-octa-1,5-diene)bis(μ_3-sulphido)-1,1,2,2-tetrakis(triphenylphosphine)diplatinum(II)rhodium(I) hexafluorophosphate–dichloromethane (1/1), $[Pt_2Rh(\mu_3\text{-}S)_2(PPh_3)_4(\eta\text{-}C_8H_{12})]PF_6\cdot CH_2Cl_2$, 851–6
Correlation between structure and circular dichroism. Structure and absolute configuration of the (−) isomer of lithium (ethylenediamine-*N*,*N*′-diacetato-*N*,*N*′-di-3-propionato)rhodate(III) pentahydrate, 861–4
Pyrazolate A-frame rhodium complexes. Crystal structures of $[Rh_2(\mu\text{-}dmpz)(CO)_2(\mu\text{-}dppm)_2][ClO_4]$ and $[Rh_2(\mu\text{-}dmpz)I_2(CO)_2(\mu\text{-}dppm)_2][ClO_4]$, 973–80
Dinuclear mono-μ-chloro-pyridyldiaza rhodium(I) complexes derived from pyridyldi-imines *via* hydrogen transfer from ethanol, 1053–6
Synthesis, properties, and multinuclear magnetic resonance (^1H, ^{77}Se, and ^{195}Pt) studies on diselenoether complexes of palladium, platinum, and rhodium, 1265–70
Fluorophosphine complexes of rhodium(I) and iridium(I): towards the design of systems with extended metal–metal interactions. The crystal structure of $[\{IrCl(PF_3)_2\}_2]$, 1295–302
New carbide clusters in the cobalt sub-group. Part 15. Synthesis and crystallographic characterization of di-μ_6-carbido-deca-μ-carbonyl-tridecacarbonyl-*polyhedro*-dodecarhodate(4−) as its tetrapropylammonium salt, $[N(C_3H_7)_4]_4[Rh_{12}C_2(CO)_{23}]$, 1309–14
Chemistry of di- and tri-metal complexes with bridging carbene or carbyne ligands. Part 33. Reactions of $[W(\equiv CMe)(CO)_2(\eta\text{-}C_5H_5)]$ with the dimetal compounds $[MRh(\mu\text{-}CO)_2(\eta\text{-}C_5Me_5)_2]$ (M = Co or Rh); *X*-ray crystal structure of $[Rh_2W(\mu\text{-}CO)(\mu_3\text{-}CMe)(CO)_2(\eta\text{-}C_5H_5)(\eta\text{-}C_5Me_5)_2]$, 1315–22
Benzamidinatorhodium complexes. *X*-Ray structures of $[Rh\{CPh(NPh)_2\}(cod)]$ and $[Rh_2\{\mu\text{-}CPh(NPh)_2\}_2(tfbb)_2]$, 1487–94
Mono- and di-nuclear rhodium and palladium complexes of macrocyclic ligands containing the 2,6-di(thiomethyl)pyridine sub-unit, 1517–22
The syntheses and characterisation of, and the determination of

RHODIUM (contd)
$^1J(^{103}Rh-^{103}Rh)$ in $[(\eta^5\text{-}C_5Me_5)_2Rh_2(\mu\text{-}CH_2)_2\{\mu\text{-}CH_2CR(CH_2CR{=}CH_2)CH_2\}]$ (R = H or Me) and $[(\eta^5\text{-}C_5Me_5)_2Rh_2(\mu\text{-}CH_2)_2\{\mu\text{-}Ph_2P(CH_2)_nPPh_2\}]^{2+}$ (n = 1 or 2), 1555–60
Synthesis of bis[μ-bis(diphenylphosphino)methane]-tri-μ-carbonyl-tricarbonyl-*triangulo*-trirhodium(1 +) perchlorate, $[Rh_3(CO)_3(\mu\text{-}CO)_3(\mu\text{-}dppm)_2]ClO_4$. An unusual 46-electron cluster possessing the A-frame structure with a bridging $Rh(CO)_3$ fragment, 1577–84
Oxygenation studies. Part 7. Catalytic dioxygenation of cyclo-octa-1,5-diene at a rhodium centre, 1591–6
The concentration and selective extraction of copper(II), rhodium(III), and iridium(III) using a copolymer functionalised with dithiocarbamate groups. Spectroscopic evidence for the nature of the binding sites, 1655–60
Preparation and vibrational spectra of $[OsX_6]^{3-}$ (X = Cl, Br, or I) and of other platinum-group hexahalogeno-complexes, 1673–6
Dirhodium(II,II) tetra-acetate complexes with axially co-ordinated triphenylstibine, triphenylarsine, and dibenzyl sulphide ligands. The syntheses, properties, and *X*-ray crystal structures of $[Rh_2(O_2CMe)_4(SbPh_3)_2]$, $[Rh_2(O_2CMe)_4(AsPh_3)_2]$, and $[Rh_2(O_2CMe)_4\{S(CH_2Ph)_2\}_2]$, 1775–80
1*H*-pyrrolo[2,3-*b*]pyridine (HL) ligands in rhodium(I) and iridium(I) chemistry. Crystal and molecular structures of $[Rh_2(\mu\text{-}L)_2(nbd)_2]$ and $[Rh_4(\mu\text{-}Cl)_2(\mu\text{-}L)_2(\mu\text{-}CO)_2(CO)_2(nbd)_2]$, 1891–8
Isolation and crystal structure of $[Rh(PPh_3)_3(MeCN)][BF_4]$, acetonitriletris(triphenylphosphine)rhodium(I) tetrafluoroborate, 1977–80
The mechanisms of the reactions between hydrido-complexes and diazonium salts. Part 2. The reactions of *cis,mer*-$[RhHCl_2(PEtPh_2)_3]$, 2067–78
Complexes of the platinum metals. Part 30. Fragmentation reactions of rhodium and iridium trichloro- and tribromo-acetates, 2113–20
Bimetallic systems. Part 12. Mixed rhodium(I)–platinum(II) acetylide complexes containing bridging $Ph_2PCH_2PPh_2$. Crystal structures of $[(MeC{\equiv}C)Pt(\mu\text{-}dppm)_2(\sigma,\eta\text{-}C{\equiv}CMe)Rh(CO)]PF_6$ and of $[ClPt(\mu\text{-}dppm)_2(\sigma,\eta\text{-}C{\equiv}CMe)Rh(CO)]PF_6$, 2121–30
Stereochemical consequences of the two-electron oxidation of a dirhodium fulvalene complex: the *X*-ray crystal structures of *trans*-$[Rh_2(CO)_2(PPh_3)_2(\eta^5\text{:}\eta'^5\text{-}C_{10}H_8)]$ and *cis*-$[Rh_2(CO)_2(PPh_3)_2(\eta^5\text{:}\eta'^5\text{-}C_{10}H_8)][PF_6]_2$, 2283–90
Reactions of co-ordinated ligands. Part 35. Evidence for carbon–carbon double-bond cleavage of cyclopropenes in their reaction with dinuclear cobalt, rhodium, and iridium complexes; crystal structure and protolysis of $[Rh_2(\mu\text{-}CO)(\mu\text{-}COCHCMe_2CH)(\mu\text{-}C_5Me_5)_2]$, 2483–92

RING OPENING
Reduction–oxidation properties of organotransition-metal complexes. Part 20. Oxidative and thermolytic cyclopropane ring-opening reactions; *X*-ray crystal structure of $[Fe_2(CO)_6(\eta^4\text{:}\eta'^4\text{-}C_{14}H_{18})]$, 699–706
Annelation of ring-opened arylcyclopropenium ions to co-ordinated cyclo-octatetraene, and the *X*-ray crystal structure of $[Fe(CO)_3(\sigma,\eta^3\text{-}C_{11}H_9Ph_3)]$, 777–82

RUTHENABORANE
Polyhedral ruthenaborane chemistry: characterization of several new ruthenaboranes by nuclear magnetic resonance spectroscopy, and the crystal and molecular structure of [5,6,6-$(PPh_3)_3$-6-H-*nido*-6-RuB_9H_{12}], 2397–406
Two unusual *closo*-type ruthenaboranes: preparation, molecular structure, and nuclear magnetic resonance properties of [1,1,1-$(PPh_3)HCl$-1-RuB_9H_7-3,5-$(PPh_3)_2$] and [1,1-$(PPh_3)_2$-1-$RuB_{10}H_8$-2,5-$(OEt)_2$], 2407–16

RUTHENIUM
Application of photoelectron spectroscopy to molecular properties. Part 19. Electronic structure of tris(α-di-imino) complexes of ruthenium(0), 43–50
Light-induced electron-transfer reactions. Part 3. Kinetics of the oxidation reaction of oxalate ion by peroxodisulphate ion, induced by irradiation with visible light of an aqueous solution containing tris(2,2′-bipyridine)ruthenium(II), 355–60
Mono- and bi-nuclear hydroxamates of bis(2-phenylazopyridine)ruthenium(II), 361–8
Nitrosoarene complexes of ruthenium and platinum and their reactions with NO and NO^+, 401–4
Complexes of the platinum metals. Part 25. Reactions of ruthenium and osmium nitrosyls $[MH(NO)(PPh_3)_3]$ and $[M(NO)_2(PPh_3)_2]$ with perfluorocarboxylic acids: *X*-ray crystal structure determination of nitrosyl(trifluoroacetato)-(trifluoroacetohydroximato-*OO*′)bis(triphenylphosphine)osmium(II)–dichloromethane (1/1), 621–8
Synthesis and *X*-ray structural studies on the cluster compounds $[RuRh_3(\mu_3\text{-}CO)_2(CO)_3(\eta\text{-}C_5Me_5)_3]$ and $[RuRh_2(\mu\text{-}CO)(\mu_3\text{-}CO)(CO)_2(\eta^4\text{-}C_8H_{10})(\eta\text{-}C_5Me_5)_2]$, 645–50
Organoruthenium(II) complexes formed by insertion reactions of some vinyl compounds and conjugated dienes into a hydrido–ruthenium bond, 873–8
Mechanism of addition of aryltrimethyl-silanes and -stannanes to tricarbonyl(cyclohexadienyl)ruthenium(II) cation, 1049–52
Absorption and emission in tris(2,2′-bipyridyl)ruthenium(II); effects of excited-state asymmetry, 1081–6
Cluster chemistry. Part 31. An $[Ru_3(CO)_{12}]$ derivative with an $[Fe_3(CO)_{12}]$-like structure: preparation and *X*-ray structure of $[Ru_3(\mu\text{-}CO)_2(CO)_6\{PPh(OMe)_2\}_4]$, 1223–8
Cluster chemistry. Part 32. Synthesis and *X*-ray crystal structure of $[Ru_5(\mu_5\text{-}\eta^2\text{-}C_2PPh_2\text{-}P)(\mu\text{-}PPh_2)(CO)_{13}]$, a complex containing an alkynyl ligand in extended interaction with an open Ru_5 cluster, 1229–34
From diarylruthenium complexes to *ortho*-metallated ketones: a mechanistic and crystal structure study, 1235–42
Photochemistry of dicarbonyl(η^5-cyclopentadienyl)-methyl- and -ethyl-iron and -ruthenium complexes in solutions at −30 °C and in frozen gas matrices at 12 K, 1365–74
Bimetallic cyano-bridged cations: preparation and hydride reduction of $[(\eta^5\text{-}C_5H_5)L_2Ru(\mu\text{-}CN)ML'_2(\eta^5\text{-}C_5H_5)]PF_6$ [L_2, L'_2 = $(PPh_3)_2$, $Ph_2PCH_2CH_2PPh_2$; M = Ru or Fe]. Formation of $[Ru(\eta^5\text{-}C_5H_5)(PPh_3)H_3]$ and *X*-ray crystal structure of $[(\eta^5\text{-}C_5H_5)(Ph_2PCH_2CH_2PPh_2)Ru(\mu\text{-}CN)Ru(PPh_3)_2(\eta^5\text{-}C_5H_5)]PF_6$, 1479–86
Kinetics and mechanism of the reductions of tris(oxalato)cobaltate(III) ion by ruthenium(II) species in aqueous solution, 1665–8
Preparation and vibrational spectra of $[OsX_6]^{3-}$ (X = Cl, Br, or I) and of other platinum-group hexahalogeno-complexes, 1673–6
A new reaction of nitrosyl complexes; one-electron reduction of *trans*-$[MX(NO)L_4]^{2+}$ [M = Ru or Fe, X = Cl or Br, L = pyridine or *o*-phenylenebis(dimethylarsine)] with hydroxylamine, 1733–4
Magnetic circular dichroism spectra of tris-chelate complexes of 2,2′-bipyridyl and 1,10-phenanthroline with iron(II), ruthenium(II), and osmium(II) at 4.2 K, 1781–8
Reaction between $[Ru_3(CO)_{12}]$ and the tridentate phosphine ligand $HC(PPh_2)_3$; *X*-ray crystal structures of complexes $[Ru_3(CO)_9\{Ph_2PCHP(Ph)C_6H_4PPh\}]$, $[Ru_2H(CO)_4(Ph_2PCHPPh_2)\{PhPC_6H_4C(O)\}]$, and $[Ru_2(CO)_4Cl(PPh_2)(dppm)]$, 1835–42
Organic chemistry of dinuclear metal centres. Part 8. Organo-iron–ruthenium chemistry. *X*-Ray structure of *trans*-$[FeRu(CO)_2(\mu\text{-}CO)_2(\eta\text{-}C_5H_5)_2]$, 1935–44
Chemistry of di- and tri-metal complexes with bridging carbene or carbyne ligands. Part 38. Ruthenium–tungsten compounds: crystal structures of $[RuW(\mu\text{-}Cl)(\mu\text{-}CMe)(Cl)(CO)_2(PPh_3)_2(\eta\text{-}C_5H_5)]$ and $[RuW_2(\mu_3\text{-}C_2Me_2)(CO)_7(\eta\text{-}C_5H_5)_2]$, 2017–24
Trimethylphosphine as a reactive solvent: synthesis, crystal structures, and reactions of $[Ta(PMe_3)_3(\eta^2\text{-}CH_2PMe_2)(\eta^2\text{-}CHPMe_2)]$ and $[W(PMe_3)_4(\eta^2\text{-}CH_2PMe_2)H]$ and related studies, 2025–36
Complexes of the platinum metals. Part 29. Pyridine-2-thiolate derivatives of ruthenium and osmium: *X*-ray crystal structures of $[Ru(C_5H_4NS)_2(CO)_2(PPh_3)]$ and $[Ru(C_5H_4NS)_2(CO)(PPh_3)]$, 2101–12
Reactions of co-ordinated ligands. Part 34. Synthesis, structure, and reactivity of cationic dieneruthenium complexes; crystal structures of $[Ru(\eta^4\text{-}C_6H_9)(CO)(\eta\text{-}C_5H_5)][BF_4]$ and $[Ru(\eta^5\text{-}C_6H_9)(CO)(\eta\text{-}C_5H_5)]$, 2145–54
Preparation and characterisation of 2,2′-bipyridine-4,4′-disulphonic and -5-sulphonic acids and their ruthenium(II) complexes. Excited-state properties and excited-state electron-transfer reactions of ruthenium(II) complexes containing 2,2′-bipyridine-4,4′-disulphonic acid or 2,2′-bipyridine-4,4′-dicarboxylic acid, 2247–62
The synthesis, magnetic, electrochemical, and spectroscopic properties of diruthenium(II,II) tetra-μ-carboxylates and their adducts. *X*-Ray structures of $Ru_2(O_2CR)_4L_2$ (R = Me, L = H_2O or tetrahydrofuran; R = Et, L = Me_2CO), 2321–6

RUTHENIUM (contd)
Metallaborane chemistry. Part 14.[1] Icosahedral η^6-arene carbametallaboranes of iron and ruthenium; molecular structures of *closo*-[1-(η^6-C_6H_5Me)-2,4-Me_2-1,2,4-$FeC_2B_9H_9$] and *closo*-[3-(η^6-C_6H_6)-3,1,2-$RuC_2B_9H_{11}$], 2343–8
Polyhedral ruthenaborane chemistry: characterization of several new ruthenaboranes by nuclear magnetic resonance spectroscopy, and the crystal and molecular structure of [5,6,6-$(PPh_3)_3$-6-H-*nido*-6-RuB_9H_{12}], 2397–406
Two unusual *closo*-type ruthenaboranes: preparation, molecular structure, and nuclear magnetic resonance properties of [1,1,1-(PPh_3)HCl-1-RuB_9H_7-3,5-$(PPh_3)_2$] and [1,1-$(PPh_3)_2$-1-$RuB_{10}H_9$-2,5-$(OEt)_2$], 2407–16
Electrophilic nitrosyls: preparation, structure, and reactivity of *cis*-chloronitrosylbis(pyridine-2-carboxylato)ruthenium and related complexes, 2427–32
Thiazyl chloride complexes of ruthenium(II), 2449–52
Linkage isomerism in penta-ammineruthenium(II),(III) complexes of benzotriazole, 2469–72
Cationic complexes of ruthenium-(II) and -(III) with uni- and polydentate ligands, 2603–8
Ligand reactivity in polypyridine complexes; the deuteriation of the bis(2,2′:6′,2″-terpyridine)ruthenium(II) cation, 2687–90

SALICYLALDOXIME
Some water-soluble Schiff-base complexes of nickel(II): the role of water, 1451–4

SALICYLIDINEIMINE
Some water-soluble Schiff-base complexes of nickel(II): the role of water, 1451–4

SCHIFF BASE
Enantioselectivity of nickel(II) and copper(II) complexes of Schiff bases derived from amino acids and (*S*)-*o*-[(*N*-benzylprolyl)amino]acetophenone or (*S*)-*o*-[(*N*-benzylprolyl)amino]benzaldehyde. Crystal and molecular structures of [Ni{(*S*)-bap-(*S*)-Val}] and [Cu{(*S*)-bap-(*S*)-Val}], 17–26
Binuclear metal complexes. Part 55. Dimanganese(II) and mixed-valence manganese(II,III) complexes of 'strati-bis' Schiff bases with two salen-like donating sites, 59–64
Synthesis and characterization of copper(II) complexes with unsymmetrical quadridentate Schiff bases derived from β-diketone, diamine, and *o*-hydroxybenzaldehyde ligands, 101–6
Synthesis and characterization of a new family of binuclear copper(II) complexes with a flexible bridge and the π–π type charge-transfer interaction with 1,3,5-trinitrobenzene, 107–10
Reaction of some aliphatic diamines with four-co-ordinated unsymmetrical ketoenamine copper(II)and nickel(II) complexes, 803–6
Mercurated and tellurated Schiff bases and phenylhydrazones, 821–6
Water photolysis. Part 1. The photolysis of co-ordinated water in $[\{MnL(H_2O)\}_2][ClO_4]_2$ (L = dianion of tetradentate O_2N_2-donor Schiff bases). A model for the manganese site in photosystem II of green plant photosynthesis, 1391–8
Some water-soluble Schiff-base complexes of nickel(II): the role of water, 1451–4
Compartmental ligands. Part 11. Copper(II) complexes of 'half-unit' Schiff bases: crystal and molecular structures of one monomeric and of one dimeric complex, 1471–8
The structure of a dinuclear copper(I) complex of a Schiff-base ligand containing a copper–copper bond, 1771–4
Metal complexes of vitamin B_6 related compounds. Crystal and molecular structures of aqua(5′-phosphopyridoxylideneglycinato)copper(II) trihydrate and bis(pyridoxylideneglycinato)nickel(II) hexahydrate, 2051–8
X-Ray crystal structures and magnetic properties of azide-bridged binuclear copper(II) complexes containing the Schiff-base ligand derived from 2-pyridinecarbaldehyde and histamine. Structure–magnetism relationship, 2095–100
Studies on spin-equilibrium iron(III) complexes. Part 1. Syntheses and magnetic properties of a new family of spin cross-over iron(III) complexes with a unidentate ligand over a wide range of the spectrochemical series and a quinquedentate ligand derived from salicylaldehyde and di(3-aminopropyl)amine. *X*-Ray crystal structure of [4-azaheptamethylene-1,7-bis(salicylideneiminato)](4-methylpyridine)iron(III) tetraphenylborate, 2575–84

SELANE
Crystal structures and rotameric forms of some diarylsulphonyl-mono-, -di-, and -tri-selanes and their sulphur analogues, 231–8

SELENIDE
The chemistry of heteroarylphosphorus compounds. Part 16. Unusual substituent effects on selenium-77 nuclear magnetic resonance chemical shifts of heteroaryl- and aryl-phosphine selenides. *X*-Ray crystal structure of tri(2-furyl)phosphine selenide, 2505–8

SELENIUM
Crystal structures and rotameric forms of some diarylsulphonyl-mono-, -di-, and -tri-selanes and their sulphur analogues, 231–8
Structural and nuclear magnetic resonance studies of short selenium–nitrogen bonds, 565–72
The molecular structure of gaseous bis(trifluoromethyl)selenium difluoride as determined by electron diffraction, 941–6
Chemistry of di- and tri-metal complexes with bridging carbene or carbyne ligands. Part 34. Reactions of sulphur and selenium with the di-irontungsten complexes $[Fe_2W(\mu_3\text{-}CR)(\mu\text{-}CO)(CO)_8(\eta\text{-}C_5H_5)]$ (R = C_6H_4Me-4 or Me); crystal structures of $[Fe_2W(\mu\text{-}CC_6H_4Me\text{-}4)(\mu_3\text{-}S)(CO)_7(\eta\text{-}C_5H_5)]$ and $[Fe_2W(\mu_3\text{-}SCMe)(CO)_8(\eta\text{-}C_5H_5)]$, 1323–30
Fundamental vibrations of six- and seven-membered selenium sulphide ring molecules, 1869–76

SELENIUM-77
Structural and nuclear magnetic resonance studies of short selenium–nitrogen bonds, 565–72
Synthesis, properties, and multinuclear magnetic resonance (^1H, ^{77}Se, and ^{195}Pt) studies on diselenoether complexes of palladium, platinum, and rhodium, 1265–70
Synthesis, properties, and ^{77}Se nuclear magnetic resonance studies of platinum metal complexes of two isomeric tris(selenoethers), $MeC(CH_2SeMe)_3$ and $Se(CH_2CH_2CH_2SeMe)_2$, 2185–90
The chemistry of heteroarylphosphorus compounds. Part 16. Unusual substituent effects on selenium-77 nuclear magnetic resonance chemical shifts of heteroaryl- and aryl-phosphine selenides. *X*-Ray crystal structure of tri(2-furyl)phosphine selenide, 2505–8

SELENOETHER
Synthesis and properties of dioxo-osmium(VI) compounds of thio- and seleno-ethers, 205–8
Static and dynamic nuclear magnetic resonance studies of complexes of trimethylplatinum(IV) halides with olefinic thio-and seleno-ethers. *X*-Ray crystal structures of $[PtXMe_3(MeSeCH{=}CHSeMe)]$ (X = Cl or I), 345–54
Synthesis, properties, and ^{77}Se nuclear magnetic resonance studies of platinum metal complexes of two isomeric tris(selenoethers), $MeC(CH_2SeMe)_3$ and $Se(CH_2CH_2CH_2SeMe)_2$, 2185–90

SELF-ASSOCIATION
Self-association of organocobalamins in aqueous solution, 1381–6

SHIFT REAGENT
Mixed-ligand complexes of trivalent lanthanides. Part 3. Complexes of heptafluorodimethyloctane-3,5-dione and pyrazine: syntheses and spectral studies, 2547–50

SILANE
Mechanism of addition of aryltrimethyl-silanes and -stannanes to tricarbonyl(cyclohexadienyl)ruthenium(II) cation, 1049–52

SILICON
Co-ordinative interactions in chelated complexes of silicon. Part 5. Chiralty of five-co-ordinate silicon compounds: crystal and molecular structures of 1-(chlorodimethylsilyl)-and 1-(dichloromethylsilyl)-1,2,3,4-tetrahydro-1,10-phenanthroline, 1–4
Co-ordinative interactions in chelated complexes of silicon. Part 7. Dynamic nuclear magnetic resonance studies of fluorine exchange at five-co-ordinated silicon, 5–8
Sila–pharmaca. Part 32. Crystal and molecular structures of the (*R*)-enantiomer and the racemate of the antimuscarinic agent (cyclohexyl)phenyl[2-(pyrrolidin-1-yl)ethyl]silanol (sila-procyclidine), 1743–6
Syntheses of organofunctional siloxanes containing metal-ligating side-chains, 2191–4

SILOXANE
Syntheses of organofunctional siloxanes containing metal-ligating side-chains, 2191–4

SILVER
Lewis-base adducts of Group 1B metal(I) compounds. Part 11. Synthesis and crystal structure of adducts of silver(I) bromide with monomethyl-substituted pyridine bases, 111–6
Lewis-base adducts of Group 1B metal(I) compounds. Part 12. Structural studies of some bis(methyl-substituted pyridine)-copper(I) and -silver(I) nitrates and perchlorates, 117–24
Lewis-base adducts of Group 1B metal(I) compounds. Part 13.

SILVER (contd)
Crystal structure determinations of tetrakis(triphenyl-phosphine)-copper(I) and -silver(I) perchlorates, bis(pyridine)bis(triphenylphosphine)copper(I) perchlorate, (2,2′-bipyridyl)bis(triphenylphosphine)copper(I) perchlorate, and tetrahydroboratobis(triphenylphosphine)copper(I)–pyridine (1/0.5), 125–34
Chemical and structural aspects of silver–triphenylarsine complexes and silver–tin complex salts, 321–32
Bimetallic systems. Part 8. Heterobimetallic di-isonitrile or isonitrile–carbonyl complexes of rhodium or iridium bridged to silver or gold by $Ph_2PCH_2PPh_2$, 511–6
The kinetics and stoicheiometry of silver(III) reduction by the octacyano-complexes of molybdenum(IV) and tungsten(IV), 1789–94
Synthesis and cryptate complexes of azathia macropolycyclic ligands based on 12-membered N_2S_2 and 15-membered N_2S_3 macrocyclic subunits, 2311–8
SOLVATION
Structure and solvation of mercury(II) iodide, bromide, and chloride in pyridine solution; refinement of the crystal structure of di-iodobis(pyridine)mercury(II), $[HgI_2(py)_2]$, 1597–604
SOLVENT EFFECT
The peroxodisulphate–iodide reaction. Reactivity and ionic association and solvation in isodielectric water–solvent mixtures, 31–4
Solvent effects on the redox potential of the uranium(VI)–uranium(V) couple, 601–4
Solvent dependence of the stereochemistry of base-catalysed solvolysis of *trans*-$[Co(NH_3)_4(^{15}NH_3)X]^{3+/2+}$ ions in dipolar aprotic solvents, 659–62
Electrostatic solvent effect on the formation of the mixed-chelate complex (acetylacetonato)(diethyldithiocarbamato)-copper(II), 987–90
Electron spin resonance studies of iron(III) complexes of ethylenediaminetetra-acetate and *N*-(2-hydroxyethyl)ethylenediamine-*NN′N′*-triacetate in co-ordinating solvents, 1077–80
Beryllocene: A microwave dielectric loss study, 1761–6
The peroxodisulphate–hexacyanoferrate(II) reaction. Reactivity and ionic association in isodielectric water–co-solvent mixtures, 1975–6
The mechanisms of the reactions between hydrido-complexes and diazonium salts. Part 1. The reactions of *trans*-$[PtH(Cl)(PEt_3)_2]$, 2059–66
The mechanisms of the reactions between hydrido-complexes and diazonium salts. Part 2. The reactions of *cis,mer*-$[RhHCl_2(PEtPh_2)_3]$, 2067–78
Copper(II) complexes of *N,N′*-bis(2-carbamoylethyl)ethylenediamine in methanol–water, 2217–20
Influence of decreasing solvent polarity (dioxane–water mixtures) on the stability and structure of binary and ternary complexes of adenosine 5′-triphosphate and uridine 5′-triphosphate, 2291–304
The effect of a dansyl group on the co-ordinative ability of *N*-protected amino acids. Part 1. Behaviour of the copper(II) ion–*N*-dansylglycinate system in aqueous and methanolic solution, 2363–8
Kinetics and mechanisms of complex formation of gallium(III) and indium(III). The reactions with 4-(2-pyridylazo)resorcinol in water and other mixed solvents, 2615–22
SOLVENT EXTRACTION
Solvent extraction of gold and platinum-group metals using 2-nonylpyridine 1-oxide, and the crystal and molecular structure of bis(2-nonylpyridine 1-oxide)hydrogen(1+) tetrachloroaurate(III), 771–6
SOLVOLYSIS
Solvent dependence of the stereochemistry of base-catalysed solvolysis of *trans*-$[Co(NH_3)_4(^{15}NH_3)X]^{3+/2+}$ ions in dipolar aprotic solvents, 659–62
SPINEL
X-Ray diffraction determination of the cation distribution and oxygen positional parameter in polycrystalline spinels, 2155–60
STABILITY
Influence of secondary ligands on the stability of metal–xanthosine complexes in solution, 239–42
The L-proline residue as a 'break-point' in metal–peptide systems, 535–40
An equation for predicting the formation constants of hydroxo-metal complexes, 723–30
The stability of nickel(II) complexes of tetra-aza macrocycles, 1877–80
The hydrolysis of metal ions. Part 8. Aluminium(III), 1967–70
On the possibility of determining the thermodynamic parameters for the formation of weak complexes using a simple model for the dependence on ionic strength of activity coefficients: Na^+, K^+, and Ca^{2+} complexes of low molecular weight ligands in aqueous solution, 2353–62
The effect of a dansyl group on the co-ordinative ability of *N*-protected amino acids. Part 1. Behaviour of the copper(II) ion–*N*-dansylglycinate system in aqueous and methanolic solution, 2363–8
Carbon-13 nuclear magnetic resonance study of the complexes formed between zinc(II) and triethylenetetramine, 2381–6
STABILITY CONSTANT
SUPERQUAD: an improved general program for computation of formation constants from potentiometric data, 1195–200
Reactions of metal ions with triketones in solution. Part 1. Formation constants for the systems of heptane-2,4,6-trione, 1-phenylhexane-1,3,5-trione, 1,5-diphenylpentane-1,3,5-trione, and 2,2′-dihydroxybenzophenone with proton, nickel(II), and cobalt(II), 1543–6
Potentiometric study of the complex-formation equilibria of manganese(II), cobalt(II), nickel(II), copper(II), and zinc(II) with ethylenediamine-*N*-acetic acid, 1605–8
The chemistry of vitamin B_{12}. Part 25. Mechanism of the β-elimination of olefins from alkylcorrinoids; evidence for an initial homolytic fission of the Co–C bond, 1613–8
The chemistry of vitamin B_{12}. Part 26. Co-ordination of the malonitrile anion by Co^{III} corrinoids: first experimental determination of equilibrium constants for the co-ordination of a tetrahedral carbanion by a transition-metal ion, 1619–22
STACKING
Influence of decreasing solvent polarity (dioxane–water mixtures) on the stability and structure of binary and ternary complexes of adenosine 5′-triphosphate and uridine 5′-triphosphate, 2291–304
STANNANE
Mechanism of addition of aryltrimethyl-silanes and -stannanes to tricarbonyl(cyclohexadienyl)ruthenium(II) cation, 1049–52
STEEL
Electrochemical studies in HNO_3–N_2O_4 mixtures: corrosion of stainless steel in HNO_3–N_2O_4 mixtures and the effect of inhibitors, 2551–4
STEREOCHEMISTRY
Solvent dependence of the stereochemistry of base-catalysed solvolysis of *trans*-$[Co(NH_3)_4(^{15}NH_3)X]^{3+/2+}$ ions in dipolar aprotic solvents, 659–62
STEREOSELECTIVITY
Optical resolution of DL-2,2′-bipiperidine through its cobalt(III) complex, 895–8
STIBINE
Dirhodium(II,II) tetra-acetate complexes with axially co-ordinated triphenylstibine, triphenylarsine, and dibenzyl sulphide ligands. The syntheses, properties, and *X*-ray crystal structures of $[Rh_2(O_2CMe)_4(SbPh_3)_2]$, $[Rh_2(O_2CMe)_4(AsPh_3)_2]$, and $[Rh_2(O_2CMe)_4\{S(CH_2Ph)_2\}_2]$, 1775–80
STRONTIUM
Magnetic exchange interactions in perovskite solid solutions. Part 5. The unusual defect structure of $SrFeO_{3-y}$, 1455–70
SUBSTITUTION
The kinetics of the displacement, by chloride, of heterocyclic nitrogen bases (am) from *trans*-$[PtL(am)Cl_2]$ (L = C_2H_4, CO, or PMe_3). The effect of steric hindrance in the leaving group on the *trans* effect of L, 27–30
Ligand substitution on (*N,N*-dimethylformamide)[2,2′,2″-tri(*N,N*-dimethylamino)triethylamine]cobalt(II), 413–6
Displacement of pyridine-2-methanol from dichloro(pyridine-2-methanolato)gold(III) in acidic solution. Ring opening at oxygen, 731–6
Electrochemically induced ligand substitutions on $[OsCl_3(PMe_2Ph)_3]$: rational pathways to osmium(II) complexes, 947–52
SULPHANE
Crystal structures and rotameric forms of some diarylsulphonyl-mono-, -di-, and -tri-selanes and their sulphur analogues, 231–8
SULPHATE
Some water-soluble Schiff-base complexes of nickel(II): the role of water, 1451–4

SULPHIDE
Niobium(IV) sulphidohalides: preparation of $Nb_2X_4S_3$ and Nb_2X_4S-*n*L[X = Br or Cl; *n* = 4, L = NCMe, SMe_2, or tetrahydrothiophene (tht); *n* = 2, L = $PhSCH_2CH_2SPh$]. Crystal and molecular structure of $Nb_2Cl_4S_2$-4tht, 417–22
Chemistry of platinum sulphido-complexes. Part 5. Synthesis and crystal and molecular structure of 3-(η-cyclo-octa-1,5-diene)bis(μ$_3$-sulphido)-1,1,2,2-tetrakis(triphenylphosphine)diplatinum(II)rhodium(I) hexafluorophosphate–dichloromethane (1/1), $[Pt_2Rh(\mu_3\text{-}S)_2(PPh_3)_4(\eta\text{-}C_8H_{12})]PF_6\cdot CH_2Cl_2$, 851–6
Synthesis, structures, and reactivities of some pentamethylcyclopentadienyl–sulphur compounds, 1303–8
Reaction of bis(morpholinothiocarbonyl) disulphide with iodine. Existence of a 1:1 charge-transfer precursory adduct in an oxidation reaction. Isolation and crystal structure of bis[3,5-di(*N*-morpholinio)-1,2,4-trithiolane] hexadecaiodide, 1349–54
Dirhodium(II,II) tetra-acetate complexes with axially co-ordinated triphenylstibine, triphenylarsine, and dibenzyl sulphide ligands. The syntheses, properties, and *X*-ray crystal structures of $[Rh_2(O_2CMe)_4(SbPh_3)_2]$, $[Rh_2(O_2CMe)_4(AsPh_3)_2]$, and $[Rh_2(O_2CMe)_4\{S(CH_2Ph)_2\}_2]$, 1775–80
High-co-ordination-number compounds of niobium and tantalum: Reactions of niobium and tantalum halides and sulphido-halides with sodium diethyldithiocarbamate. The crystal structures of $[Nb(S_2CNEt_2)_4]Br$, $Nb(S_2CNEt_2)_3S$, and $Ta(S_2CNEt_2)_3(S_2)$, 1821–8
Crystal structure and spectroscopic and redox properties of the iron–sulphur cluster compound $[NEt_4]_2[Fe_4S_4(SC_6H_4NH_2\text{-}4)_4]$, 2161–6
Synthesis and *X*-ray crystal structure of the asymmetric trinuclear complex $[Ni_3(\mu_3\text{-}S)_2(H_2O)(PPh_3)_3][PF_6]_2$, 2209–12
Electrical and magnetic properties of $K_3Cu_8S_6$, 2319–20
Preparation, characterization, and physical properties of the series MPd_3S_4 (M = rare earth), 2369–74
Fragmentation of co-ordinated carbon disulphide and dithiocarbene ligands by nucleophilic attack at carbon: the crystal structure of $[(Ph_3P)IPt(\mu\text{-}SMe)(\mu\text{-}CSMe)PtI(PPh_3)]\cdot Me_2CO$, 2595–602
SULPHONIC ACID
Preparation and characterisation of 2,2′-bipyridine-4,4′-disulphonic and -5-sulphonic acids and their ruthenium(II) complexes. Excited-state properties and excited-state electron-transfer reactions of ruthenium(II) complexes containing 2,2′-bipyridine-4,4′-disulphonic acid or 2,2′-bipyridine-4,4′-dicarboxylic acid, 2247–62
SULPHONIUM
Crystal structure of tris(methylmercurio)sulphonium perchlorate, 2457–8
SULPHOXIDE
Reactions of sulphoxide–thioether bidentate ligands with platinum(II). Determination of the mode of binding by cyclic voltammetry, 209–12
The preparation of trichloro(phenyl 2-pyridylmethyl sulphoxide)gold(III) and the kinetics and equilibria of its formation and subsequent reactions, 2091–4
SULPHUR
Reactions in mixed non-aqueous systems containing sulphur dioxide. Part 6. The reaction of metal oxides with dimethyl sulphoxide–sulphur dioxide, 99–100
Crystal structures and rotameric forms of some diarylsulphonyl-mono-, -di-, and -tri-selanes and their sulphur analogues, 231–8
Phosphorus–phosphorus bond cleavage in the cage molecule P_4S_3: synthesis and crystal structure of the trinuclear platinum complex $[\{Pt(\mu\text{-}P_4S_3)(PPh_3)\}_3]\cdot C_6H_6$, 291–6
Zinc–sulphur bond enthalpy: its determination in bis(diethyldithiocarbamato)zinc(II), 369–72
Synthesis, structures, and reactivities of some pentamethylcyclopentadienyl–sulphur compounds, 1303–8
Chemistry of di- and tri-metal complexes with bridging carbene or carbyne ligands. Part 34. Reactions of sulphur and selenium with the di-irontungsten complexes $[Fe_2W(\mu_3\text{-}CR)(\mu\text{-}CO)(CO)_8(\eta\text{-}C_5H_5)]$ (R = C_6H_4Me-4 or Me); crystal structures of $[Fe_2W(\mu\text{-}CC_6H_4Me\text{-}4)(\mu_3\text{-}S)(CO)_7(\eta\text{-}C_5H_5)]$ and $[Fe_2W(\mu_3\text{-}SCMe)(CO)_9(\eta\text{-}C_5H_5)]$, 1323–30
Intermediates in the photochemical reaction of tetraphosphorus trisulphide with organic disulphides; phosphorus-31 nuclear magnetic resonance parameters for 2,6-bis(alkylthio)- and 2,6-diiodo-3,5,7-trithia-1,2,4,6-tetraphosphabicyclo[2.2.1]heptanes and for 3,6-bis(alkylthio)-2,5,7-trithia-1,3,4,6-tetraphosphabicyclo[2.2.1]heptanes, 1707–12
Fundamental vibrations of six- and seven-membered selenium sulphide ring molecules, 1869–76
SULPHUR DIOXIDE
Reactions in mixed non-aqueous systems containing sulphur dioxide. Part 6. The reaction of metal oxides with dimethyl sulphoxide–sulphur dioxide, 99–100
Interconversion of 42- and 44-electron platinum *triangulo*-clusters using chelating tertiary phosphine ligands and the structural characterisation of [1,3-bis(diphenylphosphino)propane]-tris(μ-sulphur dioxide)bis(tricyclohexylphosphine)triplatinum–benzene (1/2), $[Pt_3(\mu\text{-}SO_2)_3\{P(C_6H_{11})_3\}_2(dppp)]\cdot 2C_6H_6$, 845–50
The co-ordination of small molecules by manganese(II) phosphine complexes. Part 3. The dependence on the nature of the halogen in $[MnX_2(PR_3)]$ (X = Cl, Br, or I; R_3 = $PhMe_2$, $PhEt_2$, $PhPr^n_2$, $PhBu^n_2$, $PhBu^t_2$, Pr^n_3, Bu^n_3, or Ph_3) on adduct formation with sulphur dioxide in the solid state and in tetrahydrofuran solution, 2661–4
SUPEROXIDE
Dismutation of superoxide ion in an aprotic solvent by 5,10,15,20-tetra-*p*-tolylporphyrinatocobalt(II), 1513–6
SUPEROXO
A new electron spin resonance spectrum for an exchange- and dipole–dipole coupled superoxocobalamin · · · free-radical pair occurring in adenosylcobalamin-containing systems, 891–4
TANTALUM
The synthesis of $NR_4[(Ta_6Cl_{12})(H_2O)_6]X_4$ (R = Me or Et, X = Cl or Br) by the spontaneous reduction of $[Ta_6Cl_{12}]^{4+}$ to $[Ta_6Cl_{12}]^{3+}$ in acidic media: *X*-ray structure analysis of $NMe_4[(Ta_6Cl_{12})(H_2O)_6]Br_4$, 455–8
The synthesis of $NR_4[(Ta_6Cl_{12})(H_2O)_6]X_4$ (R = Me or Et, X = Cl or Br) by the spontaneous reduction of $[Ta_6Cl_{12}]^{4+}$ to $[Ta_6Cl_{12}]^{3+}$ in acidic media: *X*-ray structure analysis of $NMe_4[(Ta_6Cl_{12})(H_2O)_6]Br_4$ (1985, 455), 1531–2
Axially asymmetric metal alkyls. Part 3. Chemical, electrochemical, and structural studies of group 5A $d^{0,1,2}$ metallepines $[M\{(2\text{-}CH_2C_6H_4)_2\}(\eta\text{-}C_5H_5)_2]^z$ (M = Nb or Ta; *z* = 1−, 0, or 1+); synthesis of $[Nb\{2\text{-}CH_2(C_6H_4)_2Me\text{-}2'\}X(\eta\text{-}C_5H_5)_2]$ (X = Cl or CO) and $[\{(\eta\text{-}C_5H_5)_2ClV\}_2\{(2\text{-}CH_2C_6H_4)_2\}]$, and electrochemistry of $[M(CH_2Ph)_{2-n}Cl_n(\eta\text{-}C_5H_5)_2]$ (*n* = 0, M = Nb, or Ta; *n* = 1, M = V), 1747–60
High-co-ordination-number compounds of niobium and tantalum: Reactions of niobium and tantalum halides and sulphido-halides with sodium diethyldithiocarbamate. The crystal structures of $[Nb(S_2CNEt_2)_4]Br$, $Nb(S_2CNEt_2)_3S$, and $Ta(S_2CNEt_2)_3(S_2)$, 1821–8
New carbonyl derivatives of niobium(I) and tantalum(I), 1989–96
TELLURIUM
Mercurated and tellurated Schiff bases and phenylhydrazones, 821–6
Synthesis and crystal structure of the layer compound Sb_3TeO_6Cl, 1633–6
Synthesis and characterisation of new mixed oxides of antimony and tellurium, 2225–30
TEMPLATE
The template synthesis and crystal and molecular structure of a seven-co-ordinate manganese(II) complex with 2,6-diacetylpyridine mono(2-aminobenzoylhydrazone), 215–8
The preparation and co-ordination chemistry of 2,2′:6′,2″-terpyridine macrocycles. Part 4. Structural characterisation of an intermediate in a transient template reaction, 333–6
Reactions of tris[(2*S*)-2-(aminomethyl)pyrrolidine]nickel(II) ion with alk-3-en-2-ones or 4-hydroxyalkan-2-ones: formation of an optically active tetra-aza macrocycle, 2139–44
Interactions of $[Mo(CO)_6]$ and $[Mo(CO)_3(C_6H_5CH_3)]$ with O_2N_2- and O_2N_3-donor macrocycles and the *X*-ray crystal structure of tetracarbonyl(6,7,16,17-tetrahydro-15*H*-dibenzo[*e*,*n*][1,4,8,12]dioxadiazacyclopentadecine)molybdenum–toluene (2/1), 2561–4
TERPYRIDINE
The preparation and co-ordination chemistry of 2,2′:6′,2″-terpyridine macrocycles. Part 4. Structural characterisation of an intermediate in a transient template reaction, 333–6
Ligand reactivity in polypyridine complexes; the deuteriation of the bis(2,2′:6′,2″-terpyridine)ruthenium(II) cation, 2687–90
TETRA-AZACYCLODODECANE
Proton exchange and base hydrolysis of *syn,anti-cis*-dichloro(1,4,7,10-tetra-azacyclododecane)cobalt(III) cations, 475–8

TETRA-AZACYCLODODECANE (contd)
The stability of nickel(II) complexes of tetra-aza macrocycles, 1877–80
TETRA-AZACYCLOHEXADECANE
Studies of pendant-arm macrocyclic ligands. Part 4. Two penta-aza macrocycles based on 1-(2′-dimethylaminoethyl)-1,5,9,13-tetra-azacyclohexadecane and its complexes with bivalent metal ions, 1361–4
TETRA-AZACYCLOTETRADECANE
Preparation of 11-(2′-dimethylaminoethyl)-1,4,7-trimethyl-1,4,7,11-tetra-azacyclotetradecane, and characterisation of its nickel(II), copper(II), and zinc(II) complexes, 219–22
TETRA-AZACYCLOTRIDECANE
The stability of nickel(II) complexes of tetra-aza macrocycles, 1877–80
TETRACHLOROIODATE
Thermochemistry of polyhalides. Part 5. Standard enthalpies of formation of tetramethylammonium and tetraethylammonium tetrachloroiodates, 997–1000
TETRADECAVANADOPHOSPHATE
Oxygen exchange and protonation of polyanions: a multinuclear magnetic resonance study of tetradecavanadophosphate(9−) and decavanadate(6−), 1953–8
TETRAHYDROALUMINATE
Alkyl, hydrido, and tetrahydroaluminato complexes of manganese with 1,2-bis(dimethylphosphino)ethane (dmpe). *X*-Ray crystal structures of $Mn_2(\mu\text{-}C_6H_{11})_2(C_6H_{11})_2(\mu\text{-dmpe})$, $(dmpe)_2Mn(\mu\text{-}H)_2AlH(\mu\text{-}H)_2AlH(\mu\text{-}H)_2Mn(dmpe)_2$, and $Li_4\{MnH(C_2H_4)[CH_2(Me)PCH_2CH_2PMe_2]_2\}_2\cdot 2Et_2O$, 921–30
TETRAHYDROBORATE
Synthesis, characterization, and structure of the complex $[FeH(H_2BH_2)\{CH_3C(CH_2PPh_2)_3\}]$, 605–10
Hydrogen generation by hydrolysis of sodium tetrahydroborate: effects of acids and transition metals and their salts, 307–14
TETRAHYDROFURAN
Tetrachlorobis(tetrahydrofuran)molybdate(III), a new intermediate for the synthesis of molybdenum(III) complexes: preparation and crystal structure, 1069–72
Oxoalkyls of rhenium-(V) and -(VI). *X*-Ray crystal structures of $(Me_4ReO)_2Mg(thf)_4$, $[(Me_3SiCH_2)_4ReO]_2Mg(thf)_2$, $Re_2O_3Me_6$, and $Re_2O_3(CH_2SiMe_3)_6$, 2167–76
TETRAHYDROTHIOPHENE
Sulphur ligand–metal complexes. Part 16. Copper complexes of thioethers and the single-crystal *X*-ray structure of the polymeric mixed-valence complex, penta-μ-chloro-tris-μ-tetrahydrothiophene-tetracopper(I,II), 151–8
Niobium(IV) sulphidohalides: preparation of $Nb_2X_4S_3$ and $Nb_2X_4S\cdot nL$[X = Br or Cl; n = 4, L = NCMe, SMe_2, or tetrahydrothiophene (tht); n = 2, L = $PhSCH_2CH_2SPh$]. Crystal and molecular structure of $Nb_2Cl_4S_2\cdot 4tht$, 417–22
TETRAKIS(METHYLTHIO)ETHANE
Pyramidal inversion in configurational isomers of tetracarbonyl[1,1,2,2-tetrakis(methylthio)ethane]chromium(0): a two-dimensional nuclear magnetic resonance exchange study, 2195–202
TETRAPHOSPHADITHIACYCLODOCOSANE
Macrocyclic polyphosphane ligands. Cobalt(II) and nickel(II) complexes of 5*RS*,8*RS*,16*RS*,19*RS*-tetraphenyl-5,8,16,19-tetraphospha-1,12-dithiacyclododecosane (δ-L^1) and the crystal structure of $[Co(\delta\text{-}L^1)][BF_4]_2\cdot 0.5H_2O$, 1179–82
TETRAPHOSPHADITHIACYCLO-OCTADECANE
Macrocyclic polyphosphane ligands. Cobalt(II) and nickel(II) complexes of the γ and ε diastereoisomers of 4,7,13,16-tetraphenyl-4,7,13,16-tetraphospha-1,10-dithiacyclo-octadecane (L^1) and the crystal structure of $[Co(\varepsilon\text{-}L^1)][BPh_4]_2\cdot$ EtOH, 1425–30
TETRATHIAFULVALENE
X-Ray crystal structures and properties of tris(tetrathiafulvalenium) tetrachlorodimethylstannate(IV) and tetrathiafulvalenium trichlorodimethylstannate (IV), 465–70
THERMODYNAMICS
Zinc–sulphur bond enthalpy: its determination in bis(diethyldithiocarbamato)zinc(II), 369–72
Thermochemistry of polyhalides. Part 5. Standard enthalpies of formation of tetramethylammonium and tetraethylammonium tetrachloroiodates, 997–1000
Thermochemical data for adducts of zinc, cadmium, and mercury halides with hexamethylphosphoramide, 1103–6
Solution chemistry of macrocycles. Part 4. Thermodynamics of protonation and complexation of several N_2S_2 macrocycles, 1169–72
Thermodynamics of the actinoid elements. Part 6. The preparation and heats of formation of some sodium uranates(VI), 1853–8
Thermodynamic treatment of exchange of H^+ and K^+ in α-titanium phosphate, 1865–8
On the possibility of determining the thermodynamic parameters for the formation of weak complexes using a simple model for the dependence on ionic strength of activity coefficients: Na^+, K^+, and Ca^{2+} complexes of low molecular weight ligands in aqueous solution, 2353–62
THERMOLYSIS
A kinetic study of the gas-phase thermolysis of hexaborane(10), 541–8
THIADECABORANE
Electronic structures of the *closo*-thiaboranes 1-SB_9H_9 and 1-$SB_{11}H_{11}$. Electrophilic substitution in 1-SB_9H_9, 2591–4
THIADODECABORANE
Electronic structures of the *closo*-thiaboranes 1-SB_9H_9 and 1-$SB_{11}H_{11}$. Electrophilic substitution in 1-SB_9H_9, 2591–4
THIOCARBONYL
Fragmentation of co-ordinated carbon disulphide and dithiocarbene ligands by nucleophilic attack at carbon: the crystal structure of $[(Ph_3P)IPt(\mu\text{-}SMe)(\mu\text{-}CSMe)PtI(PPh_3)]\cdot Me_2CO$, 2595–602
THIOCYANATE
The red form of bis(1,10-phenanthroline)dithiocyanatoiron(II), 991–6
Characterization of the adducts formed by Cu(CN) and Cu(NCS) with biquinoline. The crystal structure of the polymeric cyano-compound containing both linear and tetrahedrally co-ordinated copper(I), $[\{Cu_3(bq)_2(CN)_3\}_n]$, 1285–8
THIOETHER
Synthesis and properties of dioxo-osmium(VI) compounds of thio- and seleno-ethers, 205–8
Reactions of sulphoxide–thioether bidentate ligands with platinum(II). Determination of the mode of binding by cyclic voltammetry, 209–12
Static and dynamic nuclear magnetic resonance studies of complexes of trimethylplatinum(IV) halides with olefinic thio-and seleno-ethers. *X*-Ray crystal structures of $[PtXMe_3(MeSeCH{=}CHSeMe)]$ (X = Cl or I), 345–54
Copper complexes with quadridentate bis(pyrazolyl)thioether amine and tris(pyrazolyl)amine ligands. Structural characterization of the complexes $[Cu(NCS)(tpea)][Cu(NCS)_2]$ and $[CuCl(bdma)]Cl\cdot 2H_2O$, 2327–32
THIONITROSYL
Reaction of trithiazyl trichloride, $(NSCl)_3$, with triphenylphosphine or triphenylphosphine metal complexes. *X*-Ray crystal structure of aminotriphenylphosphonium chloride–dichloromethane (1/1), $[Ph_3PNH_2]Cl\cdot CH_2Cl_2$, 1043–8
THIONITROSYL CHLORIDE
Thiazyl chloride complexes of ruthenium(II), 2449–52
THIOPHENOLATE
Rhenium nitrosyl complexes with simple and with sterically demanding aromatic thiolate ligands: *X*-ray crystal structures of $[PPh_4][Re_2(SC_6H_4Me\text{-}4)_7(NO)_2]\cdot CH_2Cl_2$ and $[Re(SC_6H_3Pr^i_2\text{-}2,6)_4(NO)]$, 1533–42
THIOPYRIDINE OXIDE
Comparison between the bis(2-thiopyridine *N*-oxide) derivatives of Cu^{II} and Ni^{II}: an electron spin resonance study, 379–82
THIOUREA
Oxidation of thiourea by iodate: a new type of oligo-oscillatory reaction, 1669–72
TIN
Subvalent group 4B metal alkyls and amides. Part 7. Transition-metal chemistry of metal(II) bis(trimethylsilyl)amides $M'(NR_2)_2$ (R = $SiMe_3$; M′ = Ge, Sn, or Pb), 51–8
Tin-119 Mössbauer and nuclear magnetic resonance studies of organotin compounds. Part 1. Sterically crowded tetraorganotin derivatives, 169–76
Intercalation of alkylamines into tin(IV) bis(hydrogenphosphate) monohydrate, 213–4
Chemical and structural aspects of silver–triphenylarsine complexes and silver–tin complex salts, 321–32
The preparation and co-ordination chemistry of 2,2′:6′,2″-terpyridine macrocycles. Part 4. Structural characterisation of an intermediate in a transient template reaction, 333–6
Tin–molybdenum oxides. A study by *X*-ray diffraction, Mössbauer spectroscopy, and electron spin resonance, 451–4

TIN (contd)
X-Ray crystal structures and properties of tris(tetrathiafulvalenium) tetrachlorodimethylstannate(IV) and tetrathiafulvalenium trichlorodimethylstannate (IV), 465–70
Investigation into diphosphine oxides as ligands in diorganotin(IV) adducts. Part 3. Synthesis and crystal structure of two adducts of dinitratodiphenyltin(IV) with *cis*- and *trans*-1,2-bis(diphenylphosphoryl)ethylene, 487–92
Studies on the anti-tumour activity of di- and tri-organotin(IV) complexes of amino acids and related compounds, of 2-mercaptoethanesulphonate, and of purine-6-thiol, 523–8
The crystal structure of diaquadichlorodimethyltin(IV)–purine (1/4), 1271–4
A tin-119 Mössbauer study of tin(II) fluoride, 1275–6
Mössbauer study of the *cis–trans* isomers of tin(IV) complexes. Some considerations about the sign of the electric-field gradient, 1281–4
Bromine nuclear quadrupole resonance studies of some hexabromostannates: *X*-Ray crystal structure of pyridinium hexabromostannate(IV), $[Hpy]_2[SnBr_6]$, 1399–404
Organotin biocides. Part 2. Variable-temperature ^{119}Sn Mössbauer study of phenyl-and cyclohexyl-tin compounds, 1417–24
Methyltin hydrides in fluorosulphuric acid, 2671–6
A novel pentameric hydrolysis product of $SnMe_2Cl_2$: crystal and molecular structure of $[NHEt_3][(SnMe_2Cl)_5O_3]$, 2683–6

TIN-119
Tin-119 Mössbauer and nuclear magnetic resonance studies of organotin compounds. Part 1. Sterically crowded tetraorganotin derivatives, 169–76
Methyltin hydrides in fluorosulphuric acid, 2671–6

TITANIUM
Coloured species formed from the titanium(IV)-4-(2′-pyridylazo)resorcinol reagent in the spectrophotometric determination of trace amounts of hydrogen peroxide, 81–4
Mono-η-cycloheptatrienyltitanium chemistry: synthesis, molecular and electronic structures, and reactivity of the complexes $[Ti(\eta\text{-}C_7H_7)L_2X]$ (L = tertiary phosphine, O- or N-donor ligand; X = Cl or alkyl), 669–84
Thermodynamic treatment of exchange of H^+ and K^+ in α-titanium phosphate, 1865–8
Chemistry of di- and tri-metal complexes with bridging carbene or carbyne ligands. Part 37. Methylene group transfer to carbon–metal multiple bonds; crystal structures of $[TiW\{\mu\text{-}C(C_6H_4Me\text{-}4)=CH_2\}(\mu\text{-}CO)(CO)(\eta\text{-}C_5H_5)_3]$ and $[PtW\{\mu\text{-}C(C_6H_4Me\text{-}4)=CH_2\}(CO)_2(PMe_3)_2(\eta\text{-}C_5H_5)]$, 2009–16
Synthesis, structure, and bonding of fulvene complexes of titanium, molybdenum, and tungsten, 2037–50

TOLUENEDITHIOLATE
Toluene-3,4-dithiol (H_2tdt) complexes of Group 5B halides. Observations of lone-pair stereochemical activity and redox behaviour. Crystal and molecular structures of [AsCl(tdt)] and $[PPh_4][Sb(tdt)_3]$, 69–74

TOLYLETHENYL
Chemistry of di- and tri-metal complexes with bridging carbene or carbyne ligands. Part 37. Methylene group transfer to carbon–metal multiple bonds; crystal structures of $[TiW\{\mu\text{-}C(C_6H_4Me\text{-}4)=CH_2\}(\mu\text{-}CO)(CO)(\eta\text{-}C_5H_5)_3]$ and $[PtW\{\mu\text{-}C(C_6H_4Me\text{-}4)=CH_2\}(CO)_2(PMe_3)_2(\eta\text{-}C_5H_5)]$, 2009–16

TOLYLMETHYLENE
Chemistry of di- and tri-metal complexes with bridging carbene or carbyne ligands. Part 35. The synthesis and crystal structures of the compounds $[N(PPh_3)_2][W_2(\mu\text{-}CH)C_6H_4Me\text{-}4\}(CO)_7(\eta\text{-}C_5H_5)]\cdot 0.5Et_2O$ and $[W\{=C(H)C_6H_4Me\text{-}4\}(SnPh_3)(CO)_2(\eta\text{-}C_5H_5)]$, 1331–8

TOLYLMETHYLIDYNE
Chemistry of di- and tri-metal complexes with bridging carbene or carbyne ligands. Part 36. Reactions of the dimetal compounds $[ReM(\equiv CC_6H_4Me\text{-}4)(CO)_9]$ (M = Cr, Mo, or W) with octacarbonyldicobalt; crystal structures of $[Co_2WRe(\mu_3\text{-}CC_6H_4Me\text{-}4)(CO)_{15}]$ and $[Co_2Re(\mu_3\text{-}CC_6H_4Me\text{-}4)(CO)_{10}]$, 2001–8

TRANSIENT
Equilibrium and kinetic study of the reaction between phthalocyaninatoiron(II) and carbon monoxide in dimethyl sulphoxide in the presence of pyridine. Evidence for the formation of a transient, 1113–8

TRANSITION METAL
Synthesis and properties of the divalent 1,2-bis(dimethylphosphino)ethane (dmpe) complexes $MCl_2(dmpe)_2$ and $MMe_2(dmpe)_2$ (M = Ti, V, Cr, Mn, or Fe). *X*-Ray crystal structures of $MCl_2(dmpe)_2$ (M = Ti, V, or Cr), $MnBr_2(dmpe)_2$, $TiMe_{1.3}Cl_{0.7}(dmpe)_2$, and $CrMe_2(dmpe)_2$, 1339–48
Studies of pendant-arm macrocyclic ligands. Part 4. Two penta-aza macrocycles based on 1-(2′-dimethylaminoethyl)-1,5,9,13-tetra-azacyclohexadecane and its complexes with bivalent metal ions, 1361–4

TRICYCLOUNDECADIENYL
Annelation of ring-opened arylcyclopropenium ions to co-ordinated cyclo-octatetraene, and the *X*-ray crystal structure of $[Fe(CO)_3(\sigma,\eta^3\text{-}C_{11}H_9Ph_3)]$, 777–82

TRIETHYLAMINE
Ligand substitution on (*N*,*N*-dimethylformamide)[2,2′,2″-tri(*N*,*N*-dimethylamino)triethylamine]cobalt(II), 413–6

TRIETHYLENETETRAMINE
Carbon-13 nuclear magnetic resonance study of the complexes formed between zinc(II) and triethylenetetramine, 2381–6

TRIFLUOROACETATE
Complexes of the platinum metals. Part 23. Synthesis of the nitrosyl carboxylate complexes $[M(O_2CR)_2(NO)(PPh_3)_2]$ (M = Rh or Ir; R = CF_3, C_2F_5, or C_6F_5): crystal and molecular structures of the trifluoroacetate derivatives $[M(O_2CCF_3)_2(NO)(PPh_3)_2]$, 611–6
Complexes of the platinum metals. Part 25. Reactions of ruthenium and osmium nitrosyls $[MH(NO)(PPh_3)_3]$ and $[M(NO)_2(PPh_3)_2]$ with perfluorocarboxylic acids: *X*-ray crystal structure determination of nitrosyl(trifluoroacetato)-(trifluoroacetohydroximato-*OO*′)bis(triphenylphosphine)osmium(II)–dichloromethane (1/1), 621–8
Complexes of lanthanoid salts with macrocyclic ligands. Part 17. Synthesis and crystal and molecular structure of a hydroxide-bridged praseodymium trifluoroacetate complex with 15-crown-5-ether, $[Pr_2(CF_3CO_2)_3(OH)(C_{10}H_{20}O_5)_2]\cdot[Pr_2(CF_3CO_2)_6]$, 885–90

TRIFLUOROACETIC ACID
Complexes of the platinum metals. Part 24. The role of dioxygen in the reactions of trifluoroacetic acid with the rhodium and iridium nitrosyls $[M(NO)(PPh_3)_3]$, 617–20

TRIFLUOROACETOHYDROXIMATE
Complexes of the platinum metals. Part 25. Reactions of ruthenium and osmium nitrosyls $[MH(NO)(PPh_3)_3]$ and $[M(NO)_2(PPh_3)_2]$ with perfluorocarboxylic acids: *X*-ray crystal structure determination of nitrosyl(trifluoroacetato)-(trifluoroacetohydroximato-*OO*′)bis(triphenylphosphine)osmium(II)–dichloromethane (1/1), 621–8

TRIFLUOROMETHYL
The molecular structure of gaseous bis(trifluoromethyl)selenium difluoride as determined by electron diffraction, 941–6
A disagreement on the explanation of short and long As–O bonds of the $(As\text{–}O)_4$ ring in $As_4(CF_3)_6O_6(OH)_2$ in terms of As^{III}–O and As^{V}–O bonds, 2221–2

TRIKETONE
Reactions of metal ions with triketones in solution. Part 1. Formation constants for the systems of heptane-2,4,6-trione, 1-phenylhexane-1,3,5-trione, 1,5-diphenylpentane-1,3,5-trione, and 2,2′-dihydroxybenzophenone with proton, nickel(II), and cobalt(II), 1543–6

TRIMETHYLSILYL
Subvalent group 4B metal alkyls and amides. Part 7. Transition-metal chemistry of metal(II) bis(trimethylsilyl)amides $M'(NR_2)_2$ (R = $SiMe_3$; M′ = Ge, Sn, or Pb), 51–8
Synthesis of α-lithioarylmethanes of *m*-xylene and its α-trimethylsilyl derivatives; crystal structure of $[\{Li(Me_2NCH_2CH_2NMe_2)\}_2\{C_6H_4(CHSiMe_3)_2\text{-}m\}]$, 337–44
Purification and *X*-ray crystal structure of bis[tris(trimethylsilyl)methyl]diarsene, 383–6

TRIMETHYLSILYLMETHYL
Oxoalkyls of rhenium-(V) and -(VI). *X*-Ray crystal structures of $(Me_4ReO)_2Mg(thf)_4$, $[(Me_3SiCH_2)_4ReO]_2Mg(thf)_2$, $Re_2O_3Me_6$, and $Re_2O_3(CH_2SiMe_3)_6$, 2167–76

TRINITROBENZENE
Synthesis and characterization of a new family of binuclear copper(II) complexes with a flexible bridge and the π–π type charge-transfer interaction with 1,3,5-trinitrobenzene, 107–10

TRIONE
Chemistry of metallacyclobutanones. Part 3. Reactions of heptane-2,4,6-trione and 1,5-diphenylpentane-1,3,5-trione with some

TRIONE (contd)
carbonate complexes of platinum(II); *X*-ray crystal structures of $[Pt\{CH(COMe)COCH(COMe)\}(PPh_3)_2]$ and $[Pt\{OC(CHCOPh)CHC(Ph)O\}(PPh_3)_2]$, 549–54
TRIPHENYLSTANNYL
Chemistry of di- and tri-metal complexes with bridging carbene or carbyne ligands. Part 35. The synthesis and crystal structures of the compounds $[N(PPh_3)_2][W_2\{\mu\text{-}C(H)C_6H_4Me\text{-}4\}(CO)_7(\eta\text{-}C_5H_5)]\cdot 0.5Et_2O$ and $[W\{=C(H)C_6H_4Me\text{-}4\}(SnPh_3)(CO)_2(\eta\text{-}C_5H_5)]$, 1331–8
TRISELENOCARBONATE
Sulphur-containing metal complexes. Part 14. Reactions of carbene anions with carbon disulphide or carbon diselenide, 1963–6
TRITHIATETRAPHOSPHABICYCLOHEPTANE
Intermediates in the photochemical reaction of tetraphosphorus trisulphide with organic disulphides; phosphorus-31 nuclear magnetic resonance parameters for 2,6-bis(alkylthio)- and 2,6-diiodo-3,5,7-trithia-1,2,4,6-tetraphosphabicyclo[2.2.1]heptanes and for 3,6-bis(alkylthio)-2,5,7-trithia-1,3,4,6-tetraphosphabicyclo[2.2.1]heptanes, 1707–12
TRITHIATETRAPHOSPHABICYCLOHEPTYL
Phosphorus–phosphorus bond cleavage in the cage molecule P_4S_3: synthesis and crystal structure of the trinuclear platinum complex $[\{Pt(\mu\text{-}P_4S_3)(PPh_3)\}_3]\cdot C_6H_6$, 291–6
TRITHIOLANE
Reaction of bis(morpholinothiocarbonyl) disulphide with iodine. Existence of a 1:1 charge-transfer precursory adduct in an oxidation reaction. Isolation and crystal structure of bis[3,5-di(*N*-morpholinio)-1,2,4-trithiolane] hexadecaiodide, 1349–54
TUNGSTEN
Reaction of bis-μ-diethylphosphido-bis(tetracarbonylmetal) (*M*–*M*) (M = Cr or W) with tri-n-butylphosphine: kinetics and mechanism of a reaction involving seven-co-ordinate complexes, 91–8
Heteropolytungstates as catalysts for the photochemical reduction of oxygen and water, 395–400
Trimethylphosphine polyhydrides of tungsten and rhenium, 587–90
Formation of substituted cyclopentadienyl ligands on tungsten *via* reactions between the alkyne complexes $[W(CO)(R^1C_2R^2)_3]$ ($R^1 = R^2$ = Ph or Et; R^1 = Me, R^2 = Ph) and the alkylidyne compounds $[W(\equiv CR)(CO)_2(\eta\text{-}C_5H_5)]$ (R = C_6H_4Me-4 or Me); *X*-ray crystal structures of $[W_2(\mu\text{-}CO)_2(CO)(\eta\text{-}PhC_2Ph)(\eta\text{-}C_5Ph_4R)(\eta\text{-}C_5H_5)]$ and $[W_2(\mu\text{-}EtC_2Et)(CO)_4(\eta\text{-}C_5Et_4R)(\eta\text{-}C_5H_5)]$ (R = C_6H_4Me-4), 905–12
Reaction of transition-metal carbonylate anions and 1,1,1-tris(halogenomethyl)ethane. *X*-Ray crystal structures of tricarbonyl(η⁵-cyclopentadienyl)(1-methylcyclopropylmethyl)tungsten(II), and tetraethylammonium enneacarbonyliododirhenate(0), 931–40
Bimetallic systems. Part 9. The synthesis of and nuclear magnetic resonance studies on 10-membered ring complexes of type $[(OC)_4M^1(\mu\text{-}Ph_2PCH_2CH_2PPh_2)_2M^2(CO)_4]$ (M^1, M^2 = Cr, Mo, or W), 1009–14
Isolated CH stretching frequencies, methyl group geometry, and methyl CH bond lengths and strengths in tricarbonyl(η³-cyclopentadienyl)methylchromium(II), -molybdenum(II), and -tungsten(II), 1207–12
Chemistry of di- and tri-metal complexes with bridging carbene or carbyne ligands. Part 33. Reactions of $[W(\equiv CMe)(CO)_2(\eta\text{-}C_5H_5)]$ with the dimetal compounds $[MRh(\mu\text{-}CO)_2(\eta\text{-}C_5Me_5)_2]$ (M = Co or Rh); *X*-ray crystal structure of $[Rh_2W(\mu\text{-}CO)(\mu_3\text{-}CMe)(CO)_2(\eta\text{-}C_5H_5)(\eta\text{-}C_5Me_5)_2]$, 1315–22
Chemistry of di- and tri-metal complexes with bridging carbene or carbyne ligands. Part 34. Reactions of sulphur and selenium with the di-irontungsten complexes $[Fe_2W(\mu_3\text{-}CR)(\mu\text{-}CO)(CO)_8(\eta\text{-}C_5H_5)]$ (R = C_6H_4Me-4 or Me); crystal structures of $[Fe_2W(\mu\text{-}CC_6H_4Me\text{-}4)(\mu_3\text{-}S)(CO)_7(\eta\text{-}C_5H_5)]$ and $[Fe_2W(\mu_3\text{-}SCMe)(CO)_8(\eta\text{-}C_5H_5)]$, 1323–30
Chemistry of di- and tri-metal complexes with bridging carbene or carbyne ligands. Part 35. The synthesis and crystal structures of the compounds $[N(PPh_3)_2][W_2\{\mu\text{-}C(H)C_6H_4Me\text{-}4\}(CO)_7(\eta\text{-}C_5H_5)]\cdot 0.5Et_2O$ and $[W\{=C(H)C_6H_4Me\text{-}4\}(SnPh_3)(CO)_2(\eta\text{-}C_5H_5)]$, 1331–8
Preparation and properties of *cis*-$[W(N_2)_2(Ph_2PCH_2PPh_2)_2]$ and *trans*-$[M(N_2)_2(Ph_2PCH{=}CHPPh_2)_2]$ (M = Mo or W), and the crystal structure of *cis*-$[W(N_2)_2(PMe_2Ph)_4]$, 1523–6
Pyramidal inversions and 1,2-metal shifts in pentacarbonyl-chromium, -molybdenum, and -tungsten derivatives of dialkyl disulphides and dialkyl diselenides. A nuclear magnetic resonance investigation, 1561–8
Dynamic nuclear magnetic resonance study of Group 6 metal pentacarbonyl complexes. Pyramidal inversion and 1,2-metal shifts in the complexes $[M(CO)_5(Me_2CCH_2EECH_2)]$ (M = Cr, Mo, or W; E = S or Se), 1569–76
The kinetics and stoicheiometry of silver(III) reduction by the octacyano-complexes of molybdenum(IV) and tungsten(IV), 1789–94
Chemistry of di- and tri-metal complexes with bridging carbene or carbyne ligands. Part 36. Reactions of the dimetal compounds $[ReM(\equiv CC_6H_4Me\text{-}4)(CO)_9]$ (M = Cr, Mo, or W) with octacarbonyldicobalt; crystal structures of $[Co_2WRe(\mu_3\text{-}CC_6H_4Me\text{-}4)(CO)_{15}]$ and $[Co_2Re(\mu_3\text{-}CC_6H_4Me\text{-}4)(CO)_{10}]$, 2001–8
Chemistry of di- and tri-metal complexes with bridging carbene or carbyne ligands. Part 37. Methylene group transfer to carbon–metal multiple bonds; crystal structures of $[TiW\{\mu\text{-}C(C_6H_4Me\text{-}4)=CH_2\}(\mu\text{-}CO)(CO)(\eta\text{-}C_5H_5)_3]$ and $[PtW\{\mu\text{-}C(C_6H_4Me\text{-}4)=CH_2\}(CO)_2(PMe_3)_2(\eta\text{-}C_5H_5)]$, 2009–16
Chemistry of di- and tri-metal complexes with bridging carbene or carbyne ligands. Part 38. Ruthenium–tungsten compounds: crystal structures of $[RuW(\mu\text{-}Cl)(\mu\text{-}CMe)(Cl)(CO)_2(PPh_3)_2(\eta\text{-}C_5H_5)]$ and $[RuW_2(\mu_3\text{-}C_2Me_2)(CO)_7(\eta\text{-}C_5H_5)_2]$, 2017–24
Trimethylphosphine as a reactive solvent: synthesis, crystal structures, and reactions of $[Ta(PMe_3)_3(\eta^2\text{-}CH_2PMe_2)(\eta^2\text{-}CHPMe_2)]$ and $[W(PMe_3)_4(\eta^2\text{-}CH_2PMe_2)H]$ and related studies, 2025–36
Synthesis, structure, and bonding of fulvene complexes of titanium, molybdenum, and tungsten, 2037–50
Preparation and crystal structure of $[AsPh_4]_2[(WCl_5)_2\{\mu\text{-}NC(CF_3)_2N\}]$, 2205–8
Co-ordinatively unsaturated diene complexes of tungsten(II) and their reactions with nucleophiles to give six- and seven-co-ordinate derivatives, 2231–8
Synthesis and nuclear magnetic resonance studies of co-ordinatively unsaturated alkyne complexes of tungsten(II), 2239–46
Reaction of the anion $[WCl_4(CBu^t)]^-$ with tetrasulphur tetranitride. Formation and crystal structure of $[AsPh_4][WCl_3O(OS_2N_2)]$, 2453–6
Nitrogen-15 nuclear magnetic resonance relaxation mechanisms in dinitrogen complexes of molybdenum, tungsten, rhenium, and osmium, 2473–8
Complexes of molybdenum-(II) and -(IV) and tungsten(II) with sterically hindered thiolate ligands. Synthesis, reactivity, and *X*-ray crystal structures of $[PPh_4][Mo(SC_6H_2Pr^i_3\text{-}2,4,6)_3(CO)_2]$ and $[Mo(NNPh)(SC_6H_2Pr^i_3\text{-}2,4,6)_3(NCMe)]$, 2639–46
Molybdenum, rhenium, and tungsten complexes with bi- and tridentate phosphinothiolato-ligands; structures of $[Mo\{PhP(CH_2CH_2S)_2\}_2]$ and $[Mo(NNMe_2)\{PhP(CH_2CH_2S)_2\}_2]$, 2647–54
TYROSINE
Specific binding of the tyrosine residue in copper(II) complexes of Tyr-Pro-Gly-Tyr and Tyr-Gly-Pro-Tyr, 1201–6
URANIUM
Synthesis and structural assessment of ammonium and caesium difluorodioxoperoxouranates(VI), $A_2[UO_2(O_2)F_2]$ (A = NH_4 or Cs), and alkali-metal difluorodioxoperoxouranate(VI) monohydrates, $A_2[UO_2(O_2)F_2]\cdot H_2O$ (A = K or Rb), 409–12
Glycine complexation with uranyl ion: absorptiometric, luminescence, and *X*-ray structural studies of tetrakis-(glycine)dioxouranium(VI) nitrate, 517–22
Solvent effects on the redox potential of the uranium(VI)–uranium (V) couple, 601–4
Kinetics and mechanism of single electron oxidations of the tervalent uranium ion, U^{3+}(aq), by free radicals in aqueous solutions, 641–4
Actinide structural studies. Part 7. The crystal and molecular structures of (2,2′-bipyridyl)dinitratodioxo-uranium(VI) and -neptunium(VI), and diacetato(2,2′-bipyridyl)dioxo-uranium(VI) and -neptunium(VI), 1001–8
Thermodynamics of the actinoid elements. Part 6. The preparation and heats of formation of some sodium uranates(VI), 1853–8
Synthesis and characterisation of uranium(IV) nitrate complexes with piperazines, 1985–8
URIDINE TRIPHOSPHATE
Influence of decreasing solvent polarity (dioxane–water mixtures) on the stability and structure of binary and ternary complexes of adenosine 5′-triphosphate and uridine 5′-triphosphate, 2291–304

VANADIUM
High-field vanadium-51 and oxygen-17 nuclear magnetic resonance study of peroxovanadates(v), 1173–8
Reactions of bis(cyclopentadienyl)vanadium derivatives with nitrogen mono-oxide and the crystal structure of an oxo-bridged nitrosyl complex of vanadium, 1435–42
Axially asymmetric metal alkyls. Part 3. Chemical, electrochemical, and structural studies of group 5A $d^{0,1,2}$ metallepines $[M\{(2\text{-}CH_2C_6H_4)_2\}(\eta\text{-}C_5H_5)_2]^z$ (M = Nb or Ta; $z = 1-$, 0, or 1+); synthesis of $[Nb\{2\text{-}CH_2(C_6H_4)_2Me\text{-}2'\}X(\eta\text{-}C_5H_5)_2]$ (X = Cl or CO) and $[\{(\eta\text{-}C_5H_5)_2ClV\}_2\{(2\text{-}CH_2C_6H_4)_2\}]$, and electrochemistry of $[M(CH_2Ph)_{2-n}Cl_n(\eta\text{-}C_5H_5)_2]$ (n = 0, M = Nb, or Ta; n = 1, M = V), 1747–60
Oxygen exchange and protonation of polyanions: a multinuclear magnetic resonance study of tetradecavanadophosphate(9−) and decavanadate(6−), 1953–8
A kinetic study of the complexation of cysteine and related compounds with aqueous vanadium(II) and vanadium(III) at approximately neutral pH; the mediating role of sulphur compounds in electron transfer, 2461–8
Kinetics and mechanism of the equilibration reaction between (2,2′,2″-nitrilotriethoxy)nitrosylvanadate(1−) and cyanide. Crystal structures of sodium (2,2′,2″-nitrilotriethoxy)nitrosylvanadate(I)–sodium perchlorate tetrahydrate and of barium cyano(2,2′,2″-nitrilotriethoxy)nitrosylvanadate(I) pentahydrate, 2493–8
VANADIUM-51
High-field vanadium-51 and oxygen-17 nuclear magnetic resonance study of peroxovanadates(v), 1173–8
VINYL
Reactions of co-ordinated ligands. Part 33. Mononuclear η^2-vinyl complexes: synthesis, structure, and reactivity, 435–50
Organoruthenium(II) complexes formed by insertion reactions of some vinyl compounds and conjugated dienes into a hydrido–ruthenium bond, 873–8
VINYLOXY
Acid-induced displacement of acetaldehyde from a μ-vinyloxy-triosmium cluster, 2479–82
VITAMIN B_{12}
The chemistry of vitamin B_{12}. Part 25. Mechanism of the β-elimination of olefins from alkylcorrinoids; evidence for an initial homolytic fission of the Co–C bond, 1613–8
The chemistry of vitamin B_{12}. Part 26. Co-ordination of the malonitrile anion by Co^{III} corrinoids: first experimental determination of equilibrium constants for the co-ordination of a tetrahedral carbanion by a transition-metal ion, 1619–22
VITAMIN B_6
Metal complexes of vitamin B_6 related compounds. Crystal and molecular structures of aqua(5′-phosphopyridoxylideneglycinato)copper(II) trihydrate and bis(pyridoxylideneglycinato)nickel(II) hexahydrate, 2051–8
WATER
A kinetic study of the complexation of cysteine and related compounds with aqueous vanadium(II) and vanadium(III) at approximately neutral pH; the mediating role of sulphur compounds in electron transfer, 2461–8
XANTHOSINE
Influence of secondary ligands on the stability of metal–xanthosine complexes in solution, 239–42
***X*-RAY DIFFRACTION**
Tin–molybdenum oxides. A study by *X*-ray diffraction, Mössbauer spectroscopy, and electron spin resonance, 451–4
X-Ray diffraction determination of the cation distribution and oxygen positional parameter in polycrystalline spinels, 2155–60
Preparation, characterization, and physical properties of the series MPd_3S_4 (M = rare earth), 2369–74
XYLENE
Synthesis of α-lithioarylmethanes of *m*-xylene and its α-trimethylsilyl derivatives; crystal structure of $[\{Li(Me_2NCH_2CH_2NMe_2)\}_2\{C_6H_4(CHSiMe_3)_2\text{-}m\}]$, 337–44
YLIDE
Synthesis of gold-(I) and -(III) complexes with carbonyl-stabilized phosphorus ylides. Crystal structure of $[\{Au(PPh_3)\}_2\{\mu\text{-}C(PPh_3)CO_2Et\}]ClO_4$, 1163–8
ZINC
The crystal structure and electronic properties of the complex acetatobis(1,10-phenanthroline)copper(II) perchlorate dihydrate, acetatobis(1,10-phenanthroline)copper(II) nitrate dihydrate, and acetatobis(1,10-phenanthroline)zinc(II) tetrafluoroborate dihydrate, 141–50
Preparation of 11-(2′-dimethylaminoethyl)-1,4,7-trimethyl-1,4,7,11-tetra-azacyclotetradecane, and characterisation of its nickel(II), copper(II), and zinc(II) complexes, 219–22
Zinc–sulphur bond enthalpy: its determination in bis(diethyldithiocarbamato)zinc(II), 369–72
Co-ordination of alkali metals by open-chain polyethers in transition metal complexes. Part 4. Variation in alkali-metal ion selectivity in cobalt and zinc complexes of 1-(*o*-carboxymethoxyphenoxy)-2-(*o*-hydroxyphenoxy)ethane (HL) and the *X*-ray and molecular structure of $[Zn(NH_4L_2)_2]$, 459–64
Thermochemical data for adducts of zinc, cadmium, and mercury halides with hexamethylphosphoramide, 1103–6
Potentiometric study of the complex-formation equilibria of manganese(II), cobalt(II), nickel(II), copper(II), and zinc(II) with ethylenediamine-*N*-acetic acid, 1605–8
Carbon-13 nuclear magnetic resonance study of the complexes formed between zinc(II) and triethylenetetramine, 2381–6
Investigation into aroylhydrazones as chelating agents. Part 7. Synthesis and spectroscopic characterization of complexes of Mn^{II}, Co^{II}, Ni^{II}, Cu^{II}, and Zn^{II} with 2,6-diacetylpyridine bis(2-aminobenzoylhydrazone) and *X*-ray structure of chloro[2,6-diacetylpyridine bis(2-aminobenzoylhydrazone)](methanol)manganese(II) chloride monohydrate, 2387–92
Trigonal bipyramidal penta-aquazinc(II): crystal structure of penta-aquazinc(II) bis(3,3′,3″-phosphinetriyltripropionato)dizincate(II,II) heptahydrate, 2393–6
ZIRCONIUM
Insertion of NO into transition metal–aryl bonds: formation of zirconium complexes containing the $[ON(R)NO]^-$ (R = Ph or p-MeC_6H_4) ligand, 405–8
Crystalline zirconium(IV) hydrogenarsenate hydrogenphosphate monohydrate: synthesis, ion-exchange properties, and thermal behaviour, 1737–42